

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

584 M
NO 269

UNITED STATES DEPARTMENT OF AGRICULTURE
MISCELLANEOUS PUBLICATION NO. 369

WASHINGTON, D. C.

MARCH 1941

THE MINERAL COMPOSITION
OF CROPS
WITH PARTICULAR REFERENCE
TO THE SOILS IN WHICH
THEY WERE GROWN
A REVIEW AND COMPILATION

By
KENNETH C. BEESON
Chemist
Bureau of Plant Industry



UNITED STATES DEPARTMENT OF AGRICULTURE

MISCELLANEOUS PUBLICATION No. 369

Washington, D. C.

March 1941

THE MINERAL COMPOSITION OF CROPS WITH PARTICULAR REFERENCE TO THE SOILS IN WHICH THEY WERE GROWN

A REVIEW AND COMPILATION ¹

By KENNETH C. BEESON, *chemist, Bureau of Plant Industry* ²

CONTENTS

	Page		Page
Introduction.....	1	Factors affecting the mineral composition of plants—Continued.	
Some soil characteristics cited in relation to the occurrence of certain nutritional diseases of man and animals.....	4	Summary of effects of soils on plant composition.....	40
Studies of the relation of soils to bone diseases.....	4	Effect of fertilizers on plant composition.....	41
Studies of the relation of soils to the nutritional anemias.....	12	Nitrogen, phosphorus, and potassium.....	42
Studies of the relation of soils to "lecksucht".....	16	The minor fertilizer elements.....	47
"Alkali disease" or selenium poisoning.....	17	Summary of effect of fertilizers on plant composition.....	51
Iodine.....	17	Miscellaneous factors influencing the mineral composition of plants.....	52
Other nutritional disorders associated with soils.....	19	Age of plant.....	52
Summary of sections on soil characteristics in relation to nutritional diseases of man and animals.....	20	Climate.....	54
Factors affecting the mineral composition of plants.....	22	Irrigation.....	56
Effect of soil composition on mineral composition of plants.....	22	Variety.....	57
Effect of soils on botanical composition of herbage of pastures.....	39	Mineral content of different parts of the plant.....	58
		Literature cited.....	59
		Sources of unpublished material.....	91
		Appendix.....	92
		Tables of crop composition.....	92

INTRODUCTION

The nutritional value of such inorganic elements as calcium, phosphorus, iodine, copper, and iron has been demonstrated by many investigators, and it is generally recognized that quantitative variations of these elements in foods and feeds are important factors in human and animal health. These and other mineral elements, especially those occurring in trace amounts, are receiving greatly increased

¹ The work represented by this publication was supported by the Bankhead-Jones special research fund. It was initiated in the Bureau of Chemistry and Soils (now the Bureau of Agricultural Chemistry and Engineering) and was later transferred to the Bureau of Plant Industry.

² The author wishes to express his appreciation to the several persons who have assisted him. Many helpful suggestions were received from C. A. Browne, under whose direction the work was originally organized, and from Charles E. Kellogg, in charge of the Division of Soil Survey, Bureau of Plant Industry, who originally proposed the study. J. A. LeClerc, of the Bureau of Agricultural Chemistry and Engineering, took an active part in the early stages of the work and contributed much time and effort to it. L. A. Maynard, director of the United States Regional Plant, Soil, and Nutrition Laboratory, and H. G. Byers, in charge of the Division of Soil Chemistry and Physics, both of the Bureau of Plant Industry, made many helpful suggestions and carefully reviewed the manuscript. For her part in the translations of the Italian and Spanish literature, in the systematic handling of the abstracts and cross-references (several thousand such references were involved), in the compilation of the data, and in the careful reading of the manuscript for verbal accuracy, the author is deeply grateful to Ruth B. Rice. The files of the Bureau of Home Economics were very helpful.

attention in both popular and scientific literature dealing with soils, fertilizers, plant growth and composition, food quality, and animal and human health and nutrition.

It is also significant that greater and greater emphasis is being placed upon the problem of the interrelationships among these fields of investigation. It is believed, and the evidence will be cited in this review, that many nutritional diseases are caused by deficiencies or excesses of particular minerals in food plants grown in different soils, in different localities, and with different cultural practices. Although there is abundant evidence, as shown by this literature survey, that these relationships are real and profound, the data are wholly insufficient for defining such relationships in definite practical terms that can be translated into recommendations for agricultural practice on specific soils for the culture of specific food plants. Aside from the fact that the many unsolved aspects of this problem are certainly responsible for much poor nutrition and poor husbandry, this kind of situation is leading to all sorts of conjecture on the part of the public and to many unwarranted claims in regard to the special advantage or disadvantage of certain soils, fertilizers, and food plants. The Department of Agriculture does not have the necessary facts to judge all of these claims.

It is not the purpose of this review to discuss animal nutrition, plant physiology, pedology, or other disciplines closely related to this general problem, as such. It cannot be held to be within the scope of any one worker's experience or capabilities to venture to discuss the special techniques in such a broad variety of fields. Furthermore, within each field, certain facts are well known or appreciated, and it is neither necessary nor desirable to repeat or duplicate the readily available surveys that have already been made. For example, remarkably complete bibliographies on the minor elements and on the relation of mineral deficiencies to animal diseases are available (486, 589, 595).³ A few surveys of the role of minerals in plant or animal nutrition have been made (87, 88, 168, 194, 243, 295, 305, 360, 388, 405, 410, 412, 439, 494, 605), but the discussion in these surveys centers generally around the requirements of plants and animals for minerals. Orr (443) has made a survey of pasture research in the British Empire relating to the composition of herbage and the health of the animal, and brief reviews of the importance of soils in the United States to human and animal nutrition have been made by Maynard (402), Beeson and LeClerc (53), Browne (93), and Auchter (36).

In the investigation of any nutritional disorder in plants or animals, the difficulties of identification of the cause of that disease multiply as the mobility of the subject increases. Thus, a plant, being stationary as far as its orbit of activity is concerned, represents a relatively simple subject for investigation. Any discrepancies or abnormalities in its growth or appearance can be readily discerned and checked against its source of nutrients, the soil. As a result, the effects of more than 40 elements on the growth and health of plants have been investigated and are more or less known.

The typical domestic animal is, likewise, confined to a relatively small area, particularly in modern times. The animal apparently

³ Italic numbers in parentheses refer to Literature Cited, p. 59, and such numbers followed by "x" refer to Sources of Unpublished Material, p. 91.

develops some sense of values in regard to the healthfulness of its foods, and when, as in ancient times, it was allowed to roam over large areas of the better lands, nutritional disorders were probably not often observed. However, the ever-increasing population forced the adoption of fundamental changes in the habits of the herdsman. He was required to confine his cattle to that portion of the landscape that he controlled, or he was required to move them to an entirely new and probably much less desirable environment to which they were not adapted. Those who were fortunate enough to have chosen or to have preempted the better lands experienced no difficulties with their cattle, but those to whom fell the less desirable lands or who were driven to some other locality deficient in some respect soon found that their cattle developed disorders that could not be combatted by any known methods. The fact that cattle thrived better on some lands than on others was thus quickly noticed, and we have many references as early as the eighteenth century of the adverse effects of grazing of cattle on certain lands while often adjacent lands were found to be healthful.

Man, in general, has derived his food, in modern times, from such a variety of localities that mineral deficiencies are seldom traceable to any specific soil cause, although such disorders as goiter are confined largely to persons living in areas where the soils and waters are deficient in iodine. Because of the varied diet and the modern methods of food processing and transportation, the fundamental problem of the relation of sources of food to human disease is more difficult to solve. However, some of our people do live in restricted localities and on highly restricted diets, and it should be possible to approach the problem of human nutritional diseases in such localities in much the same manner as one would for animals. Such relationships between the soil as an environmental factor and the occurrence of human disorders have already been observed, but it is very probable that some of these are quite accidental or at least very indirect, such as the observation of Virtanen (576) in 1928 that tuberculosis in Finland is more prevalent among people who live on acid soils than among those living on soils that are more nearly neutral. Many such observations of cause and effect are, however, quite direct, as, for example, the fact observed by Byers (102) that in certain localities in Mexico the inhabitants exhibit the characteristics of selenium poisoning, while in the selenium areas in South Dakota "a high incidence of symptoms pointing to gastric or intestinal dysfunction, and a few instances of apparent hepatic dysfunction, both probably the result of continual selenium ingestion," have been reported (531).

Thus, even cursory observations throughout the past two or three centuries have indicated that certain soils were good while others were poor, and Corlette (123) has stated:

In working through veterinary literature on osteomalacia, rickets and associated conditions of disease, I found my attention over and over again attracted to the character of the soil, climate and vegetation in their relation to the prevalence of cases.

The nutritional diseases of animals which have been traced to soil characteristics may be divided into two general classes. The first, and more important, are those reported to be due to a deficiency of one or

more of the inorganic nutritional elements in the food, eventually resulting in bone maladies, anemia, goiter, and related diseases. The second are those claimed to be the result of excesses of certain elements in the soils and in the plants growing therein. Of the latter class, only selenium has been extensively studied, although investigations would doubtless indicate that other serious disturbances may really be due to poisoning of the animal.

The problem that confronts us is more complicated and insidious in nature than that of correcting observable deficiencies in the domestic animal, for although the humane and economic aspects of this part of the difficulty are great, the more important aspect is man, who depends upon a high quality of food for his growth, health, and reproduction.

It is probably true that not only are a large portion of our people existing on a diet deficient in the protective foods such as milk, eggs, fruits, and vegetables, but also a large portion are obtaining only a minimum amount of these foods. If this is true, we are then confronted with the problem of the variation in the nutritive value of such foods as the fruits and vegetables in particular. Are these fruits and vegetables as highly nutritious as is possible? The calcium content of cabbage, one of the most commonly used protective foods of the vegetable class, may vary from 0.40 to 1.60 percent of the dry weight (refer to appendix), and the average is about 0.73 percent. Comparable variations in other nutritive elements in this and other foods are often reported. It is entirely probable, therefore, that the value of a minimum diet of protective foods may be reduced significantly below the minimum requirement through the use of inferior foods. The necessity for maintaining these foods at an optimum nutritive level is thus apparent.

The problem is intensified because of the location of our large cities and of our truck farms that supply these cities with food. Most of our large cities are located on the well-leached Podzols and podzolic soils that are acid in reaction and low in the bases. The truck farms are of necessity located near the cities on the same soils, and large quantities of truck crops and fruits are shipped to these cities from localities where nutritional disorders due to mineral deficiencies in the soils have been noted in both humans and animals.

Although many factors other than the poor quality of food probably are responsible for dietary difficulties, this factor is believed to be an important one; and if it is important, then it certainly remains the duty of the agriculturalist to supply for at least the protective portion of the diet a food of the highest nutritional quality.

SOME SOIL CHARACTERISTICS CITED IN RELATION TO THE OCCURRENCE OF CERTAIN NUTRITIONAL DISEASES OF MAN AND ANIMALS

In the discussion that follows, the occurrences of certain nutritional deficiency diseases are cited only as indicators of observed differences in soils. No attempt has been made to discuss the diseases themselves.

STUDIES OF THE RELATION OF SOILS TO BONE DISEASES

The occurrence of many disorders in animals, believed to be due to deficiencies of elements such as calcium or phosphorus, are often suspected to be related to deficiencies of one or more of these elements in

the soil. Direct evidence of such relationships is, however, generally difficult to obtain, because of the numerous factors involved and the overlapping of these factors. Thus, climatic conditions may be such as to cause in some years a significant reduction in the quantity of an element in the plant or a marked change in the usual ratio of one element to another. Mineral deficiencies in the animal may also be due to an inadequate supply of food in general, to the unavailability to the animal of the mineral elements in the plant, or to the species of plant available to the grazing animal. Many of the factors causing variations from the normal in plants are very closely interrelated, and a discussion of them will be found in the second part of this review. It is generally agreed, however, that frequently there are differences in soils, often in the same locality, that are associated with the occurrence of disorders in animals, and it is the purpose of this part of the review to cite some of the observed characteristics of these soils.

An appreciation of differences in the capabilities of soils to produce healthful food for animals is not a recent achievement, for Neyen (436) stated that Gleditsch in 1787 noted the occurrence of a bone trouble of cattle in certain localities in Europe, and other early observations were made by Le Vaillant (555) and by Lichtenstein (350) in their accounts of their travels in South Africa at the beginning of the nineteenth century. Von Gohren (217) reported in 1861 that a disease of the bones of cows occurred over a considerable region in Germany in 1859, a year of drought. He associated the disease with a low content of inorganic elements in the feed, and he reported that the addition of bone meal to feeding stuffs cured the disease. Among other early investigators of the relationship between the composition of the forage and the occurrence of bone diseases were Grouven (230) in 1865, Karmrodt (310) in 1868, Roloff (498) in 1869, Klimmer and Schmidt (317) in 1906, Scheunert, Schattke, and Lötsch (511) in 1911, and Zuntz (607) in 1912. All of these workers reported that the calcium or phosphorus content of pasture or hay in the abnormal areas differed from that in the healthful areas, and in many cases their analyses indicate that these differences were real and significant. In other cases, however, particularly where differences in calcium are cited as being important, the minimum value of calcium given is often not less than that now believed to be the minimum required by animals (403, p. 354), which may indicate, of course, that certain of the conclusions of these early investigators were not justified. It is interesting to note, nevertheless, that considerable importance was attached to soils and their properties as factors in animal health. Later investigators may have been more accurate in some of their classification of diseases, but many of them have not improved their soil investigation technique to any important degree.

A bone disease of livestock was first observed in Australia in 1895 by Potts (471) following a severe winter and a hot, dry summer. A note of the investigator indicates that the soils involved were largely formed from granites and that there was a scarcity of lime. Henry (260, 261, 262) has reported that the occurrence of bone disease in Australia was related to definite areas and soils that were often sharply marked off from those parts of the same country where the disease did not occur.

Guthrie and coworkers (234) stated in 1914 that bone diseases in Australia were generally most acute where the soils exhibited a marked

deficiency in lime, and that when animals in one portion of a district were affected and in another unaffected a notable difference in the lime content of the soils of the two localities usually existed. The normal areas were described as being mostly dark-colored soils, well supplied with lime and possessing a porous subsoil. The deficient areas had light-colored, poor soils and clayey subsoils. The diseases were reported to be most prevalent on geological formations like sandstones, aplites, and granites, which yield silicious sandy soils, poor in lime. The normal areas were located on limestone, basalt, dolerite, diorite, and gabbro formations, which are well supplied with lime. Unaccountable exceptions to these general observances were reported, however. A comparison of soils in some of these good and poor regions is shown in table 1.

TABLE 1.—Comparison of composition of soils from coastal areas of New South Wales on which animals were affected and on which they were unaffected with bone disease

[Guthrie and coworkers (234)]

Character of area	Reaction of soil	Clay	N	CaO	K ₂ O	P ₂ O ₅
Affected (17 soils).....	Usually strongly acid....	Percent 43.2	Percent 0.157	Percent 0.168	Percent 0.087	Percent 0.047
Unaffected (17 soils).....	Neutral to acid.....	58.0	.282	.465	.143	.184

Although observations in South Africa more than a century ago indicated that cattle were subject to a bone disease, no formal investigations seem to have been conducted until fairly recently, when in 1918 Viljoen (575) studied the occurrence of lamziekte, a disease later described by Theiler and others (554) as "an indirect consequence of phosphorus deficiency, the direct cause of the disease being a 'ptomaine poisoning' brought about by infection of carcass debris with a specific toxigenic saprophyte; an anaerobic bacterium reminiscent of, but not identical with, *Bacillus botulinus*." Viljoen reported that in the Bechuanaland area lamziekte was most prevalent on limestone and dolomite formations. Sandy soils, especially shallow ones or those mixed with limestone, seemed to be the most common formation on which the disease occurred. He stated that lamziekte was unknown or very rare on farms where the soil formation is a deep alluvial one. He believed that the moisture relationships of the soils were very important, for he observed that the disease appeared most commonly on farms where the soil formation is such that very little moisture is retained. Plants growing in such soils quickly responded to changes in the climatic conditions.

This relationship between climate and the occurrence of bone disease had already been indicated by Von Gohren (217), and Lewite (348) had found in 1907 that oat plants grown in dry years, characterized by outbreaks of bone diseases near Leipzig, contained less phosphorus and more calcium than did the normal plant. Viljoen, however, appears to be the first to relate this phenomenon to such definite soil characteristics as structure and the water relationships.

Further investigations by Theiler, Green, and du Toit (554) in 1924 confirmed the effect of climate as well as soil on plant composition and the occurrence of deficiency diseases. Thus, in table 2 the phosphorus contents of the vegetation consumed by cows suffering from osteomalacia and lamziekte in South Africa are given for different times of the year.

TABLE 2.—Composition of dry matter in vegetation from Armoedsvlakte veld with special reference to phosphorus

[Theiler, Green, and du Toit (554)]

Date	Composition of dry matter						
	Crude protein	Ether extract	N-free extract	Crude fiber	P	Estimated energy value (starch=100)	Ratio of starch equivalent to P
	Percent	Percent	Percent	Percent	Percent		
Nov. 10, 1919.....	19.4	5.5	41.0	22.5	0.26	56	100:0.46
Dec. 8, 1919.....	14.3	5.6	46.8	25.6	.14		
Jan. 15, 1920.....	13.8	5.5	48.0	25.0	.10	52	100:0.19
Mar. 4, 1920.....	7.2	3.4	49.8	23.7	.10		
Apr. 19, 1920.....	4.9	2.4	51.6	35.0	.05	32	100:0.16
May 11, 1920.....	4.1	2.2	52.9	34.9	.03		
June 8, 1920.....	4.0	2.0	53.7	33.1	.04	25	100:0.16

It is stated that except for the very early grass this veld is deficient in phosphorus as well as protein and other desirable constituents all the year round. That this deficiency may be due to the unavailability rather than to a deficiency of phosphorus in this soil is apparent from the fact that the available P_2O_5 in the soil is reported to vary from 0.0005 to 0.001 percent as compared with a total P_2O_5 content of 0.03 to 0.12 percent. These authors cite another soil in the same area that contains 0.009 percent of available P_2O_5 and that is said to support a pasture that will maintain healthy cows.

The Armoedsvlakte soil is described as a shallow soil, varying from dolomitic outcrop to a few feet of leached material. According to the extent of leaching it varies from a heavy dolomitic loam to a sandy soil containing very little carbonate, although still alkaline.

More detailed descriptions of soil characteristics, particularly parent material, as related to the occurrence of bone diseases in animals have been made in New Zealand than in any other locality. The earliest reference by Reid and Aston (479) in 1910 cites this abnormal condition among pastured animals as being the result of "an excess of organic matter and a deficiency of salts in the soil upon which the animals were depastured [sic]."

The interest in soils probably arose from the observation that adjacent pastures often had quite different effects on the health of the animal. Thus, Aston (33) in his study of the occurrence of osteomalacia in the Wairarapa district observed, in 1930:

The district is a very large one, in which, while it contains the most phosphorus-deficient pastures yet met with in New Zealand, certain areas afford soils of great fertility and pastures of high nutrient value. *The evidence of the relationship between soils and animal health is convincing and continuous from the soil, the pasture, and to the animal.*

In a further investigation of the soil characteristics in relation to the occurrence of osteomalacia in the Wairarapa district, no definite relationship between soil structure and the occurrence of disease was noted, for it was stated in 1931 by Aston and others (34):

The soil-analyses reveal no soil of excessively coarse nature, such as pumice, sandy silts or even the coastal dune sands, so that from a mechanical point of view the soils involved show no inherent structural defect. The finest textured soils on which the disease occurs are silt clays, and the coarsest are fine sandy loams.

The investigations of the New Zealand workers in connection with the occurrence of bone diseases included many analyses of both pastures and soils (34), observations of the characteristic flora of good and affected pastures (32, 34), and descriptions of soils with particular reference to parent materials (27, 32, 34). Thus an affected soil at Martinborough was described as being quite heavy with a hardpan formation in the flat lands, while a good soil in the same area was classified as a clay loam derived from limestone (34). Many difficulties with bone diseases were reported as occurring on certain soils derived from an air-borne volcanic material in a region of heavy rainfall (32). The pH-value range was reported to vary from 5.0 to 5.8 for the affected soils and 6.1 to 6.3 for the better soils.

An occurrence of a bone disease referred to as osteomalacia was investigated in Norway by Tuff (567) in 1923. The disease was more severe in dry years, but Tuff stated that there are also typically osteomalacia districts where the disease occurs as an enzootic, independent of the amount of rain.

He reported that if hay from normal farms was given to the stock affected with osteomalacia, or if the owner of a threatened farm hired summer pasturage on a normal farm, his animals were not subject to the disease. Conversely, animals moved to pasture on deficient soil acquired the disease. Tuff examined the mountain rocks in the abnormal areas and reported that they contained only 0.002 percent of P_2O_5 , as compared with 2 percent in the substratum of the normal areas.

In 1926 Elliot and his coworkers (165), from their quite extensive investigations in Great Britain of the effect of the mineral composition of pastures on the health of the animal, concluded that a comparison of the

health, quality, and number of sheep grazing a pasture with the analytical data for the individual (pasture) samples seem to indicate that the highest mortality amongst the sheep * * * is found where the pasture shows the lowest percentage of mineral matter. It is interesting to compare the results for two very poor pastures and a very rich pasture with the average results for cultivated pastures. [See table 3.] As types of very poor pastures, the samples from the island of Lewes and the Falkland Islands are taken. * * * there is a high mortality amongst sheep in the Falkland Islands. Samples taken from the paddocks at Lord Astor's racing stables at Taplow, where the grass has received special treatment to render it suitable for grazing for race horses, have been chosen as representative of a very rich pasture.

TABLE 3.—*Composition of some British pastures*

[Analyses of William Godden (211)]

Locality	Composition of dry matter					
	Crude protein	K	Na	Ca	P	Cl
	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
Island of Lewes (eaten).....	8.37	0.563	0.280	0.204	0.106	1.340
Island of Lewes (not eaten).....	6.43	.448	.281	.211	.077	1.029
Falkland Islands.....	10.31	1.644	.215	.161	.213	1.650
Taplow pastures.....	22.26	1.991	.518	1.767	.435	3.562
Average for cultivated pastures.....	17.69	2.637	.182	.718	.321	2.830

The deficiency of mineral elements in the poor pastures is apparent. Even the "eaten" areas in the island of Lewes contain about one-third as much calcium and phosphorus as well as only one-half the

protein found in the average pasture, while a comparison with the Taplow pasture is even more unfavorable.

Orr (444) in 1929 reported on the difference in composition of good and poor pastures in the hilly districts in the Western Highlands and in the southwest of Scotland, where it is claimed a bone disease, locally referred to as *croitich*, in sheep occurs. Some of his data are given in table 4, and they indicate that both calcium and phosphorus in the affected areas may be deficient (403, p. 354).

TABLE 4.—Composition of good and poor Scottish pastures

[Orr (444)]

Description	Composition of dry matter						
	Crude protein	Ash	K	Na	Ca	P	Cl
	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
Healthy hill pasture.....	15.62	5.85	2.21	0.27	0.46	0.29	0.64
"Croitich" area.....	12.80	2.82	1.25	.09	.15	.13	.52

Elliot, Orr, and Wood (164) reported in 1926 little or no difference in the energy values of good and poor pastures and, in agreement with other investigators (126) in England and other localities, emphasized that the poor nutritive values of some pastures were due in part to their botanical composition. A more complete discussion of the influence of soils on botanical composition will be presented in subsequent sections.

Aphosphorosis, or sickness from deficiency of phosphorus, has been reported in certain areas in Ontario, particularly in dry years (384), and soil descriptions have been made in some cases. Thus the Brookston silt loam in Canada has been reported as a soil on which grazing animals do well, and herbage growing on this soil, which in one locality has an available phosphorus content of 368 pounds per acre, according to Atkinson and Woodward (35), contains nearly twice as much phosphorus as does herbage growing on the Berrien sandy loam, which contains, according to these authors, only 84 pounds of phosphorus and is considered to be less desirable for pasture. This is in contrast with the experience in Michigan (229), where the medium-textured soils such as the Brookston and Gilford loams are claimed to support plants of low phosphorus content and on these soils nutritional disorders in cattle are reported to occur. Such anomalies appear to be common, and their occurrence suggests that much of fundamental importance as to soil and plant and animal relationships is as yet obscure.

In France, Maume and Monteil (400) in 1938 observed animal diseases in some areas where soils were developed from gneiss, mica schists, or granites. The authors stated that the forage grown on these soils was deficient in calcium and phosphorus. In India, Davis (143) states that it has long been known that the native cattle of Bihar are small and give far less milk than the cattle in many other districts where it is claimed the soils are more favorable so that good nutrition is possible. Other observations of the importance of soil characteristics in the incidence of bone diseases have been made by Crawford (127) working in Ceylon and by Rubino (499) in Uruguay.

Although American investigators have generally recognized that soil conditions may be responsible for bone diseases in animals grazing

certain pastures, few careful descriptions of abnormal soils have been made. Faville (173) in 1896, a report of the Bureau of Animal Industry for 1898 (571), and Parker (449) in 1904 refer to possible environmental conditions as being important in the occurrence of nutritional diseases. In 1920 Cary (111) described a deficiency disease in southern Alabama along the Coastal Plain region, sometimes called the wire grass region. The terms "sweeny" or "creeping sickness" were applied to the disease, which Cary states was actually osteomalacia. The soil was described as being sandy and deficient in lime. Schmidt (512) in 1924 investigated a fatal disease of cattle in Texas known as "loin disease" or "down-in-the-back." He observed that the occurrence of the disease was "limited to the low, flat area bordering on the Gulf of Mexico and extending inland to the north to a distance of about 100 miles. The soil is usually noncalcareous." Becker (52) in 1933 stated in connection with his investigation of "bone chewing" in Florida that the deficient areas were associated with certain of the "clay soils of the lower Appalachian Mountain regions, soils of the flatlands more distant from the flood plains of rivers that drain more fertile fields, and the muck and lighter sandy soils of the coastal plains."

Welch (582) in 1924, in his report on bone chewing in Montana, stated that throughout large areas in Montana, cattle showed symptoms of a deficiency disease. The occurrence of the disease was usually associated with the feeding of wild hay and usually that grown on low ground. As a rule the hay contained redtop, wire grass, bluejoint, and various swamp grasses. Cattle that were fed alfalfa, clover, or other legumes were not often observed to chew bones, even though legume forage may have been used for only a part of the year. Although the disease had been observed since 1910, it did not become acute until the better lands were taken over for farming and the poorer lands only were reserved for grazing.

Scott (516) in 1929 examined the forage in several localities in Montana where depraved appetite of cows had been observed. The data on grasses from normal and deficient areas indicate that the cattle must obtain in any of the areas studied in Montana a forage relatively low in phosphorus and that there is little difference in the composition of forage from "good" or "poor" lands. The data in table 5 illustrate the point.

TABLE 5.—*Calcium and phosphorus content of forage crops in Montana*

(Scott (516))

Forage	Year sampled	Description of area	Number of analyses ¹	Analysis of dry matter	
				Ca	P
				Percent	Percent
Alfalfa.....	1925	Normal.....	11	1.44	0.181
Do.....	1927	do.....	9	1.54	.164
Do.....	1926	Deficient.....	6	2.32	.161
Do.....	1926	do.....	3	2.02	.160
Do.....	1927	do.....	1	1.49	.141
Clover.....	1925	Normal.....	3	1.05	.170
Do.....	1927	do.....	2	1.23	.123
Do.....	1926	Deficient.....	1	1.82	.147
Do.....	1927	do.....	1	1.46	.139
Wild grass.....		Normal.....	9	2.44	.157
Do.....		Deficient.....	14	2.94	.113

¹ Maximum and minimum values are not given by the author.² The season of 1927 was wet. No information available for other years.

The phosphorus content of the alfalfa from even the better areas is about as low as that found anywhere in the world for this plant when harvested at the one-half bloom stage. (See appendix.) The calcium is slightly below average for alfalfa, but the Ca:P ratio is 10 as compared with 7.5, the ratio of the average values shown in the appendix. The phosphorus contents of the clovers from both regions are not of sufficient difference to be significant. The analyses of the wild grass samples do indicate a difference of phosphorus, although the amount found in the deficient areas is not so low as it was in the pasture grasses studied by Theiler (table 2). It is instructive to note that the calcium content of forages from deficient areas was generally found to be higher than in those from the normal areas, which may mean that a Ca:P ratio less favorable to the animal is a factor to be considered.

According to Eckles, Becker, and Palmer (161), mineral deficiencies in cattle have been noted in 32 counties in Minnesota. The soils, which at one time were covered with grass, are of glacial origin and are described as young gray drift. While the phosphorus contents of the forages from these areas show some differences according to the prevalence of disease, it must be admitted that they are all very low. For example, the average phosphorus content of 9 samples of alfalfa from normal regions in Minnesota was 0.257 percent, and that from abnormal regions was 0.189 percent, while the average of all available data for alfalfa at one-half bloom stage is 0.272 percent. The average prairie hay in abnormal regions contained only 0.100 percent of phosphorus, while that from normal regions contained 0.108 percent. Such differences are probably negligible, and the forages in the good regions are probably on the border line as far as their nutritional value is concerned.

In harmony with the results from other deficient areas, it was found that osteomalacia was more severe in years of drought than in wet years. Likewise, the forage contained more phosphorus in the years of higher rainfall.

Later work from Minnesota by Eckles, Gullickson, and Palmer (162) showed extreme cases of phosphorus-deficient forage in prairie hay grown on the Nygaard farm. From 1925 to 1931 the phosphorus contents of the hay from this farm varied from 0.047 to 0.094 percent. Such values compare very well with the results of Theiler in South Africa, and they are lower than the values found in Montana. It was estimated by the authors that a cow receiving this hay along with the usual ration of oats would receive only 68.6 percent of the phosphorus necessary for maintenance and milk supply.

Huffman and Taylor (281) in 1926 surveyed areas in Michigan where depraved appetite occurred and found that soils in these areas contained approximately one-half as much phosphorus as did normal soils. Analyses of three samples of forage collected by these investigators do not indicate a phosphorus deficiency, but the rather unusual circumstance is presented of a low phosphorus content in their forage samples collected in the early spring followed by a 40-percent increase by mid-summer and a further increase by fall. They state that depraved appetite in Michigan cattle seemed to be most severe in the spring and abated during the fall. They further found that, although a mineral mixture similar to the mineral combination found in alfalfa hay failed to relieve depraved appetite, well-cured alfalfa hay effected a cure in all cases tried.

Ten samples of Michigan alfalfa hay examined by Reed and Huffman (478) had a phosphorus content well below the average for alfalfa shown in the appendix of this publication.

Phosphorus-deficient areas have been reported from various localities in Wisconsin by Hart and others (242), but apparently no relationships have been determined between the occurrence of the disease and the soils or the nature of the forage. Mineral-deficient areas have also been suspected in Merced County, Calif., according to Hart and Guilbert (245). Soil studies have not been reported, however, from this area.

A recent survey in Pennsylvania (193) is reported to have failed to reveal any deficiency of phosphorus in the forage of that State. It had been assumed that since a large proportion of the soils in the State are responsive to phosphate fertilization, there might be a deficiency in the forages in some places. The conclusions were that the data obtained did not support this assumption.

STUDIES OF THE RELATION OF SOILS TO THE NUTRITIONAL ANEMIAS

The properties of soils in many parts of the world have been described in connection with the occurrence of troubles in which anemia has been reported as a symptom. From the discussion that follows it is not to be inferred, of course, that a direct connection with soil conditions has been established, but it will be noted that evidence has repeatedly been produced that would indicate significant differences in soils associated with healthy animals and those associated with affected ones. McGowan and Smith (367) state in 1922, for example, that a disease of sheep referred to as "pining" was noted in the southern part of Scotland as early as 1807, and a recognized cure was to shift the sheep to a more succulent herbage or pasture grown on limed soils. Sheep on moors or moss-covered lands were always susceptible to the disease, but no symptoms were ever noticed in sheep grazing on the steep and rocky lands where herbage was sweet and short. Regions underlain by porphyritic rock were associated with the occurrence of pining, and it appeared occasionally in the sandstone regions, but the limestone regions were almost always free of it. Investigations of soils in these areas by Greig and others (226) in 1933 seemed to indicate that the iron content was low in affected soils, although it was reported by Godden and Grimmett (212) in 1928 that additions of iron to these soils did not result in any increase in the iron content of either oats or mustard. The work of Corner and Smith (124) in 1938 indicates that cobalt may be the limiting factor in these areas.

A nutritional anemia referred to as bush or Tauranga disease was known in New Zealand in 1900, when Gilruth (210) stated that in localities where the disease occurred there were "healthy" and "unhealthy" lands, and that vegetation grew as luxuriantly on one as on the other. Aston (30) examined these soils in 1912 and found that abnormal soils were composed largely of air-borne pumice, an acid lava that had been so mixed with gases in the molten state that on cooling it presented a spongy structure with the hardness of rock. He reported also that the available phosphorus was generally low and that manganese was high in these soils.

Manurial experiments (475, 476) indicated that applying iron compounds to the soil did not always alleviate the disease, although

improved health of the cattle could usually be obtained by dressing pastures with superphosphate, blood, bones, and guano. Heavy applications of superphosphate to soils with a low content of available iron were reported to be sometimes successful in combatting the disease, and the use of iron salts as a drench in advanced cases was sometimes satisfactory (477).

More recent investigations of the soils associated with bush sickness showed that most of them were well leached and that they did not contain any appreciable amounts of calcium and manganese likely to be antagonistic toward the availability of iron. Taylor (549) believed that leaching was primarily the reason why certain Mairoa farms were unsound for stock. He inspected 12 farms with a definite history of abnormality and found that

The predominating structure in each case was volcanic ash upon porous sandstones or loose sands. In the neighboring healthy areas the sands were practically absent, and the ash was underlain by limestone with its attendant residual clay, or by Jurassic mudstones.

Grange et al. (218) later claimed that profiles of six typically "bush-sick" soils showed evidence of distinct podzolization. They were all coarse-textured soils containing only small percentages of clay. The chemical analyses of the soils showed that some contained a moderate supply of available phosphorus and potash, while others were definitely low in both of these constituents. However, normal volcanic soils were also found to be low in these constituents, and they were often of coarse texture. Texture, in fact, does not seem to be an important factor, for Aston (31) states that certain coarse sandy soils near Rotorua are recognized locally as "hospital" lands and supply a nutritious food for animals.

In their study of bush sickness at Glenhope, Nelson, New Zealand, in 1932, Askew and Rigg (26) reported that the deficient soils were well leached and occurred mostly over granitic formations. Evidence that a deficiency of iron in these soils was a factor in the disease was not demonstrated (487).

Dawson (144) seems to have been the first to associate "salt-sick," an anemic condition of cattle, to certain characteristics of some of the soils of Florida when in 1906 he stated: "The disease is usually confined to regions where the predominating soil is light, sandy, and more or less unproductive of nutritious grasses." He believed the disease to be comparable to that occurring on Cape Cod Peninsula and in certain counties in southern Texas. He observed that salt-sick occurs in nearly all parts of Florida.

In 1931 Becker, Neal, and Shealy (51, 429) found that "'salt-sick' is a naturally occurring nutritional anemia in cattle, the feed of which has been restricted largely to grass forage grown on certain white and gray sandy soils, residual muck and peat soils not subject to overflow from more fertile watersheds." No trouble was experienced on clay soils.

More detailed soil work by Bryan and Becker (94) in 1935 showed that soils from healthy areas contained over twice as much silt and clay in the first foot and over three times as much in the second foot as did those soils in the salt-sick area. Their survey revealed that for many years cattlemen "have found it necessary to transfer 'salt-sick' animals to healthy ranges, sometimes known locally as 'hospital'

lands. Where this practice was observed, the so-called hospital lands have been either on sandy loam or on one with a sandy clay strata from 15 to 30 inches below the surface."

It was found that salt sickness occurred on both calcareous and low-lime soils, and it was believed that this indicated that the disease was not related to calcium deficiency. The soils associated with salt-sick were Dade sand, Leon sand, Leon fine sand, Portsmouth fine sand, Norfolk sand, and Norfolk fine sand. Those in the healthy areas were the Bladen fine sand, Fellowship sandy loam, Gainesville fine sand, Hernando sandy loam, Hernando sand, Lakewood sand, Norfolk sand, Norfolk fine sand, Orangeburg loamy sand, Orangeburg fine sandy loam, Orangeburg fine sand, and Ruston sand. The average composition of each of these groups is given in table 6. The authors explain that the Norfolk soils have a wide range of types and phases which vary greatly in texture and natural fertility. This fact may account for this group of soils occurring in both the "salt-sick" and healthy ranges.

TABLE 6.—*Mineral composition of first foot of Florida soils in salt-sick and healthy areas*

[Bryan and Becker (94)]

Description of area	CaO	P ₂ O ₅	Fe ₂ O ₃	Cu	Silt and clay	pH
	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>P.p.m.</i>	<i>Percent</i>	
"Salt-sick"-----	0.050	0.027	0.057	3.97	4.3	5.17
Healthy-----	.227	.130	.604	8.56	9.9	5.24

Neal and Becker (428) examined samples of wire grass in 1933 from many of the healthy and salt-sick areas in Florida. Their data show that the mineral content of forage growing on burned areas is higher than that growing on unburned areas, and that the calcium and phosphorus are higher in forage from healthy areas than from salt-sick areas. Unfortunately, their extensive iron data are open to question because of the method of preparing the samples.

A nutritional anemia in calves described as a deficiency of iron was reported by Archibald and coworkers (12, 13) in 1938. They concluded from their analyses that the iron content of the roughage available to the animals in the abnormal areas was somewhat below that for hays from other sections of the State and of the country where such trouble is not experienced.

A trouble referred to as "pica," "coast disease," or "coastiness" was first observed in Australia about 1913, when it was claimed, according to Filmer (182), that in certain areas of the Denmark district of Western Australia farmers had been unable to rear calves since first settling in the area. In some cases certain paddocks only were affected, while adjacent ones were normal. Soil descriptions were rather sketchy, but they indicate a similarity to some soils in New Zealand. The disease is said to occur in various parts of Australia (386). The investigations of Filmer and Underwood (183) in 1934 indicated that a deficiency of iron in the forage was not a cause of coastiness, and in 1935 these authors (570) reported that a deficiency of cobalt in the soil and herbage was a factor, while later investigations by the same authors indicate that nickel is of importance (184).

In Dartmoor, England, it was reported by Patterson (451) in 1937 that soils on which sheep suffer from "pining" have a mean cobalt content of 3.9 p. p. m., while lowland soils on which sheep recover have a cobalt content of 16.7 p. p. m. Analyses of the pastures show that those in the unhealthful areas have a mean cobalt content of 0.20 p. p. m. and those in the recovery areas 0.45 p. p. m. It was also indicated by Kidson (316) that the cobalt contents of soils in this area are in general related to the magnesium content of their parent rocks.

In New Zealand (22, 24, 25, 148) it was reported that the cobalt content of soils in bush-sick areas contained only a trace of cobalt as compared with the healthful areas, although "soil results indicate that a low cobalt status of the soil is not always a satisfactory index of the need for cobalt supplements for stock." Generally, however, Kidson (315) believed that "soils affected with 'bush-sickness' and allied stock ailments have comparatively low cobalt contents, often less than 2 p. p. m. of cobalt." It was reported by Josland (308) that 1 mg. CoSO_4 per 200 gm. of live weight of the animal is not toxic.

A disease of cattle known as "nakuruitis" has been reported from certain areas of Nakuru, in the Kenya Colony in East Africa, by Orr and Holm (446). "The top soil in the affected areas consists of material which has obviously been erupted from the adjoining extinct Menengai volcano."

Neal and Ahmann (427) in 1937 reported that cobalt was not detected in forage from certain areas in Florida associated with a nutritional disturbance in cattle that was cured when cobalt was fed as a supplement, but later work by Rusoff and others (500) failed to demonstrate that cobalt was present in forage from either salt-sick or healthy areas. In fact, a spectrographic estimation of copper and several other trace elements was reported as giving no indications of differences in samples of wire grass collected from these areas.

A nutritional anemia among children in certain areas of Florida was investigated by Abbott, Neal, and Bryan (2) in 1934. In a brief report they state that in one section situated on soils overlying hard rock phosphate subsoils only 3 percent of the children had anemia, but within a distance of 6 miles, where the soil is for the greater part Leon, located in the flatwoods country, 96 percent of the children were affected. They add: "Hookworm and malaria are known to contribute to this condition, but it was found that the number of children infected with these parasites was comparable in the two localities."

In a later study by these authors (1) it was reported that the Leon, Portsmouth, and Norfolk series (all low in iron) prevail in those districts having the highest percentage of anemic children, and that the sands and sandy loams of the Hernando and Hoffman series (all high in iron) predominate in the districts having the lowest percentage of anemia.

Riceman, Donald, and Piper (484) have recently described what may be a copper deficiency along the southeastern coast of South Australia. The soil is derived from blown calcareous sand containing over 60 percent of CaCO_3 in the surface soil. Sheep grazing the area were reported to suffer from an ailment similar to coast disease. Normal vegetative development was attained with the use of CuSO_4 , but no statement was made relative to any improvement in the sheep.

The clinical aspects of a deficiency disease in sheep in Western Australia have been studied by Bennetts (54), who reported in 1937 that copper seemed to be the limiting factor. In their investigations of the soils in this district Hosking and Greaves (279) state: "The presence of this deficiency disease (enzootic ataxia) is closely associated in Gingen with any soil underlain by Cretaceous rocks." Country adjacent that is underlain by granite and gneiss is apparently unaffected, although "recent investigations show that the disease sometimes occurs on granitic formations."

Indications of a copper deficiency in Florida forages was reported by Becker, Neal, and Shealy (51) in 1931. These investigators stated that sometimes copper in addition to iron was necessary in order to prevent or to cure salt sickness.

In 1934, Svanberg (544) discussed the environmental relationships of an anemia of horses grazing on the very acid soils of northern Sweden. The pH value of these soils was found to be about 4.8 or less, and they are described as being sterile alum soils (546). Over comparatively large areas these soils are used for hay production and the crops serve as horse fodder. Although most of the major constituents and the iron, aluminum, and copper contents of the forages are normal, the manganese content is strikingly high, often 10 times as much in abnormal as in normal areas. Svanberg states that favorable natural conditions under which anemia in horses will not occur seem to include a soil rich in lime and containing not too much manganese.

Further investigations of this disease of Swedish horses by Carlstrom and Hjärre (105) in 1938 have attempted to show a relationship between vitamin B₁ and manganese. Carlstrom and Hjärre claim that this anemia is

due to an insufficient content of vitamin B₁ in the hay and this insufficiency is probably caused by too large a proportion of manganese in the hay. Everything leads to the belief, in effect, that this element destroys through oxidation the vitamin B₁ of the hay.

Further investigations will no doubt be necessary to clarify many of the factors involved in the occurrence of this disease.

STUDIES OF THE RELATION OF SOILS TO "LECKSUCHT"

A relationship between some property or properties of soils and the occurrence of a disturbance in cattle in Europe, known as lecksucht, has long been suspected (287, 288, 289, 296, 297, 318, 346, 435, 448, 457, 470, 534, 545, 564). In some districts it was reported that the poor soils were developed from granites or mottled sandstone, whereas the unaffected or good soils were derived from gneiss. In other areas the poor soils were peaty in character and were low in potassium as well as in phosphorus, and the acidity was reported as being quite high.

Sjollema (526, 527) reported in 1933 that the copper content of hay-fed cows suffering from what he calls lecksucht was much less than that fed cows in good health. Nicolaisen and Seelbach (437) in 1938 reported that the application of copper sulfate to pastures not only improved the pastures but also the health of the animals suffering from what they also refer to as lecksucht.

"ALKALI DISEASE" OR SELENIUM POISONING

In 1934, as a result of a preliminary field survey of "alkali disease," Franke and coworkers (197) concluded that

Soil seems to be an important factor in the occurrence of this diseased condition of livestock. Every case investigated occurred on certain soils that were called Pierre clay or Pierre clay loam, or could be attributed to grain or hay grown on such soils. These soils have developed from the geological formation known as Pierre shale.

It hardly seems necessary to discuss extensively in this report alkali disease, or what is now known to be selenium poisoning, inasmuch as such complete summaries as those of Byers (100, 101, 104) and Moxon (419) are available and well known. It is sufficient to state that the disease is closely related to certain soils, and that the work of identifying these soils and the relative toxicity of their selenium contents has now practically been completed by Byers and his coworkers in the Division of Soil Chemistry and Physics of the Bureau of Plant Industry. A brief discussion of their work with plants and soils will be presented in later sections.

IODINE

Probably the first reference to the relationship of goiter to iodine deficiency in soils is that of Chatin (112), who in 1851 observed that the air is less iodized in the Alps than in Paris; the same is true of the rain waters. *The iodine content of tillable soil and its products is also higher than in the countries where goiter is more prevalent.*

Chatin offered no data to substantiate his statements.

Many investigations have been made of the iodine content of foods grown in goitrous regions and in goiter-free regions (3, 45, 174, 175, 265, 363, 445, 548), but only those dealing with actual soil conditions will be discussed here.

Von Fellenberg (176) in 1924 reported that the soil at Effingen, Switzerland, a goiter-free area, contained 11.9 p. p. m. of iodine, while that at Hunzenschwill, where 56.2 percent of the populations suffered from goiter, contained 0.620 p. p. m. of iodine.

Goiter is reported to be quite prevalent in New Zealand, and "its incidence is in general roughly inversely proportional to the average iodine in the soil," according to Hercus and others (264). Hopkirk and coworkers (277) showed in 1930 that the iodine content of soils in a certain goitrous region of New Zealand varied from a few parts up to 900 parts in 10 million. They found little correlation between the iodine contents of the soil and those of pastures grown on them. They state that, generally speaking, alkaline soils do not give up iodine to plants, while acid soils do so very readily. They found, however, that cows' milk from the goiter-free regions contained three times as much iodine as did that from affected regions.

Sykes (547) reported that limestone regions in particular were characterized by complete absence of goiter. Aston in 1930 reported the soil in a goiter region to be low in iodine (33). He made the

interesting observation "that stock make every effort to break bounds in order to escape, it is thought, from the iodine deficient pasture to other areas which are not deficient."

It was observed in New Zealand by Hercus and others (263) "that variation in the average amount of iodine in soils containing more than 10 p. p. m. has little effect on the small incidence of goiter which there exists; but as the amount of soil iodine decreases so the incidence of goiter rises." The same report states that the most important




FIGURE 1.—Relationship between the average amount of iodine in the soils and the regional incidence of goiter among school children in New Zealand. (Based on data from Hercus, Benson, and Carter (264).)

causes of variation are (1) the combination of the iodine in the soil (an analogy to the observation of Byers (104) concerning selenium), and (2) the presence of abundant iodine in the drinking water in regions in which the soil is relatively poor in that element.

The data of Hercus, Benson, and Carter (264) have been plotted (fig. 1) in order to bring out these approximate relationships as clearly as possible. The positions of the points indicate that several notable exceptions occur in the inverse relationship between incidence of goiter and soil iodine, but that there is a general agreement.

A striking inverse correlation between the quantity of iodine in the soils and foods and the incidence of goiter was reported by Hercus and coworkers (263) in 1931. Their data are given in table 7.

TABLE 7.—*Relation of iodine in food to incidence of goiter and to soil iodine in New Zealand*

[Hercus (263)]

District	Goiter incidence	I in soils	I in grass	I in foods	
				Eggs	Milk
	<i>Percent</i>	<i>P. p. m.</i>	<i>P. p. b.</i>	<i>P. p. b.</i>	<i>P. p. b.</i>
Samoa.....	Nil	285	326	394	11
Otago.....	19	64	68	58	8
Canterbury.....	64	25	67	41	6

Many anomalies occur in relating the iodine content of the soil and plant to incidence of disease, but Green (225), in his discussion of minerals in relation to disease, states that "a direct iodine deficiency is not the only cause of thyroid hyperplasia, and any factor which reduces absorption of iodine or increases the utilization of the thyroid hormone may also cause goiter." Similar views have been stated by Marine (395) and by McCarrison and coworkers (361, 362).

The incidence of goiter has received a great deal of attention in the United States, but studies relating this to the iodine content of soils or food plants are exceptional. In 1917 Smith (530) defined some areas in Montana in which sows had farrowed hairless or otherwise defective young, and observed that in many instances the rancher could save his pig crop by moving his pregnant sows a mile or two out of the affected district during the gestation period.

Welch (583) in 1928 described soil conditions in Montana that were associated with goiter, but his geological statements do not agree with the known facts, and his reasons for iodine deficiencies in these areas may, therefore, not be accurate. He attributes a lack of iodine in these soils to the leaching action of former lake waters on what he calls sedimentary soils. Actually, however, the soils in the eastern half of Montana, the goiter area, are Chestnut soils developed from a highly calcareous glacial drift, and it is quite probable that in such an alkaline soil the availability of the iodine is very low.

Kalkus (309) in 1920 noted that goiter and associated conditions were common among domestic animals grazing the valleys of certain rivers and lakes in north-central Washington. He found that "wild animals which range in the mountains above the goitrous districts are not affected. These animals visit the so-called 'deer-licks' that contain high percentages of iodine."

Several investigators in the United States have attempted to relate the iodine content of different crops to that of the soil, but their data will be discussed in the section "Factors Affecting the Mineral Composition of Plants" of this publication, inasmuch as they are not related to the incidence of disease.

OTHER NUTRITIONAL DISORDERS ASSOCIATED WITH SOILS

A nutritional disturbance believed to be caused by molybdenum was described by Ferguson, Lewis, and Watson (180) in 1938:

A large area under pasture in Somerset cannot be grazed by cows during the normal grazing season (April–October) without milk yields falling rapidly, causing marked loss of condition and even death. The available information seems to rule out bacteria, parasites, water supply or a particular herb as causal agents, and points to the presence of some constituent of the soil taken up by the herbage.

An extensive and systematic spectrographic examination of samples of herbage in abnormal and normal areas has shown that the molybdenum content of affected herbage is much higher than what is believed to be normal. Soluble molybdenum fed to cows produced symptoms similar to those observed in cows grazing affected pastures. In connection with this it might be noted that Stanfield (538) found 7 p. p. m. of molybdenum in alfalfa and 89 p. p. m. in barley hay, while 317 p. p. m. of molybdenum was found in *Atriplex nuttalli* S. Wats. growing in Steele shale in Wyoming (48).

Losses of cattle due to eating of oat hay have recently been observed in Wyoming by Bradley, Beath, and Eppson (86). It is not known what influence, if any, the soil may have upon the development of the toxicant, but "the localization of the problem suggests that local influence must be of basic significance." The difficulty is apparently believed not to be related to the occurrence of selenium.

Other nutritional disturbances confined to certain areas have been observed in British Guiana by Bone (82) and in Morocco by Velu (574). An endemic disease of horses in the mountainous districts of the island of Hondo off Japan has been observed recently (563). The appearance of the disease coincides with the growth of vegetation—May to November—and the pasturing of the animals during this period. A botanical examination of the pastures failed to establish that selenium-loving plants were present. No definite conclusions have been drawn as to the cause of the disease.

A chronic copper poisoning in sheep grazing in the vicinity of a nickel smelter has been investigated by Bisset (74). Copper, which was present in large amounts in the livers of older sheep, was believed to have been ingested with the soil rather than as a constituent of the grass.

A few observations have been made on the effect of deficiencies of other elements such as magnesium, manganese, and zinc on animals (55, 376, 485, 507), but relationships to environment have not been observed. Among the rather interesting although inconclusive studies is one from Czechoslovakia (433) in which it is claimed that 60 gm. of gold per metric ton was found in the horn of a roe grazing on a soil containing 0.1 gm. of gold per ton. No gold was found in the horn of an animal grazing on plants grown in a soil that contained no gold.

Several investigators in Europe (145, 146, 152, 491, 513) purport to have found inverse relationships between the magnesium content of the soil and the occurrence of cancer, but their studies are not of sufficient extent to be thoroughly reliable.

SUMMARY OF SECTIONS ON SOIL CHARACTERISTICS IN RELATION TO NUTRITIONAL DISEASES OF MAN AND ANIMALS

The inability of some soils to supply in proper amounts and proportions those elements essential to the well-being of man and animals has been known for more than a century. Investigations of certain nutritional diseases of animals during the last 25 years have emphasized the importance of soils, for, throughout the world, the occurrence of bone diseases, nutritional anemias, or of the effect on animals of excessive quantities of some inorganic elements has been characterized by the interspersing of normal and abnormal areas.

This fact has encouraged the study of soils in these areas, and many references in the literature describe the materials from which the soils were developed and the relation of these geological differences to the occurrence of disease. Unfortunately, however, in only a very few instances were the soils studied and described by competent soil scientists. Most of the descriptions give insufficient information about the true soils for their proper classification, and it is almost certain that many of the apparent anomalies could have been resolved with the aid of a competent soil taxonomist.

In general, it has been found that the low, poorly drained lands of the humid coastal regions, the moors, and the peaty soils are associated with bone diseases. Other physiographic regions are not free of soils on which these diseases occur, for many of the mineral soils over the entire world are deficient in nutrients, or, because of climatic conditions, fail to supply sufficient quantities of available nutrients essential to plant and animal life. Investigations of the forage growing on these soils indicate that two factors operate to prevent the animal from obtaining adequate supplies of minerals. (1) The pasture flora of fertile soils often are quite different from those of infertile ones on which the acid-loving plants of low nutritive requirements are able to crowd out plants such as the Leguminosae that require more of the nutritive elements. (2) Actual differences occur in the composition of the plant, and studies such as that of Greaves and Greaves (219) show that these variations have an important bearing on the health of the animal.

Nutritional anemias ascribed directly or indirectly to deficiencies of some of the minor elements such as iron, copper, and cobalt have been found to occur on many soils in the United States, New Zealand, Australia, and other countries. In both the Kenya Colony and New Zealand many of the soils have developed on erupted volcanic materials. These volcanic showers usually lie upon porous sandstones or loose sands in the abnormal regions and upon limestone in the normal regions, and many of these soils show some podzolization. In Florida the disease has been observed on the white or gray sands such as the Leon fine sand, Everglade mucks, and the young, poorly drained soils developed from marl in the vicinity of Homestead.

References to osteomalacia and anemia in human beings that may be correlated to the soil are few, but studies in Florida indicate that a nutritional anemia in children has been correlated with the soils from which the children derived the major portion of their food.

The causes of some nutritional diseases such as licking disease have not been well defined, although in some areas, at least, copper is said to be the limiting factor. Whatever the cause of this disease, descriptions of soil parent materials are often made, and, although true soil descriptions are lacking, some distinctive differences between deficient soils and normal soils have been noted. Thus, the soils in abnormal regions were often found to be developed from granites or mottled sandstone, whereas those in normal regions were developed mostly from gneiss, a potash-bearing rock, and in general, deficient soils were described as low moor soils or very light sands. A vast amount of work has been done in Europe on the composition of forages in connection with the occurrence of licking disease.

The presence of undesirable or toxic elements in the soils and their relation to disease has nowhere been studied in the systematic and comprehensive manner in which the selenium survey was conducted by the United States Department of Agriculture and some of the State experiment stations. That selenium poisoning is related to soils and their characteristics through the vegetation growing on them seems to have been demonstrated beyond doubt, while a nutritional disturbance in a certain area of England has been ascribed to the presence of excessive quantities of molybdenum in the herbage.

There are probably more contradictory data relating to the effect of soils on the occurrence of goiter troubles than to the occurrence of any other nutritional disturbance. A careful survey of the facts indicates, however, that the iodine in soils, either because of deficiencies or because of its low availability when present, must play some part in animal and human nutritional disorders.

Actually contradictory data are found in studies of all of these diseases considered in this review. The solutions of problems of this nature are difficult because of the large number of variables, and it may be necessary in the future to undertake studies statistical in nature, because the minute quantities and the catalytic character of the mineral elements involved often make it difficult to determine by chemical or physical means the role these elements play. Furthermore, modern techniques for soil examination and classification must be employed if the part played by soils is to be properly evaluated.

FACTORS AFFECTING THE MINERAL COMPOSITION OF PLANTS

EFFECT OF SOIL COMPOSITION ON MINERAL COMPOSITION OF PLANTS

The complex relationships between the composition of the plant and that of the soil have never been fully understood by man, although investigations of those relationships have been conducted for more than 300 years. Many problems confronting the agriculturist of the seventeenth century were not unlike those of today; and for this reason, and because of the remarkable ingenuity of the early workers, a few references to these investigations will be quoted from the unpublished portion of a manuscript prepared by C. A. Browne⁴ for the Yearbook of Agriculture, 1938, Soils and Men.

Nehemiah Grew was the first who attempted to make quantitative determinations and analyses of plant ashes. In his "Anatomy of Plants," 1682 [(228)], he recognized three classes of salts in vegetable materials: (1) The mineral or marine salts, such as common salt, (2) the essential salts, which occur in plants, such as tartar, and (3) the lixivial salts or water-soluble ash of plants and plant products. The influence of soil upon the yield of lixivial salts is shown in the experiments with scurvy grass, which in a garden soil yielded 1.69 percent of lixivial salt and in a soil near the seashore 7.29 percent. In his studies of plant composition and growth Grew recognized the important relationships of soils, which he classified into mellow, sandy, clayey, chalky, etc., and mixtures of the same. He recommended that experiments upon the growth of plants be conducted on single soils of types, "that it may appear how far any of these may contribute to the growth of a plant, or to one above another."

John Clayton's remarks [(114)], in 1688, upon the relationship of the quality of tobacco to the soils upon which it is grown is the earliest correlation of this character in the history of American agriculture. "The same sort of seed in different earth," writes Clayton, "will produce tobacco much different as to goodness."

⁴ Supervisor of chemical research, Bureau of Chemistry and Soils, U. S. Department of Agriculture.

Theodore de Saussure, of Switzerland, in 1804 made the first approximate analyses of the ashes of the leaves, stalks, bark, wood, straw, seeds, fruits, etc., of 79 different plant products, the results of which were assembled in the earliest published tabulations of this character [(504)].

Saussure's work demonstrated the falsity of the old view that plants could produce any of their mineral constituents by the transmuting action of vital forces. His numerous experiments established the following facts:

(1) The mineral constituents of plants are obtained in solution through the roots from the soil.

(2) Plants do not absorb mineral constituents in the same proportion in which they occur in the soil solution, but they have a marked selective power, some elements being assimilated in greater amounts than others.

(3) Herbaceous plants assimilate more mineral matter and exhaust the soil more markedly than do woody plants.

(4) Different soils have a profound influence upon the quantity and composition of the mineral constituents of the same species of plants.

(7) The different organs of plants differ in the content and composition of their mineral constituents.

(8) The percentage of ash in herbaceous plants reaches its maximum before the flowering stage and thereafter begins to decline.

Carl Sprengel, a German agriculturalist, was the first (1838) to attempt an enumeration of the various mineral elements of the soil that are necessary for the normal growth of crops [(535)]. Because of the general occurrences of the 10 mineral elements, potassium, sodium, calcium, magnesium, iron, aluminum, manganese, sulfur, phosphorus, and chlorine, in the ashes of all plants, he drew the conclusion that the elements named were all essential for the growth of crops. Sprengel also remarked that other mineral elements not yet discovered in plants, as fluorine, bromine, iodine, lithium, and copper, even though occurring in minute amounts, might also be necessary—a statement that foreshadowed modern investigations of a century later upon the so-called trace or minor elements of soils and crops.

Sprengel maintained that in addition to the soil elements which promoted the growth of plants, there were other elements, as lead, arsenic, and selenium, which injured the growth of plants if they existed in combinations that were easily soluble in water. He stated, however, that all plants were not affected alike by these injurious elements, one species having a greater power of resistance than another. Sprengel seems to have been the earliest to call attention to the possible injurious effects of selenium upon the growth of plants.

Of the early developments in the United States, Browne states that "the first scientists to make studies of the correlation of differences in American soils with the mineral constituents and nutritive value of the crops were the State geologists." Among these men was Amos Eaton, who "made an agricultural survey of Rensselaer County, N. Y., as early as 1821. Hitchcock, Jackson, Emmons, Shepard, Owen, Booth, and the other early State geologists made chemical analyses of the soils and crops of their respective States." Browne continues:

No writer has more clearly expressed the vital need of correlating the mutual chemical relationships of soils and crops than has Jackson. In the Patent Office Report on Agriculture for 1858 [(296)], Jackson published a comparative research upon the relationship of the chemical composition of soils in Massachusetts and Maryland to the chemical composition of the ash of the tobacco which was grown thereon. The ash of Maryland tobacco was higher in magnesia and lower in lime than the ash of the Massachusetts tobacco, a circumstance which led Jackson to the conclusion that these bases might replace each other to a certain extent without alteration in the healthy condition of the plant. He remarks that the limits of such substitution can be determined only "by making a long series of analyses of the same varieties of plants grown on peculiar soils, or on such as are artificially prepared for the experiments."

There are many factors that may operate to modify the mineral composition of the plant. These would include differences in soils, cultivation and the use of soil amendments, rainfall and other climatic

influences. These factors are overlapping in their effect or work simultaneously. One factor may influence another, as, for example, the climate is a dominant influence in the development of the soil, and through the soil, therefore, climate influences the plant. Differences in these factors will naturally operate to modify the plant in different ways. For example, it is possible (1) for the mineral composition of two plants of the same variety growing in different soils to be significantly different without there being any important difference in their size or the distribution of their parts such as leaf, stem, or seed head; (2) for the growth of the same variety of plants to vary (differences in yield) in different soils without any important differences in the proportions of the plant; (3) for two plants of the same variety growing in different soils to have quite different distributions of leaf, stem, or head; and (4) for properties of soils to so modify the quantities of plant constituents such as protein, carbohydrates, lignin, and cellulose as to influence the percentage distribution of other constituents, such as a greater deposition of starch or formation of lignin with a consequent reduction in the percentage composition of the mineral elements. The properties of two soils may be such as to modify the natural flora, and thus to produce plants quite different in mineral composition. The evaluation of these factors is still proceeding, and much valuable information of fundamental importance is being obtained.

For the purpose of this discussion, the subject of the variation in the mineral content of plants and the factors that affect it will be treated in three sections, namely:

(1) The soil. The summarization of this literature will be as complete as possible.

(2) Fertilizers. Typical investigations, particularly as they relate to our own soil types, and to the effect of the addition of one element on the absorption of others, will be reviewed.

(3) Miscellaneous. A few pertinent papers relating to the following factors will be reviewed: Stage of maturity, climate, irrigation, different varieties of the same species, and the composition of different parts of the same plant.

In discussing the effect of soils on plant composition, it must be remembered that almost all work in this field should really be classified as general environmental studies, since the effect of locality instead of the soil as an isolated entity has more often been the subject of inquiry.

In 1873, Fliche and Grandeau (186) determined the principal inorganic constituents in needles of the maritime pine (*Pinus pinaster* Soland) grown in two soils in France. They found that the variations of phosphorus, magnesium, and sodium were small when compared with those of calcium and potassium, which are given in table 8. The data show that the calcium content of the needles of the tree grown in the calcareous soil is higher than that of the other needles. Although the effect of the calcareous soil at Bas-du-Cellier is reflected in the calcium content of the plant growing on that soil, no explanation is readily apparent as to the differences in potassium, except that it is known that the availability of potassium in soils varies to a far greater degree than does that of calcium.

TABLE 8.—*Calcium and potassium in pine needles and in the soils in which the trees grew*

[Fliche and Grandeau (186)]

Locality and material	Constituent as percentage of ash		Locality and material	Constituent as percentage of ash	
	CaO	K ₂ O		CaO	K ₂ O
Quatre-Arpents:	<i>Percent</i>	<i>Percent</i>	Bas-du-Cellier:	<i>Percent</i>	<i>Percent</i>
Soil.....	0.35	0.07	Soil.....	3.25	0.04
Subsoil.....	.20	.03	Subsoil.....	24.04	.16
Pine needles.....	40.20	16.04	Pine needles.....	56.14	11.09

Liebscher and coworkers (352) in 1898 determined the potassium, nitrogen, and phosphorus contents of oat plants grown in pots containing 24 different German soils. The tests were conducted under fairly uniform conditions, and the results may reasonably be expected to show differences in the soils. The phosphorus content of the plants, for example, varied from 0.141 to 0.419 percent, and the potassium from 0.648 to 2.610 percent. The high value for potassium in the plant agrees with a high content in the soil in which it was grown, but with the lower values there is no agreement. Likewise, there is no correlation between the phosphorus content of the plants and of the soil.

One of the first attempts at a systematic study of the soils of the United States was initiated in 1895 by Wiley, Moore, and Ewell, of the Division of Chemistry of the United States Department of Agriculture (23x). Forty-four soils, some virgin and some cultivated, were collected from widely separated localities and brought to Washington to be used in greenhouse-pot tests. Oats and buckwheat in one series of tests and beans and buckwheat in another were the rotations used each year over a period of about 8 years. The crops were grown under uniform conditions, except for the soils used, and the results (unpublished) show wide variations in the nitrogen, potassium, and phosphorus contents of these crops. Thus the potassium content of the oat plant varied from 1.23 to 3.34 percent of the dry weight, depending on the soil in which it was grown, while the phosphorus varied from 0.15 to 0.52 percent in the same series. For example, the data show that all crops grown in a virgin sandy loam from Massachusetts had very low contents of both potassium and phosphorus, while those grown in an adobe soil from California were high in both of these elements. Of course these pot tests do not reflect local conditions such as the natural moisture content of the soil, climatic factors, and other environmental characteristics found in the field; but they do indicate some fundamental differences in soils and their ability to supply, under ideal conditions, those elements essential to the health and well-being of the animal consuming the products of that soil.

Other work of Moore's includes pot and field tests made in 1903 of soils in the northern part of the United States (415) and an unpublished report of a field investigation of about 5 different soil types in 26 localities in Kentucky (16x). The data from the latter report show differences up to 150 percent in the phosphorus and potassium contents of the hemp plant, and even greater differences in magnesium

contents. It is interesting to note that the highest phosphorus contents were found in hemp grown in what Moore's descriptions indicate to be the Maury silt loam, a soil derived from phosphatic limestone.

As early as 1909, Russian agriculturalists were using soil characteristics as units in the study of plant composition. Kossowitch (320) determined the absorption of phosphorus by the grain and straw of flax, oats, and mustard and by red clover and hays. He reported, for example, that a greater quantity of phosphorus was absorbed by the mustard grown in a Chernozem than in any other soil.

Ames (7) in 1910 analyzed several wheat crops grown in different soils from the fertility plots at the Ohio Agricultural Experiment Station. His data indicate little or no difference in the phosphorus contents of either grain or straw grown in soils at Wooster and at Strongsville, but the potassium in the straw from Strongsville is nearly 50 percent higher than that in the Wooster soil, and this difference is in agreement with the values reported for potassium soluble in N/5 HNO₃ in each of the soils. No difference in available phosphorus was found, which is also in accord with the similar phosphorus content of the plants.

In their study of the composition of bluegrass in different parts of Ohio in 1910, Forbes, Whittier, and Collison (195) observed that

while the mineral content of vegetable crops is without doubt the resultant of a considerable number of varying factors, the most important of these is the composition of the soil. While it seems unlikely that the variations in soils would cause acute or immediately perceptible effects on animals in Ohio, it seems entirely probable that in the course of the lifetime of animals they do cause, through a gradual molding of the growth, differences of considerable practical importance.

They reported that samples of bluegrass (collected when first coming into bloom) taken from some pastures in Ohio contained more than twice as much phosphorus or calcium as did those from other pastures. They also observed that while the inherent fertility of the soil was important, the use of manure and other fertilizers on poor soils often resulted in bluegrass rich in mineral nutrients.

Pot tests similar to those of Wiley and Moore were conducted by Lemmermann, Einecke, and Fischer in 1911, using German soils (344). They grew oats, barley, rye, and wheat, and in every case the straw from the plant grown in a sandy soil from near Petkus contained the lowest amount of calcium. This was not typical of sandy soils, however, for another soil of this type produced crops that compared favorably with loamy sands and sandy soils of very high humus content. The differences in plant composition were distinctly not related to soil structure.

Gile (208) in 1911 reported that the ash of plants grown in calcareous soils differs from the ash of plants grown in noncalcareous soils chiefly in containing a larger amount of calcium and a smaller amount of iron. Chlorotic pineapple plants were produced in Puerto Rican soils to which limestone had been added, and the iron content in these plants was less than half that in plants grown in the more acid soils.

Cohen (116), investigating in 1914 the composition of tobacco grown in Javanese soils, found some relationship between available phosphorus content of the soil and that of the plant, but none between the potassium of the soil and plant. In general the heavier soils, poor in citric acid-soluble phosphorus, produced a tobacco with a lower

phosphorus content than tobaccos grown in light soils. Some relationship was also established between the "available" lime in the soils and the calcium content of the plant.

Robinson, Steinkoenig, and Miller (495) in 1917 made a number of analyses of plants collected on different soils. Their analytical work is the most complete that has ever been published in this field, but, unfortunately, comparisons of the different samples cannot always be made in order to determine differences in soil properties, because the stage of growth of the plants differs. Specific references to most of their analyses are given in the appendix, and an examination of their data reveals interesting variations in some of the minor element contents of the plants studied. Thus, the manganese content of wheat grown in the Durham soil is very high as compared with that of wheat grown in the Penn silt loam, but the manganese content of the Durham soil is about one-eighth of that of the Penn soil. The magnesium content of the red clover grown in a Penn silt loam was higher than that of the clover grown in a Hagerstown loam, although the relative magnesium contents of the two soils are the reverse of that of the plants. Similar results are shown for the manganese, strontium, and aluminum contents of this plant. Both soils are classified as gray-brown podzolic soils, but the Penn silt loam is developed from shale while the Hagerstown loam is developed from limestone. It is evident from these data that no direct relationship necessarily exists between any single characteristic of the soil and any characteristic of the plant.

These authors also studied the occurrence of a number of unusual elements such as rubidium, which was not detected at all in the wheat plant from a Penn silt loam, whereas 2.7 p. p. m. was found in the plant from the Durham loam. In other samples of vegetation these authors found caesium in plants grown in two soils known to contain caesium beryls, whereas they reported that most of the vegetation did not contain this element. Chromium was occasionally found, though in very small amounts. Vanadium was detected in only six instances, and then only in traces.

Maschhaupt (397) determined the composition of summer barley, field beets, oats, two varieties of winter barley, potatoes, and rye grown in rotation in several Dutch soils from 1909 to 1922. His soils, transported from their original environment, included peat soils, loams, and clays and lay adjacent to one another but separated by concrete walls. He reported that the influence of soil type on the content of nitrogen and the ash constituents was greater on grain straw, potato leaves and beet leaves than on the grain itself, beet roots, or potato tubers. The silica content of the straws from the clay soil were nearly five times as high as those of the straws grown in a "heidebode," and significant variations were noted in the phosphorus contents of potatoes and the calcium contents of beets grown in different soils.

Carefully sampled and prepared specimens of crabgrass (*Digitaria sanguinalis* (L.) Scop.) obtained from a garden soil and from a limestone roadway were analyzed by Buckner (96) in 1919. The samples grown on the roadway soil contained about 50 percent more calcium and more phosphorus and magnesium in the ash than did the other samples, but calculated on the dry basis rather than on the ash, the

difference in composition between the two samples is negligible, which illustrates the possibility of misjudging the composition of a plant when it is calculated as percentage of the ash.

Headden (256) in 1921 determined titanium, barium, strontium, and lithium in alfalfa grown in several soils of Colorado. He found over 20 times as much titanium in plants grown in one soil as in those grown in another, and variations of 200 to 1,000 percent in strontium were found. The barium contents were more uniform, and, except in one case, only traces of lithium were found.

In 1923, Neidig, McDole, and Magnuson (430) determined the sulfur and nitrogen contents of alfalfa grown in greenhouse tests in six different soils from localities in Idaho. The sulfur contents varied from 0.140 to 0.300 percent.

In some unpublished work done on fertility problems at the New York State (Cornell) Agricultural Experiment Station (2x), samples of three soils—Dunkirk silty clay loam, Volusia silt loam, and Petoskey gritty sandy loam—were placed in lysimeter tanks. Corn, oats, barley, wheat, and timothy were grown in the soils, and the crops were analyzed for potassium, calcium, magnesium, phosphorus, and sulfur. The data obtained for 3 years on corn stover grown in two of these soils are presented in table 9. The data indicate little or no difference in the potassium and phosphorus contents of the two crops, but there is a significant difference in the magnesium contents of the stovers grown in the unfertilized soils. Furthermore, the addition of manure modified the magnesium content of the stover grown in the Dunkirk silty clay loam to a greater degree than it did that of the crop grown in the other soil. It is equally possible, however, that these relative differences in the crop composition are due to differences in the climatic effects on the soil, for the various fertilizer treatments were studied in different years.

TABLE 9.—Mineral composition of corn stover¹ grown in New York soils in lysimeter tanks (2x)

Year	Fertilizer	Composition of stover grown in—							
		Dunkirk silty clay loam				Volusia silt loam			
		K	Ca	Mg	P	K	Ca	Mg	P
1915.....	None.....	Percent 1.39	Percent 0.20	Percent 0.01	Percent 0.27	Percent 1.19	Percent 0.15	Percent 0.08	Percent 0.22
1920.....	Manure.....	1.39	.30	.21	.28	1.76	.10	.12	.42
1925.....	Manure and Na ₂ HPO ₄77	.21	.16	.37	.82	.24	.22	.32

¹ Moisture-free basis.

Neller, investigating the influence of sulfur on the yield and composition of legumes (432), used two Washington soils in his greenhouse tests in 1925. His analyses of alfalfa cut in the early blossoming period are given in table 10. The sulfur content of the Ritzville soil

was found to be 0.020 percent, as compared with 0.043 percent for the Sagemoor soil, and this difference seems to correlate very well with the sulfur content of the vegetation.

TABLE 10.—*Mineral composition of alfalfa*¹ (average of duplicate experiments grown in pots without fertilizer treatments)

[Neller (432)]

Soil	Yield	Ash	N	K	Ca	P	S
	<i>Gm.</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
Ritzville silt loam.....	31.8	8.30	1.64	3.13	1.38	0.230	0.120
Sagemoor very fine sandy loam.....	25.1	-----	2.57	2.32	1.92	.139	.242

¹ Moisture-free basis.

Meinck (406), in studying the relationships between the iodine content of the soil, water, and foods in goitrous regions in Germany, reported in 1927 little correlation between the iodine content of potatoes and either the total or hydrochloric acid soluble iodine in the soil. Potatoes grown at Schwalbach in a soil having 168 p. p. b. of iodine soluble in acid contained more than three times as much iodine as did potatoes grown at Mammolsheim in a soil having 502 p. p. b. of soluble iodine. The total iodine in the soil in both cases was nearly the same.

The variations in the composition of soybean forage grown in six Michigan soils were determined by Austin (37) in 1928. The plots, located in different parts of the State, were all planted and harvested at the same time, so that the age factor was eliminated. In determining the composition of the plants the entire above-ground portion of the plant was used for analysis. The data obtained on samples from the unfertilized plots are presented in table 11. A study of the data indicates that some of the constituents vary to a marked extent, and that certain trends during the whole season are observable. The calcium content of the plants grown in the Hillsdale soil is without exception higher than the average throughout the season, while that of the plants grown in the Kewanee is below average throughout the season. The average seasonal value for calcium in the plants grown in the Hillsdale is 2.206 percent, as compared with 1.729 percent for that grown in the Kewanee. The magnesium content of the plants grown in the Brookston soil is in some cases nearly double that of plants grown in other soils, and it remains high throughout the season. It is apparent that the plants grown in the Kewanee soil are consistently high in potassium throughout the season as compared with those grown in other soils. The plants grown in the Miami are likewise low in phosphorus, while those grown in the Brookston and Coloma soils are consistently above the average in this respect. The differences (0.278 compared with 0.411 and 0.425 percent respectively) are important.

TABLE 11.—*Mineral composition of soybean plants (above-ground portion) ¹ grown in Michigan*

[Austin (37)]

Stage of growth and soil type on which grown	K	Ca	Mg	P	S
35 days after seeding (July 13):	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
Hillsdale sandy loam.....	0.695	2.702	0.586	0.234	0.232
Brookston clay loam.....	.635	2.162	1.083	.469	.289
Coloma sand.....	.504	2.799	.694	.411	.318
Kewanee loam.....	.812	2.081	.663	.345	.275
Miami loam.....	.663	2.370	.832	.312	.258
Average.....	.662	2.423	.772	.354	.274
73 days after seeding (Aug. 19):					
Hillsdale sandy loam.....	.722	2.050	.846	.287	.283
Brookston clay loam.....	.627	1.792	1.086	.374	.291
Coloma sand.....	.729	1.760	.644	.403	.321
Kewanee loam.....	.623	1.813	.643	.383	.236
Miami loam.....	.774	1.834	.773	.242	.290
Fox sandy loam.....	.511	1.894	.665	.377	.347
Average.....	.664	1.857	.776	.344	.290
110 days after seeding (Sept. 26):					
Hillsdale sandy loam.....	.576	1.865	.719	.329	.249
Brookston clay loam.....	.713	1.681	.795	.390	.326
Coloma sand.....	.591	2.115	.607	.462	.286
Kewanee loam.....	.738	1.294	.536	.423	.277
Miami loam.....	.532	1.291	.637	.280	.229
Average.....	.630	1.649	.659	.377	.273
Averages for each soil:					
Hillsdale sandy loam.....	.664	2.206	.717	.283	.255
Brookston clay loam.....	.658	1.878	.988	.411	.302
Coloma sand.....	.608	2.225	.648	.425	.308
Kewanee loam.....	.724	1.729	.614	.384	.263
Miami loam.....	.656	1.832	.747	.278	.249

¹ Moisture-free basis.

Fonder in 1929 investigated the composition of the leaves and stems of several plants grown in different soil types in different areas in Michigan, and he attempted to correlate his data with the composition of the soil solutions (188, 189, 190, 191). Some of his data are presented in table 12, because they show how the mineral composition of one part of the plant may vary when the plants are grown in different soils. The data for alfalfa leaves only are given here, but the discussion would also apply to the composition of the stems of alfalfa or of leaves and stems of the green bean plant or pea plant which Fonder also studied. The data show that alfalfa leaves grown in the Plainfield loamy sand contain less calcium than the average throughout the growing season, while those growing in the Fox sandy loam contain more than the average. The average calcium throughout the season in the plant leaves growing in the Plainfield soil is 1.648 percent, and that of the plant leaves growing in the Fox soil is 2.609 percent, a significant difference. The magnesium in the plant leaves grown in the Brookstons loam is low throughout the season, in contrast with the data of Austin (table 11), while that in plant leaves grown in the Fox sandy loam (table 12) is high. The potassium data are not so consistent throughout the season, but well-defined trends are observable. The data show also that calcium accumulates

at a rapid rate in the leaves, in contrast with potassium, which decreases appreciably, while magnesium increases more slowly throughout the growing season.

TABLE 12.—Variations in the potassium, calcium, and magnesium content of alfalfa leaves¹ grown in different soil types

[Fonder (188, 189)]

Stage of growth and soil type on which grown	K	Ca	Mg
	Percent	Percent	Percent
May 8:			
Plainfield loamy sand.....	4.63	1.103	0.256
Coloma loamy sand.....	2.98	1.211	.224
Hillsdale sandy loam.....	2.70	1.182	.226
Fox sandy loam.....	2.22	1.571	.268
Conover loam.....	2.40	1.353	.313
Brookston loam.....	3.02	1.161	.241
Miami silt loam.....	1.62	1.468	.265
Average.....	2.80	1.293	.256
May 22:			
Plainfield loamy sand.....	1.69	0.826	.285
Coloma loamy sand.....	1.82	1.683	.309
Hillsdale sandy loam.....	2.18	2.120	.244
Fox sandy loam.....	1.90	1.737	.297
Conover loam.....	1.64	1.710	.250
Brookston loam.....	2.05	1.748	.203
Miami silt loam.....	1.50	1.917	.284
Average.....	1.83	1.677	.267
June 7:			
Plainfield loamy sand.....	3.02	2.140	.258
Coloma loamy sand.....	2.02	1.695	.199
Hillsdale sandy loam.....	1.56	2.339	.241
Fox sandy loam.....	1.22	3.108	.397
Conover loam.....	.67	3.830	.444
Brookston loam.....	1.14	2.568	.263
Miami silt loam.....	.88	2.148	.262
Average.....	1.50	2.547	.295
July 2:			
Plainfield loamy sand.....	2.26	2.522	.337
Coloma loamy sand.....	2.13	2.930	.321
Hillsdale sandy loam.....	1.34	4.060	.330
Fox sandy loam.....	.95	4.020	.587
Brookston loam.....	1.39	2.250	.263
Miami silt loam.....	2.20	2.610	.298
Average.....	1.71	3.065	.356
Averages for each soil:			
Plainfield loamy sand.....	2.90	1.648	.284
Coloma loamy sand.....	2.24	1.880	.263
Hillsdale sandy loam.....	1.94	2.425	.260
Fox sandy loam.....	1.57	2.609	.387
Conover loam.....	1.57	2.298	.336
Brookston loam.....	1.90	1.932	.242
Miami silt loam.....	1.55	2.036	.277
Average.....	1.95	2.118	.293

¹ Moisture-free basis.

In his investigation of the soil solution, Fonder reported that although the amounts of calcium and magnesium present in the several soil solutions varied greatly at the beginning of growth, by the time the plant had reached the mature stage the amounts had been reduced until there was but little difference among the soil types. The reduction in the concentration of the two elements in the solution was not rapid and was very uniform, the soil types maintaining the same order with respect to each other during most of the growth period. He concludes: "In general, the calcium contents of the

plants grown in two soils of like or nearly the same degree of acidity and of similar texture vary with the calcium content of the soil solution." The data in table 13 illustrate this point.

TABLE 13.—Relation between the calcium content of the leaves of the pea plant¹ and that of the soil solution

[Fonder (191)]

Soil	pH of soil	Ca in soil solution	Ca content of leaves after—				
			3 weeks	6 weeks	8 weeks	Budding	Maturity
Light soils:		<i>P. p. m.</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
Onaway	7.40	24.55	0.259	0.286	0.260	0.462	0.536
Plainfield	7.35	10.60	.193	.285	.285	.379	.384
Kewanee	6.25	44.70	.196	.359	.274	.292	.418
Hillsdale	5.10	14.62	.185	.332	.255	.292	.282
Heavy soils:							
Miami	5.00	30.06	.197	.384	.278	-----	.582
Brookston	7.05	19.00	.193	.293	.336	-----	.436

¹ Moisture-free basis.

Brown (91, 92) analyzed apples grown in different soils in England during two seasons (1926 and 1929). Her results show that the seasonal variations are comparatively small, whereas quite large and significant differences occurred in those fruits grown in different soils. Differences in the calcium and phosphorus contents were particularly noticeable.

Increasing interest in soil-plant relationships has led to considerably more active investigations in this field since 1930 than had been evidenced during any other period. The investigations during this time will be taken up in their geographical order, so that a clear picture can be had of the work on any group of soils.

Coleman and Ruprecht (119, 5x) in 1935 analyzed tomatoes, celery, potatoes, oranges, grapefruit, string beans, cabbage, and lettuce grown in different soils in Florida. Their data, included in the compilation in the appendix, show clearly the influence of soils on the calcium content of the plant. Thus, the calcium content of tomatoes grown in the Bladen soil, a strongly acid soil with a small amount of organic matter, was reported as 0.22 percent, while that of tomatoes grown in the Hernando soil, developed from a phosphatic limestone, was 0.45 percent. Similar variations were reported for cabbage and other vegetables.

Bryan and Neal (95) in 1936 reported slight variations in the phosphorus contents of vetch and sorghum grown in three Florida soils in pots. The variations, although consistent in both crops, are not large. Thus, the vetch grown in Norfolk sand contained 0.114 percent of phosphorus, in Norfolk fine sand 0.092 percent, and in Orangeburg fine sand 0.087 percent.

Bishop (71) grew a number of vegetables in some Alabama soils in the greenhouse, and her analyses, published in 1934, show enormous variations in the calcium and phosphorus contents of the edible portions of these crops. Thus, cabbage grown in a Eutaw soil contained 0.208 percent of phosphorus, while that grown in a Hartsell soil contained only 0.086 percent. In general, the calcium contents of the vegetables were found to be the lowest in those grown in the Norfolk

soil, while the phosphorus contents were consistently the lowest in those grown in the Cecil soil. Bishop's data also show that the calcium and phosphorus contents vary inversely, so that the calcium-phosphorus ratio fluctuates even more markedly.

Weathers' (580) recent analyses of lespedezas from three soils of varying fertility in Tennessee show quite clearly the effect of soil and locality as contrasted with that of variety. The data tabulated in the appendix show that the hay grown in the Maury silt loam, a gray-brown podzolic soil developed from a phosphatic limestone, contains the highest quantity of calcium and phosphorus, while that grown in the Clarksville loam contains the least quantity. The phosphorus content of the lespedeza grown in the Clarksville loam, a nonphosphatic soil, is less than half that grown in either the Maury or the Cumberland soils, and there is a corresponding disturbance in the calcium-phosphorus ratio.

Pierre (17x) analyzed a number of grass, clover, and alfalfa samples growing in association in old established pastures in West Virginia, and his data for alfalfa are presented in table 14. The collections of grass and clover samples were not made at any stated stage of maturity nor at any uniform time, and they are not, therefore, comparable. Likewise, his soils were chosen in different localities, so that rainfall, management, and other factors might have influenced the composition of the plant. These data indicate that throughout the season all cuttings of alfalfa grown in the Hagerstown soil were lowest in phosphorus as well as calcium. This is in relative agreement with the available phosphorus in the soils, although the results as between DeKalb and Hagerstown are contradictory to what would be expected unless amendments were used liberally on the former.

TABLE 14.—*Calcium and phosphorus in alfalfa*¹ *grown in different soil types in West Virginia*

[Pierre (17x)]

Cutting	Ca			P		
	DeKalb silt loam ²	Holston silt loam ³	Hagerstown silt loam ⁴	DeKalb silt loam	Holston silt loam	Hagerstown silt loam
	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
First.....	2.80	3.22	1.92	0.184	0.166	0.139
Second.....	1.85	1.98	1.43	.324	.245	.171
Third.....	2.18	2.15	1.79	.268	.240	.195
Average.....	2.28	2.45	1.71	.259	.217	.168

¹ Harvested at one-third to one-half bloom. Moisture-free basis.

² Available phosphorus, 16 p. p. m.; pH, 6.83.

³ pH, 7.30.

⁴ Available phosphorus, 9 p. p. m.; pH, 6.20.

Magistad (15x) analyzed five varieties of grass, each grown in four different soil types in Hawaii. His data (unpublished) are very consistent in indicating definite relationships between the calcium and phosphorus contents of all varieties and each of the soil types. Thus, all varieties grown in certain soils are low in these constituents, while those grown in other soils are high. The predominance of soil over variety is quite marked in these experiments.

Hester and coworkers made a number of fundamental studies on three soils occurring in Virginia, a Portsmouth loamy fine sand, a Bladen sandy loam, and a Norfolk loam (266, 267, 268, 269). They found significant differences in the power of plants (greenhouse tests) to absorb the nutrient from each of the soils because of the differences in the exchange capacity of the soils for the nutrients. The Bladen soil, for example, was found to possess a high power for fixing potash in a nonreplaceable state, whereas plant absorption of this element from the Portsmouth is high. The utilization of phosphorus from the different soils by the plants was in the descending order of Portsmouth, Bladen, Norfolk, the reverse of the phosphorus fixation capacity of these soils. These authors conclude that

The presence of a large amount of a particular replaceable base in the soil colloidal complex influenced the elemental composition of a plant material even though the yields were affected but little. A high replaceable calcium content suppressed the absorption of potassium, nitrogen and magnesium; a high replaceable magnesium content suppressed the absorption of potassium, calcium, and nitrogen; and a high potassium content suppressed the absorption of calcium, magnesium, and nitrogen.

Garner, McMurtrey, Bowling, and Moss (201) found that the magnesium content of tobacco leaves was closely correlated with the occurrence of magnesium deficiency diseases and with the particular soils of Maryland, Connecticut, and North Carolina in which the plants grew.

The influence of soils on the composition of several vegetables grown under different soil conditions in Maryland and in Virginia was studied by Davidson and Le Clerc (140) in 1936. Their data, compiled in the appendix, do not indicate many consistent differences in the mineral composition of the crops. However, these soils had been heavily fertilized, and it is recognized that the Norfolk and the Sassafras are both quite responsive to fertilization.

The effect of soil type on the composition of corn stover was studied by Wimer (596) in his recent experiments in Illinois. No significant differences in composition of the stover were found in plants grown in Saybrook silt loam and in the deep phase of this soil. The soils are, however, identical except for variations in depth. In another field in which more significantly different soils occur, differences in the composition were found. In discussing this point the author states:

Drummer clay loam is rich in total nutrients, has a high content of available phosphorus, and requires no limestone to grow sweet clover. This soil yielded 1,514 pounds more stover and 20.2 bushels more grain per acre than Sidell silt loam, which is less fertile, lower in available phosphorus, and more acid in reaction.

The composition of the stover from each soil in this field is given in table 15.

TABLE 15.—*Mineral composition of corn stovers¹ grown in different (untreated) soils in the same field*

[Wimer (596)]

Soil type	N	P	K	Ca
	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
Drummer clay loam.....	0.83	0.115	0.79	0.24
Sidell silt loam.....	.86	.082	.31	.21

¹ Moisture-free basis.

According to the author,

The wide variations in climate in different parts of Illinois have made it very difficult to study the effect of soil types, throughout the area of their occurrence, on the composition of corn stover. Widely different soil types ordinarily do not occur on the same experimental field; therefore, seldom under the same seasonal conditions of rainfall, temperature, and other climatic factors. Because of these and other limitations (particularly variety of plant), comparisons that will show the effect of soil type on the composition of stover are indeed rare.

Grizzard (229) determined the variations in composition of the leaves and the stems of alfalfa plants grown in different soils in different localities of Michigan. Differences of 105 percent of phosphorus and 73 percent of calcium in the leaves of the plants were found. The author states:

The heavy soil types, such as Brookston clay loam, tend to give a higher nitrogen content in the alfalfa plants than the lighter soil types, such as Isabella sandy loam, both when fertilized and unfertilized. On the other hand, the lighter soil types tend to give a decidedly higher phosphorus content in the alfalfa plants than the heavy soil types, this being true both in the fertilized and the unfertilized condition. In the case of calcium, there is a tendency for this element to be decidedly higher in the intermediate soil types, such as the Gilford and Brookston loams, than in either the heavy or light soil types. Apparently, these data indicate that soil type predominates over fertilizer treatments in the composition of alfalfa; although fertilization tends to influence the composition of alfalfa, especially in certain soils.

Some of the factors responsible for the variation in the calcium and phosphorus contents of peas were studied by Peterson and coworkers (4x). A summary of their unpublished data is given in table 16. The data indicate that peas grown in the Knox soil contain less calcium than those grown in the other two, and that variety of peas may influence the absorption of calcium.

TABLE 16.—*Calcium and phosphorus contents of Wisconsin peas*¹

[Peterson and others (4x)]

Variety of pea and soil type	Ca	P	Variety of pea and soil type	Ca	P
Smooth Alaska:	<i>Percent</i>	<i>Percent</i>	Perfection:	<i>Percent</i>	<i>Percent</i>
Miami silt loam.....	0. 263	0. 649	Miami silt loam.....	0. 226	0. 611
Carrington silt loam.....	. 203	. 578	Carrington silt loam.....	. 174	. 640
Knox silt loam.....	. 179	. 582			

¹ Moisture-free basis.

Holtz (275) in 1930 studied the variations in calcium and phosphorus contents of oats and red and white clover grown in different western Washington soils in different localities. He concludes that "the calcium content of oat hay is influenced by the calcium content of the soil in any particular group," that is, residual, glacial or alluvial soils. The composition of oat hay and red clover both follow the available phosphorus in the residual soil, but in the upper glacial soils the composition of oat hay responds to the total phosphorus content of the soil, whereas clover follows the available phosphorus. The results of the experiments were influenced, of course, by climate and other factors as well as by the soil itself. Other investigations of Washington soils include those of Baker and Vandecaveye (44) in 1935 and Vandecaveye and Bond (572) reported in 1936.

Greaves and Anderson (220) reported in 1936 that the copper content of wheat, barley, and oats (grains) varied tremendously as grown in different localities. No correlation was found, however, between the content of copper in the wheat and that of the soil. These investigators also determined the copper content of several varieties of wheat grown in the same soil and concluded "that variety is more important than soil, except where the copper content of the soil falls below a certain minimum." The variations that they found on different soils are in direct contradiction to some unpublished work of Bailey and Hutchinson (1x), who found practically no variation in the copper content of wheats grown over a large part of the northwestern wheat country.

The sulfur contents of two samples of alfalfa grown in two localities in Utah were not significantly different, according to Evans and Greaves (169). No descriptions of the soils were given, but the sulfur contents of the soils were quite different.

A recent comprehensive survey of New Mexico grasses by Watkins (579) indicated that there were 12 counties in New Mexico in which the average phosphorus content of the range grasses for the year 1932 was insufficient for normal growth and reproduction of range cattle. However, the grasses were collected late in the fall and early in the spring, so that the samples might have been leached, and undoubtedly they did not represent whole plants of any particular stage of maturity.

The effect of soil type on the composition of clover was studied by Myers and Metzger (425) in Kansas, but no indication is given that the samples chosen were consistent in stage of maturity.

Poehlman (469), investigating in 1935 the analyses of plant juices as indicators of the nutrient needs, found that

the phosphorus and potassium concentration in the plant juice differs significantly with the different soils [in Missouri] upon which the plants were grown and the soil type is an important factor in determining the concentration of these elements in the plant juice.

He states, however, that uncontrolled factors in his experiments include "variations due to age or maturity of the plant, moisture content of the plant, rate of growth, soil heterogeneity, and climatic factors such as light, temperature, rainfall, humidity and soil moisture."

Daniel and Harper (134), in studying the composition of prairie grasses in Oklahoma in 1934, found that the moisture relationships in the soil were very important with respect to the phosphorus content of the plant. Thus, a sample of grass (*Andropogon furcatus* Muhl. and *A. scoparius* Michx.) collected from an alluvial soil where moisture conditions were favorable during the growing season contained more than twice as much phosphorus as usually occurs in the average sample of upland prairie hay.

Variations in the nutrient content of soils have less effect on the seed than on any other part of the plant. A comparative analysis of Chinese and American rices by Davidson and Chambliss (139) in 1932 did not "indicate any striking differences in composition between the Chinese rices, grown on a soil presumably cropped for thousands of years, and the rices grown under the American method of cropping."

Lincoln (354) studied differences in soil and plant relationships by examining samples of *Stenotaphrum dimidiatum* Bronger, growing in Mauritius pastures. Samples collected at the same stage of growth varied considerably in composition, depending upon the character of

the soil. Thus, the calcium content of this grass growing in red soil averaged 0.38 percent as compared with an average of 0.67 percent for that grown in calcareous soils and 0.54 percent for that grown in the dark-brown soils.

Depardon (152) in 1938 reported on the effect of soil, climate, and geology of three regions in France on the composition and quality of wheat, forage beets, and potatoes. He concluded that the predominant factor in the production of plants was distinctly the soil. Thus in every case studied, the phosphorus content of plants grown in Perche was lower than that in plants from either of two other localities.

A few studies have been made that were concerned with the more unusual minor elements in the soil. The most comprehensive study of this nature is that conducted by the Department of Agriculture and some of the State experiment stations on selenium. The importance of selenium in relation to the health of the animal has already been noted, and a brief discussion will be presented concerning the relation between selenium in soils and its occurrence in plants growing in the soils.

Byers and Knight (103) reported in 1935 on the selenium content of soils and two species of grass. All of the soils and grasses contained selenium, and variations occurred in the amounts, but the greatest variations by far were between the different species of grasses.

Beath and coworkers (48) published the selenium contents of a number of crops grown in certain shales in Montana. Their data seem to indicate that certain species, such as alfalfa, do not absorb selenium, and that crops grown in soils developed from the Niobrara shales contain less selenium than do those grown in soils from Steele shales.

Byers, in a series of reports on the occurrence of selenium-bearing soils in South Dakota, Colorado, Kansas, and Montana (100, 101, 104), states:

The distribution of selenium in soils appears to be general. No true soils containing colloids in any significant quantity have been found in which the presence of selenium cannot be demonstrated. The source of the selenium is believed to be the residual selenium derived from the soil parent material, supplemented by that derived from direct absorption from the air by rain. It has been demonstrated that selenium may exist in soils as the element; as a substituent in sulfide minerals, particularly pyrites; as selenite, particularly basic ferric selenite; as selenate, particularly as calcium selenate; and as organic selenium compounds of undemonstrated composition. Of these forms apparently those most available to the soil solution are the organic and selenate forms.

The distribution of selenium within the soil profile is extremely variable, and definite general statements are not possible concerning these variations.

When plants grow in seleniferous soil they seemingly absorb selenium to some degree in all cases, though under many circumstances the quantity so absorbed is vanishingly small. For a given species [of plant] the quantity found by analysis appears to depend upon a number of variables. These include at least the following: The quantity and character of the selenium in the soil [(282)] and its distribution in the soil; the portion of the plant examined, whether seed, blossom, leaves, stem or root; the stage of maturity of the plant; and seasonable variations which might be attributed to rainfall. It is also certain that variations in selenium absorption are brought about by variation of available sulfur [(285)].

In a later publication (331), Lakin, Williams, and Byers state: "A lack of quantitative correlation between the selenium content of the soil and the vegetation has been evident throughout the entire investigation of seleniferous areas." They reported, for example, that the Yunes silt loam of Hawaii contains from 2 to 10 p. m. of selenium

in the surface horizon, while a soil in Teton County, Mont., contains only 0.2 p. p. m. Examination of the vegetation from the Hawaiian soil showed that none contained more than 1 p. p. m., while *Astragalus pectinatus* growing in the Montana soil contained 190 p. p. m. of selenium.

Boron has been studied quite extensively, both because of its injurious effect and its essentiality for plant growth. The boron contents of leaves of citrus trees suffering from a disease was shown by Kelley and Brown (312) in 1928 to be several hundred times as high as those of healthy trees. They stated in this connection:

The results of this investigation strongly indicate that boron is readily absorbed by citrus and walnut trees and that this element tends to accumulate in the leaves of these species. Because of this fact, a determination of the boron content of the leaves affords a valuable indication as to the boron conditions of the soil.

They concluded that irrigation water is responsible for some difficulties with boron, and Eaton (160) in 1935 supports this when he states:

Notwithstanding evidence of the occasional occurrence of appreciable concentrations of boron in unirrigated soils in the San Joaquin Valley, the general inference to be drawn from the appearance of crops is that boron carried by irrigation water is for the most part responsible for such depression in yield and such crop injury as have occurred.

Eaton further concludes that "much of the boron found in soils, whether irrigated or not, is not very soluble."

Woodbridge (598) in 1937 found that "a low boron concentration in tissues can be correlated with high incidence of disease [of apples]. A correlation between low boron in the soil and incidence of disease was not established."

Definite relationships between manganese in the soil and that in the plant are few. Wester (586) in 1923 determined manganese in several soils in the Netherlands along with that in digitalis leaves growing in the soils. His data are given in table 17.

TABLE 17.—*Manganese in some Netherland soils and in digitalis leaves growing thereon*

[Wester (586)]

Material	Mn per 100 gm. of dry matter in—		
	Noordwyk	Delft	Haag
Soil.....	2.4	14.4	10
Digitalis leaves.....	.94	1.32	2.57

Carlyle (106) in 1931 determined the manganese contents of several plants grown in pots containing different Texas soils. He reported differences of 200 to 500 percent in the manganese content, according to the soil used.

Leeper (340) stated that "manganese deficiency disease is confined to soils of pH 6.7 or more and occurs especially on heavily limed podzols."

Iodine has been extensively discussed in relation to the occurrence of goiter, but it is interesting to note that McHargue (377) in 1933 reported 250,000 p. p. b. of iodine in one sample of corn, whereas 18 other samples from other soils in Kentucky contained from 98 to 650

p. p. b. Bishop and Lawrenz (72) reported that they detected cobalt in different species of plants, as well as in some species grown in certain soils. Robinson, Whetstone, and Scribner (497) found most of the rare earths in hickory leaves taken from a tree growing in a pegmatite vein of the Moorefield Mine, Amelia, Va. Later unpublished work by Robinson (18x) seems to indicate that this group of elements may normally be present in the hickory tree wherever it may be found. Dingwall, McKibbin, and Beans (155) reported, in 1934, that molybdenum was found in plants grown on one farm, while none was found in those grown in any other soils. The molybdenum, they state, "may have been present in the parent rock or it may have been carried down with drainage waters." Dingwall and Beans (154) also reported the presence of chromium in some plants grown in certain localities. Nemeč, Babicka, and Oborsky (433) recently claimed to have found considerable quantities of gold in certain plants growing in gold-bearing soil in Czechoslovakia. Their data indicate that *Equisetum palustre* L. accumulated gold to the extent of 610 gm. per ton of plant ash.

EFFECT OF SOILS ON BOTANICAL COMPOSITION OF HERBAGE OF PASTURES

In 1873, Müller (420) reasoned that pastures grazed by animals suffering from nutritional diseases must, either positively through poisoning or indirectly through mineral deficiencies, adversely affect the animal. A botanical analysis of some pastures revealed no great difference in the amount or quality of the different forage plants in normal and abnormal pastures, but in other cases an unusual number of undesirable plants such as "sauergräsern" and marsh plants were found in abnormal pastures, whereas sweetgrasses and clovers prevailed in the better pastures. Similar observations were made by Lawes and Gilbert (333) in 1900 and by Armstrong (14) in England, who found that white clover, ryegrass, crested dogstail, and fiorin (*Agrostis stolonifera* L.) were most abundant on the best grazing lands. The herbage of the inferior types of grassland in the same districts consisted very largely of bentgrass (*A. vulgaris* With.) and various weeds, while white clover and ryegrass were present in comparatively small quantities. Woodman, Blunt, and Stewart (599) noted seasonal variations in the botanical composition of pastures.

Barnes (47), in 1911, found that treating Ohio pastures with limestone and acid phosphate was accompanied first by a marked increase in percentage of legumes. This was later followed by a renewal of the bluegrass sod, which, however, did not reach its optimum until at least the end of the fourth summer after treatment. Brown (89, 90) reported in 1933 that fertilization with phosphorus, lime, and potassium, as compared with other systems of manuring, resulted in a maximum increase of white clover in Connecticut pastures.

Robinson (492), in his recent comprehensive investigations of pasture flora and soil fertility in West Virginia, concluded that

A comparison of samples from the same pastures shows that, in general, soils supporting Kentucky bluegrass are higher in pH, percentage base saturation, total exchangeable bases, available phosphorus, total nitrogen, and exchangeable potassium, than are soils supporting *Danthonia*. Base exchange and available phosphorus are the limiting factors, however. In the Hagerstown soil there is a high correlation between the percentage of Kentucky bluegrass plus white clover in the pasture and the amount of phosphorus in the soil.

SUMMARY OF EFFECTS OF SOILS ON PLANT COMPOSITION

Investigations designed to study the effect of the single factor, soils, on the composition of plants are rare indeed, which is unusual when one realizes for what length of time the importance of this one factor has been understood. Probably not more than one or two field experiments and a small number of greenhouse experiments may really be considered to have demonstrated that differences in soils alone are responsible for differences in such constituents as calcium and phosphorus in the plant.

It is recognized, of course, that there are two approaches to any study of the capabilities of soils to supply the proper amount of minerals to plants. First, a field study of the soil in its natural environment over a period of several years will indicate how that soil, in the sense that the soil is a summation of its environmental factors, in relation to all other soils in that or other localities, will affect plant composition. In such studies the elimination of climatic and physical differences is not desirable. The advantage of such a study is that information is obtained as to the properties of the one soil being studied in relation to all other soils. The disadvantage of such a study lies in the fact that since several variables are not accounted for or eliminated, no predictions based on any one or several factors may be made concerning the behavior of this one soil type in another locality, even though it is recognized that over a long period of time the climate will not differ greatly on different areas of the same soil type. Thus, if a plant grown on a certain soil be deficient in cobalt, that deficiency may be due, among other causes, to an inherent deficiency or unavailability of this element in the soil, or it may be due to climatic factors. If the first factor be the cause, one would naturally suspect that that soil type would be deficient in cobalt wherever found. If the latter factor be the cause, one would expect that, over relatively short periods of time, this soil type in some localities would support plants with a normal quantity of cobalt. If one wished, therefore, to determine the effect of soil alone on plant composition, that soil must be compared with others under exactly the same conditions of climate, relief, and management practices.

Climate and soils are, moreover, so interrelated that greenhouse tests or other methods designed to bring soils together under the same conditions cannot be depended upon to give accurate information as to the ability of the soil in situ to supply the proper amounts of any nutrients required for the development and health of the plant and the animal or human consuming it. It must be recognized, however, that in spite of their limitations, translocation tests of soils, whether in greenhouse or open air, are the only means of differentiating between effect of soil and climate, unless the soils are situated in such a way as to eliminate climatic and other factors. While such tests are of limited value in determining differences in soil, they are of much greater value in studying the means of modifying with amendments soils that fail to support plants of the highest nutritive value.

There is an abundance of evidence that plants and their parts vary greatly and significantly in the principal elements when grown on different soils in different environments, or even in environments that differ only in small degree.

Conclusive evidence is to be found in the literature to prove that significant differences occur in the minor or trace element content of the plant or its part, that these differences may be directly attributed to differences in soils, and that they are responsible for nutritional disorders. Thus, the selenium content of a plant may be related to the amount of selenium occurring in a soil; but enormous differences are also observed that are the result of (1) differences of distribution of the selenium in the profile, (2) differences of availability of the selenium compounds that occur in the soil, (3) the presence of inhibitory agents such as certain compounds of sulfur, or (4) the water relationships in the soil. Information of such complete nature is available for no other element except selenium, but the sporadic deficiencies or occurrence of other trace elements such as boron, iodine, chromium, and even gold, may be taken as an indication that similar relationships would be found for all elements. This and other investigations emphasize moreover that there is probably no direct relationship between any single characteristic of the soil and any characteristic of the plant or of plant production.

The influence of soils on the botanical composition of flora of pastures has long been known. This is also an important factor in soil genesis, for the character of a soil will vary tremendously, depending upon whether it is developed under a high-calcium grass, a high-sodium grass, or under forests, for example, of beech-maple or of conifers.

Difference of species of pasture herbage should also affect the health of the animal, for, as will be shown in the appendix, these different species vary markedly in their chemical composition.

That a relationship should obtain between the soil taken as a whole and the composition of the plant grown therein is a reasonable assumption, for soils, being the products of their environments, show marked variations because of differences in climate, vegetation, parent material, and relief. Thus, the Podzolic and Lateritic soils are well-leached acid soils generally quite low in bases and phosphorus, while the Chernozem and Chestnut soils are not leached, are high in base constituents, and are nearly neutral in reaction.

Within any soil group, however, there will be variations in fertility and reaction because of differences in parent material or relief. Thus, in the Chernozem soils we may find deficiencies in potash and phosphorus where the parent materials were very low in these elements, and in Kentucky and Tennessee bluegrass regions we find variations from the well-leached acid soils common to the gray-brown Podzolic group. In the rolling or hilly sections of these States, erosion removes the acid and leached materials as rapidly as they are formed. The soil that remains is young and relatively immature, and calcareous soil lies within the reach of the plant roots. These soils often have the additional advantage of being formed from parent materials relatively high in phosphorus, and a very evident result of an abundance of these bone-forming materials is the excellent type of horse developed in those localities.

EFFECT OF FERTILIZERS ON PLANT COMPOSITION

Fertilizer experiments have, for the most part, been concerned with the study of yields, and, in comparison with the number of investigations of this nature, very few have included the problem of modifying the composition of the plant. That the composition of the plant is

modified by the use of fertilizers has been known since the earliest soil fertility studies were made, and, in fact, plant composition has often been used as a guide for determining the nutrient requirements of a soil. One of the first and most comprehensive investigations of this nature ever undertaken is that of Lawes and Gilbert (332) in 1884. They analyzed 92 wheat-grain and the same number of wheat-straw samples collected over a period of years. Every sample was from a crop of known history of growth, as to soil, season, and manuring. Nine different fertilizer treatments were investigated, and much valuable information was obtained concerning yields and crop quality. They concluded, for example, that

the proportion of both potassium and phosphorus in the grain, which if normally ripened is assumed to be a comparatively uniform product, varies very considerably according to season; whilst at any rate as between the grain grown with farmyard manure and that grown without manure, the mean proportion over the sixteen years differs but little. The grain grown under the influence of ammonium salts alone shows not only a wide range in the proportions of both potassium and phosphorus according to season, but much lower mean amount than either with farmyard manure or without manure.

Among other early workers in this field of investigation was Robert Peter (454), who in 1884 reported that calcium sulfate applied to a Kentucky soil caused an increase in the calcium and potassium content of the hemp plant. Anderson (9) in Alabama, Lechartier (334) in France, and Godlewski (213) in Austria were also pioneers in this work of determining the composition of plants grown in fertilized and unfertilized soils. Hills made quite extensive investigations, in 1898, of the composition of potatoes and of corn and corn stover grown in Vermont soils with and without fertilizers (271, 272). His data do not show any marked differences in the composition of the plants when different fertilizer elements or combinations of these elements were applied to either a sandy loam or a clay loam soil. Similar results on potatoes were reported by Davidson (142) in 1898 in connection with his work in Virginia.

NITROGEN, PHOSPHORUS, AND POTASSIUM

It does not seem necessary to consider in detail the great number of experiments that dealt with the change in the nitrogen, phosphorus, and potassium contents of plants when these plants were grown in differently fertilized soils. Few of the investigations included observation on soils, climate, or other pertinent factors, and they are, therefore, of little value in any general interpretation. For the most part, therefore, only those fertility experiments of rather recent date that deal with well-identified soils, and those that deal with the specific effects of each, or definite combinations of the elements, on the mineral composition of the plant, will be discussed.

Nolte (438) reported in 1923 that fertilization with potash, either with or without phosphates, did not materially modify the composition of the grain or straw of barley, field beans, or field bean straw, or beets or beet leaves, when these crops were grown in soils of quite different character in Germany.

Price (472) in 1923 reported a 500-percent increase in the calcium content and a 300-percent increase in the phosphorus content of the first cutting of alfalfa hay when he applied 2 tons of limestone per acre.

The addition of 250 pounds of superphosphate resulted in a further increase in phosphorus.

Austin (37) found that the phosphorus content of the soybean plant grown in the Brookston clay loam was actually reduced when phosphates alone were applied to the soil. In general, however, the effects of fertilizer treatments upon the composition of the plants were small, except possibly that of plants grown in the Coloma sand.

Lagatu and Maume (324, 325, 326, 327, 328, 329, 330) have made a number of fundamental studies in recent years of the requirements of the plant for the mineral nutrients. They describe a method of sampling which consists of taking the two leaves just below the fruiting branch of the plant. They claim that these samples give a true index of the plant's fertilizer requirements, and they made the following observations from their results:

1. The lack of one of the three fertilizer constituents [nitrogen, phosphorus, or potassium] increases the absorption of the other two above the amounts ordinarily absorbed from a medium containing a balanced supply of plant nutrients.

2. In the presence of an incomplete fertilizer the plant fails to absorb from the soil the missing constituent in amounts as great as that from an unfertilized plot.

The work of Lagatu and Maume has been supported by that of Thomas (557, 558, 559, 560), using the potato vine grown in a Pennsylvania soil. He showed that the two observations of Lagatu and Maume were "contrary to Liebig's 'law of the minimum,'" and that whatever effect the absence of an element from a fertilizer has on the yield, "it is not due to a depression of absorption of the other elements, but, on the contrary, to a nutritional lack of balance due to increased absorption of the remaining elements."

The results obtained by Thomas on leaves of potato vines are presented graphically in figure 2, and his summary of the indications given by the graphs follows:

Nitrogen graphs.—The content of nitrogen in the leaves of plants which received mineral nitrogen applications is higher throughout the whole period than in those which received none.

Phosphoric acid graphs.—The content of phosphoric acid in the foliage of plants which received phosphate applications is higher throughout the whole period than in those which received none.

Potash graphs.—The content of potash in the foliage of plants which received mineral potash applications is very much higher than in those which did not.

In his concept of the nutrition of the plant, Thomas makes use of two terms, viz, *quantity* and *quality* of the mineral constituents. "The concept of *quantity* or intensity of nutrition of the selected leaves consists of the sum ($N + P_2O_5 + K_2O$) of each element at the moment of sampling, expressed as a percentage of the dried material. The *quality* of nutrition is the ratio of these entities to each other at the moment of sampling." The quantity of nutrition may be expressed as follows:

$$S = X + Y + Z,$$

where X is the percentage of N , Y the percentage of P_2O_5 , and Z the percentage of K_2O . The quality of nutrition may then be expressed as

$$1 = \frac{X}{S} + \frac{Y}{S} + \frac{Z}{S}.$$

This expression, according to Thomas, gives "the physiological ratios between N , P_2O_5 , and K_2O ." When this proportion of elements is

expressed as milligram equivalents, and the terms of the equation are multiplied by 100 to avoid fractions, Thomas obtains his expression for the NPK unit, and the properties of this expression are such that "for each simultaneous value of N, P, and K there is one point and only one on an equilateral triangle each of whose sides equals 100 [units]."




FIGURE 2.—Influence of different fertilizer treatments on the nitrogen, phosphoric acid, and potash content of potato leaves at different stages of growth. (From Thomas, 560.)

Thomas assumes that a fertilizer may modify the composition of a plant by (1) "an increase in the sum, $(N+P_2O_5+K_2O)$, i. e., in the intensity of nutrition, or (2) a change in the composition of the NPK-unit, or (3) a change in both (1) and (2) simultaneously."

His data, plotted in figure 2, when analyzed on the NPK-unit basis show that compared to a balanced fertilizer:

(1) The omission of nitrogen from the complete fertilizer has increased the potash and phosphoric acid in the NPK-unit of the leaf throughout the whole period, resulting in decreased yields.

(2) The omission of potash from the complete mixture has increased the nitrogen and the phosphoric acid in the NPK-unit, resulting in decreased yields.

(3) The omission of phosphorus has increased the potash in the NPK-unit of the leaf at the expense of the nitrogen, resulting in a slight increase in yield.

(4) The omission of both phosphorus and potash has increased the nitrogen in the NPK-unit, resulting in considerably decreased yields.

(5) The omission of both nitrogen and potash has increased the nitrogen and the phosphorus in the NPK-unit, with considerable reduction in yields.

(6) The omission of both nitrogen and phosphorus has resulted in increasing the potash in the NPK-unit, resulting in decreased yields.

McCool and Weldon (366) in 1928 noted that 36 days after planting, the phosphorus content of the juice of barley plants had increased with both phosphorus and potassium fertilization. After 90 days, however, a lower phosphorus content of the juice of both leaves and

stems accompanied potassium fertilization. They also found that the potassium content of the juice paralleled the potassium applied as fertilizer, whereas the phosphorus fertilization decreased the potassium content in young, rapidly growing plants.

Mather (398) claimed in 1929 that fertilization increased the inorganic, but not the organic, phosphorus content of hays grown in Minnesota and Canada.

Murphy (422), working with the Kirkland sandy loam in Oklahoma, in 1930, reported an increase of about 20 percent in the phosphorus content of wheat grain when he applied superphosphates. Neither the application of potash nor of nitrogen alone resulted in any change in the phosphorus content of the grain.

Hester and Shelton (267, 268, 269), investigating the relationship between the replaceable bases in Virginia soils and the composition of vegetable crops, concluded that a high absorption of calcium at high pH value to the exclusion of the other ions (potassium in particular) was probably responsible for depressed yields. In general, he observed that

The presence of a large amount of a particular replaceable base in the soil colloidal complex influenced the elemental composition of plant material even though the yields were affected but little. A high replaceable calcium content suppressed the absorption of potassium, nitrogen and magnesium; a high replaceable magnesium content suppressed the absorption of potassium, calcium and nitrogen; and a high potassium content suppressed the absorption of calcium, magnesium and nitrogen.

The investigations of Tyson (569) in 1930 of the relationship between soil conditions in Michigan and the chemical composition of sugar beets indicated that the addition of potassium to unlimed plots resulted in increased potassium content of the leaves and roots and caused a decrease in the percentages of calcium, magnesium, sodium, and phosphorus. Applications of calcium resulted in a decrease in the potassium content and increased the phosphorus, sodium, and calcium content. Applications of phosphorus caused an increase in phosphorus and a decrease in potassium in the plant.

Wimer (596), studying the composition of mature corn stover in Illinois, reported in 1937 that

The general tendency of limestone was to raise the nitrogen, phosphorus, potassium and calcium contents of the stover and to increase the stover and grain yields. The chief exceptions were in the stover yields on soils naturally well supplied with carbonates, where the addition of limestone might be expected to have little or no effect.

Rock phosphate, when used in addition to residues (green manures) and limestone, resulted in increased nitrogen and phosphorus contents of the stover, but little or no change was observed in the contents of potassium and calcium. The use of potassium resulted in an increase in the potash content, a decrease in nitrogen and phosphorus, and no change in the calcium content.

Opitz, Rothsack, and Morgenroth (442), working with barley grown in a German soil, stated in 1937 that the omission of phosphorus from a complete fertilizer caused a reduction in the phosphorus content of the plant, a slight increase in the potash, and a significant increase in nitrogen. The omission of potassium resulted in a decrease in potassium and a slight increase in nitrogen, but no material change in phosphorus. Byczkowski (99), working with Polish soils, found that

the amount of potassium absorbed by a plant was dependent upon the anion of the salt used, the chloride being more readily assimilated than the sulfate.

Stalé (537) in 1937 reported that phosphates alone increased yields as much as did phosphorus and potassium together on certain glacial soils in Switzerland that were rich in available potash. An increase in the phosphorus content and a decrease in calcium content of hay was observed with phosphate fertilization. No consistent effect was noted on the magnesium and potassium contents. Potash alone resulted in no increased yields, a marked increase in the potash content, and a decrease in the calcium and magnesium content of the plant. No effect was observed on the phosphorus content of the plant.

Increases in the phosphorus content of hays, as a result of fertilization, have been reported by several investigators in Minnesota (6), Montana (98, 440), Florida (95), and Germany (321). Bishop (71) obtained increases in the phosphorus content of a number of vegetables (both leaves and roots) when phosphates were applied to Alabama soils, but the increases in phosphorus in plants grown in some soils as a result of fertilization were not large enough, in many cases, to produce plants that contained as much phosphorus as did those grown in other soils. Thus, a marked increase in the phosphorus content of turnip tops was obtained by the use of 1,000 pounds of superphosphate per acre of Oktibbeha soil, but still the phosphorus content did not equal that of turnip leaves grown on the Eutaw soil without phosphorus fertilization.

The effect of the liming of soils has been studied by a number of investigators. Gile and Ageton (209) in 1914 made extensive investigations of the effect of liming on the composition of crops grown in Puerto Rican soils. They reported that

The limestone apparently had no effect on the amount of nitrogen, potassium and phosphorus contained in the various plants, but did increase slightly the total carbon-free ash in all plants except rice. The calcium, magnesium and iron contents were all modified, the iron content sometimes being greatly reduced.

Ames and Schollenberger (8) showed in 1924 that the effect of liming on plant composition varied markedly with the soil. In general, however, they noted a tendency for liming to result in a decreased phosphorus content in the plant. Hjorth-Hansen (273) in 1931 reported that the contents of phosphorus, potassium, and calcium in the plant were influenced to a great extent by a change of soil reaction resulting from liming. The magnesium content remained quite constant.

Emmert (167), investigating liming practices in Kentucky, reported that the average percentages of the phosphorus and manganese in the plants grown in the more acid soils were greater than the average of those grown in the more nearly neutral soils, while Lott (358) found that calcium carbonate greatly decreased the absorption of zinc by plants.

Carolus (108) in 1933 reported that even small amounts of limestone increased the absorption of magnesium under conditions of excessive rainfall in Virginia. The available magnesium content of soils under such conditions is readily depleted by leaching, unless the soil contains in its internal colloidal phase an abundance of organic matter. Rendering the soil more alkaline with limestone results in an increase in this necessary organic material.

Naftel (426), investigating Alabama soils in 1937, found that liming with increasing increments of CaCO_3 increased the percentage of calcium in the plants but decreased the magnesium, potassium, phosphorus, manganese, and iron. Where soils were limed with dolomite there were similar decreases in mineral content, with the exception of magnesium, which was increased.

THE MINOR FERTILIZER ELEMENTS

Carolus (109) found that the addition of magnesium to the soil had a more pronounced effect on the magnesium content of the plant than it did on the yield or the nitrogen or calcium contents. Scharrer and Schropp (508) observed, in 1938, that the efficiency of magnesium absorption is clearly correlated with the natural magnesium content of the soil, decreasing with an increase in soil magnesium. An increase in magnesium fertilization, they state, resulted many times in an increase in the magnesium content of the plant, but the highest efficiency in absorption was found when the least magnesium was applied. Garner, McMurtrey, Bowling, and Moss (201) showed that the use of 120 pounds of magnesium applied to the Collington sandy loam in Maryland increased the magnesium content of tobacco leaves and prevented symptoms of magnesium hunger.

The application of magnesium resulted in increased content of this element in corn stover, oats (grain and straw), barley (grain and straw), millet, and buckwheat, but no increase was found in rutabagas (roots or tops) or Sudan grass, according to the experiments of Beaumont and Snell in Massachusetts (50).

Wiaddrowska (591), in her experiments with magnesium, using Polish soils, reported no increase in the magnesium content of lettuce, cabbage, carrots, or beans, although the calcium content of all vegetables except cabbage was depressed as a result of magnesium fertilization.

Wheeler, Hartwell, Kellogg, and Steel (590) reported in 1906 that sodium salts applied to Rhode Island soils increased the percentage of phosphorus in the plant. The carbonate was more efficient in this respect than was the chloride.

Deficiencies of sulfur have been studied extensively in the western part of the United States, and some of these investigations have included determinations of the effect of sulfur on the chemical composition of the plant.

In 1923, Neidig, McDole, and Magnuson (430) determined from pot tests that the sulfur content of alfalfa grown in several Idaho soils could be doubled by the application of sulfur at the rate of 100 pounds per acre. Even greater increases were obtained when 500 pounds of sulfur was applied. Smaller increases in sulfur were obtained when gypsum or lime and sulfur were added to the soil. All forms of sulfur produced an increase in the total nitrogen removed by the alfalfa. They stated, furthermore, that "the arid soils gave a greater [yield] response to applications of phosphorus and less to sulfur than did the humid soils." All applications of phosphorus increased the percentage of sulfur in the alfalfa and produced in all but two of these soils slight increases in the nitrogen content of the alfalfa.

Reimer and Tartar (480), in 1919, obtained an increase in both the sulfate-sulfur and organic-sulfur content of alfalfa grown in three Oregon soils when they fertilized with 300 pounds of sulfur per acre. Very large increases in the sulfur content of alfalfa were obtained by Neller, using two Washington soils, the Ritzville and the Palouse silt loams. The calcium content of the alfalfa was lowered and the phosphorus content was increased, but no significant change was observed in the potassium or nitrogen contents. Ligon (353), on the other hand, reported in 1935 that 1,000 pounds of sulfur applied per acre of muck soil in Michigan resulted in a decrease in the phosphorus as well as the calcium in the plant, with little or no change in potassium or nitrogen. Increases in the sulfur content of alfalfa following fertilization with sulfur, gypsum, or ammonium sulfate were found by Evans and Greaves (169).

No increase in the sulfur content of the plant was found by Barbier (46), who investigated the influence of sulfur and chlorine on the mineral nutrition of the plant. In the case of beets, an increment in the chloride content brought about by fertilization with chlorides was accompanied by an increment in the basic ions.

The addition of aluminum to culture solutions containing phosphates did not result in an appreciable increase in the aluminum content of lettuce, but the aluminum content of the roots was more than doubled, according to some investigations of Pierre and Stuart (465), reported in 1933. They state:

The results obtained definitely show that the problems of aluminum toxicity and phosphate availability are very closely interrelated. The so-called toxic action of soluble aluminum in very acid soils can be ascribed to two general causes. First, aluminum in high concentration directly injures the root hairs and thus causes stunted and poor development of the roots. Secondly, the aluminum absorbed by the plant precipitates part of the phosphate present, which interferes with phosphate translocation and assimilation. The addition of large amounts of phosphorus to the soil precipitates some of the aluminum out of the soil solution and enables the root to obtain a sufficient amount of phosphorus for nutritional purposes.

Hester (266) found that aluminum appeared in the soil solution at pH 3.9 to 4.4 in Portsmouth loamy fine sand, 4.2 in Bladen sandy loam, and 5.2 and below in Norfolk fine sand. He claimed that this was directly responsible for the low yields at these pH values, a problem that was overcome by applying lime to the soils. Other studies of aluminum in plants include those of Myers and Voegtlin (424) in 1914, Burgess and Pember (97) in 1923, Winter and Bird (597) in 1929, and Meunier (407) in 1935.

Densch and Hunnius (149) reported, in 1924, that fertilization with copper sulfate, although increasing yields of barley grain, caused no change in the mineral content of the plant except to depress the amount of iron assimilated.

Elvehjem and Hart (166) reported in 1929 increases of from 10 to 148 percent in the copper content of different vegetables when 50 pounds of copper sulfate per acre was used. Miller and Mitchell (411) in 1931 were able to increase the copper content of lettuce fourfold by adding 20 pounds of copper sulfate per acre in four equal doses. Much smaller increases were obtained in spinach. Copper had a slight depressive action on manganese and a stimulative action on the assimilation of iron, under the conditions of their experiments. Arnd and Hoffman (15) and Coleman and Ruprecht (119) reported that

fertilization with copper resulted in increased copper assimilation by the plant, but Rademacher (473) observed no such increase, even when increased yields resulted from the use of copper.

Iodine added as a fertilizer seems to be easily assimilated by the plant, and some investigators have reported manyfold increases in the iodine content of plants in this way. The work of Stoklasa (542) in 1926, Scharrer and Strobel (510) in 1927, Scharrer and Schwaibold (509) in 1927, Pfeiffer and Courth (463) in 1929, and Balks (45) in 1935 in Germany, Orr and coworkers in England (447) in 1928, and Beaumont and Karns (49) in the United States in 1932, may be cited as examples. McHargue, Young, and Calfee (383) in 1935 added potassium iodide from time to time to soils in which corn plants were growing, and they obtained, according to their report, an iodine content in the leaves of these plants of as much as 30,000 p. p. b. at one time during the season.

When no iodine was included in the fertilizer mixture, neither McHargue (383) nor Mack and Brasher (385) reported any relationship between the fertilizer treatment and the iodine in crops. The latter investigators, however, obtained large increases in iodine in beans and turnips by applying potassium iodide about 30 days after planting. Dietz (153) in 1938 was able to increase the iodine content of head lettuce (grown in pots) from 4,620 p. p. b. to 91,560 p. p. b. by applying 20 pounds of potassium iodide per acre 4 days before harvest.

Claims that Chile saltpeter contains sufficient iodine for plant growth have been disputed and affirmed by several authors (150, 206, 604). In general, it would seem that there are so many factors involved in determining the specific value of any trace element in such a material that no sound conclusions could be reached as to its effect on plant growth or composition.

Cook and Wilson (121) reported in 1917 the presence of boron in wheat grown in soils fertilized with borax and colemanite. No boron was found in plants grown without the use of this element as a fertilizer. Increased concentrations of boron in both leaves and stems of alfalfa were obtained by McLarty, Wilcox, and Woodbridge (387) after treating the soil with boric acid. The boron in the leaves of untreated plants was 11.4 p. p. m., while that of treated plants ranged from 214 to 734 p. p. m., depending upon the amount of boron added. Askew, Thomson, and Chittenden (29) reported in 1938 that the boron content of apples in areas top-dressed with boron was appreciably greater than that of the control fruit. Applications of 100 pounds of borax per acre had no greater influence on the boron content of the apple, however, than did 50 pounds of borax per acre. Hudig and Lehr (280), studying soils in the Netherlands, claim to have shown that the boron in Chile saltpeter was responsible for increased growth and boron content of plants.

Studies of the effect of arsenic on plant composition have originated with the problem of spraying. Pirotta (468) investigated this problem in Italy in 1906 and reported that the application of a dilute arsenical spray to plants resulted in an increase in the arsenic content of the plant. Increase of arsenic in the plant resulting from spray on the soil has not been reported, and Jones and Hatch (307) stated in 1937: "Analytical data on commercial orchard soils and fruits indicate

plainly that orchard welfare is not directly affected by soil spray residue accumulations, however long the period of accumulation." Their conclusions, of course, relate only to the fruit of trees and not to any crop such as a legume or cereal that might be planted in the soil. Hurd-Karrer (283) has observed that arsenic injury is inhibited by phosphorus applications to certain soils that do not fix the phosphorus in an unavailable state.

Stewart and Smith (540) in 1922, using a gravelly bench loam for pot tests, reported enormous increases in the arsenic content of pea, radish, wheat, potato, and bean plants. Disodium arsenate was added to the pots after the plant was fairly well developed.

McHargue and Shedd (382) studied the effect of manganese, copper, zinc, boron, and arsenic on the growth of oats. Their data, reported in 1930, show the effect of manganese only, for in only one experiment did they use a single element. In the remainder of the tests, manganese was used with copper or with copper and zinc, etc. The manganese content of oat straw was increased from 0.004 to 0.0644 percent, but no other significant change in the composition of the plant was found with manganese fertilization. Coleman and Ruprecht (119) in 1932 reported that increases in the manganese content of vegetables grown in Florida soils was obtained when manganese was applied as a fertilizer element.

Olsen (441) found that the absorption of manganese was related to the degree of acidity of the soil, and that increased quantities of manganese were assimilated by plants when the manganese content of the nutrient solution was increased. This observation is in harmony with those of Svanberg (544, 545, 546) and of Leeper (340).

Haselhoff (249) found traces of strontium in barley and bean plants grown in pots with soils to which had been added strontium carbonate. Calcium was found by Hurd-Karrer (284) to inhibit injury to plants by strontium.

Geilmann and Brunger (205) found germanium in plants grown in soils in pots to which germanium had been added.

Nemec and Kas (434) in 1923 obtained very slight increases in the titanium content of plants as a result of using Na_2TiO_3 in pot tests.

The influence of lead compounds on the growth of barley was investigated by Keaton (311) by adding lead nitrate or carbonate to soils in pots. His experiments show that while nearly all of the lead was fixed in an insoluble state in the soil within a few days, some lead was absorbed by the plant (2.53 to 5.40 p. p. m.) and a very large amount was absorbed by the roots (94.2 to 1475.0 p. p. m.).

Lepape and Trannoy (345) claimed to have found an increase in the radium content of plants grown in soils to which radium ores had been added.

Hart, Phillips and Bohstedt (244) studied the relation of soil fertilization with superphosphate and rock phosphate to the fluorine content of plants. They concluded that

Plant materials from plots receiving fluorine-carrying phosphates, such as rock phosphate and acid phosphates, for periods of sixteen to thirty-six years, did not show consistent or greatly increased fluorine content over plant materials grown in plots receiving a low fluorine-carrying phosphate such as bone meal.

A rather interesting survey was conducted in 1932 by Hartwell (247). He submitted three questions to several thousand agricul-

tural workers concerning the effect of fertilizers on the quality of the crops. A brief tabulation of some of the replies follows.

1. Effect of fertilizers on carbohydrates:
 - (a) Phosphorus increases the starch content of barley and alfalfa.
 - (b) Potassium produces greater hardness of rice kernels.
 - (c) Calcium and magnesium, by making potassium more available, increase carbohydrate content.
 - (d) Manganese increases the sugar content of beets.
2. Effect of fertilizers on proteins:
 - (a) Nitrogen generally increases the protein content of seeds and tubers.
 - (b) Phosphorus increases phosphorus and nitrogen of cereals.
 - (c) Phosphorus usually decreases gluten of wheat.
 - (d) Potassium tends to reduce protein content.
 - (e) Sulfur increases protein content of barley and wheat.
3. Effect of fertilizers on fats:
 - (a) Phosphorus has no influence on fat of flax.
 - (b) Potassium favors production of fat in flax, mustard, and pecans.
4. Effect of fertilizers on fiber:
 - (a) Too much nitrogen affects fiber unfavorably.
 - (b) Potassium increases length and strength of cotton fiber.
5. Effect of fertilizers on enzymes and vitamins:
 - (a) Phosphorus hinders enzymatic activity in plant tissues.
 - (b) Lack of manganese in spinach is correlated with decreased vitamin A.

Ashton (17) observed that the strength of oat straw was associated with a relatively high content of certain ash constituents, particularly silica, and a low fiber content. Studies on the relationship between the calcium content of the plant and the organic acids in the plant have been made by Iljin (290) in 1938; and Pfutzer and Pfaff (464) have recently reported on the effect of minerals on the vitamin content of plants.

Garner, McMurtrey, Bowling, and Moss (202) reported in 1930 that

From the standpoint of quality the chloride content of the cured [tobacco] leaf may affect its properties either favorably or adversely depending on the quantity of chlorine present, the use to be made of the tobacco, and the stage in handling and manufacturing operations which it has reached.

SUMMARY OF EFFECT OF FERTILIZERS ON PLANT COMPOSITION

Out of the confusion of contradictory results of fertilizer experiments one can draw a very few definite conclusions. First among these is the important fact that no fundamental study has ever been made of the effect of soil amendments on plant composition taking into consideration that most important of factors, the characteristics of the soil used in the experiment. It is true that physical characteristics of the soil such as structure have been observed, and that the availabilities of a few of the principal plant nutrients have been arbitrarily measured; but fundamental studies of what changes take place in the soil when the fertilizers are applied and the effect these changes will have on the plant are lacking.

One or two recent empirical investigations have quite definitely shown that Liebig's "law of the minimum" never represented the mechanism of absorption of nutrients by plants, and that the actual facts seem to indicate that when one of the principal nutrients is deficient in the soil solution the others are taken up by the plant in amounts greater than normal, while even less of the deficient nutrient is absorbed than would be expected normally. The necessity for a balanced source of nutrients for the plant is thus indicated, and the

variance in results by most investigators would tend to prove that this is a problem that can be solved only by study of the characteristics of each soil type.

In general it can be stated that in many instances known deficiencies of some particular element can be remedied by adding that element to the soil. Deficiencies of phosphorus and iodine are good examples of such a situation that seems to be easily remedied.

The use of soil amendments to prevent injury to plants by certain elements seems to be a promising field of investigation, although the experience of Franke and Painter (196) with selenium emphasizes the necessity for knowing in what form the toxic element may exist in the soil, or, as in the case of the inhibition of aluminum injury by phosphorus, whether or not the phosphorus may become "fixed" by the soil before it reacts with the aluminum.

Liming of soils has varied effects on the composition of the plant, but in most cases it exerts a depressing effect on iron and manganese. Depending upon the soil, for example, the phosphorus in the plant may be depressed, unchanged, or increased by liming. The same observations apply to the effect of liming on magnesium and other elements. The Virginia work (108) cited in this report shows clearly how indirect effects of soil treatment may result in conclusions being made in one place that are directly opposed to those in some other locality.

The effect of the applications of minor elements to the soil has been studied mainly from the viewpoint of increasing that element in the plant, and not from the viewpoint of modifying the general plant composition. The latter type of investigations would seem to be of paramount importance.

The influence of the fluorine in phosphatic fertilizers on the fluorine content of the plant has been determined in a very limited area. What the effect is on plants grown in soils of different types than those studied is entirely unknown.

Some observations of the effect of fertilizers on the quality and the technological uses of crops have been made.

MISCELLANEOUS FACTORS INFLUENCING THE MINERAL COMPOSITION OF PLANTS

AGE OF PLANT

It has already been noted that changes in the mineral composition of the plant as it matured were observed in the early part of the nineteenth century, and many investigations since that time have emphasized the importance of this factor in relation to animal nutrition problems. A few of these investigations are cited here in order to demonstrate the necessity of making observations on the age of the plant in interpreting soil and fertilizer studies.

Hills (270) in 1893 noted that both the phosphorus and potassium contents in corn decreased rapidly with the age of the plant, until, in some cases, less than half of the amounts present in the plant at tasseling stage were found in the mature plant.

Recent investigations include those of the sunflower by Neidig and Snyder (431) in 1925 and of the alfalfa plant by Sotola (532) in 1927. Sotola determined the chemical composition, digestibility, and protein utilization of the first, second, and third cuttings of alfalfa

grown in Washington. Each of these three cuttings was studied at the one-fourth, one-half, and three-fourths bloom. His data show that the protein content of the hay decreased and the fiber content increased as the plant matured. Plants cut at one-half bloom stage contained more calcium than similar samples harvested at the one-fourth and three-fourths stages.

Radu (474) in 1936 reported a maximum concentration of calcium in the alfalfa plant during its growth, but Bettini (69), studying the composition of alfalfa in Italy, found a progressive increase in calcium throughout the life of the plant until the advanced-bloom stage, after which it remained constant until the seeds were mature.

Determination of the variation in composition of spinach was made by Fondard and Margailan (187) in France in 1928. They reported that the phosphorus content increased for about the first 30 days of growth, after which it rapidly decreased until it was less in 48 days than it had been at the end of 10 days. The potash increased during the first 17 days but remained quite constant thereafter. According to Tuorila (568), the potash content of potatoes grown in Finland decreased with maturity.

A comprehensive study of the composition of the different parts of the soybean plant at different stages of growth was reported by Borst and Thatcher (84). Their data show that a reduction of calcium, magnesium, potassium, and phosphorus occurred in the stems as the plant approached maturity. A reduction in the potassium and phosphorus contents was also found in the leaves, but the other elements remained fairly constant.

A number of investigations in changes in the composition of grasses have been conducted by workers in Great Britain (171, 172, 601). In general, except for timothy, they reported that the phosphorus content is reduced with the age of the plant.

Van Itallie (292), working with three Dutch soils, reported in 1934 that the differences in composition of one variety or species of grass at different stages of growth are greater than differences between different species at the same stage of growth.

Murphy (423) in 1936 found that both the calcium and phosphorus content of the cotton plant decreased with age, and that the rate of decrease in calcium was accelerated after the bolls started to form.

Considerable attention was given in the compilation of the mineral composition of crops to classifying the crop as to its age or stage of maturity (see appendix). Thus, with alfalfa the average contents of calcium and of phosphorus at the bud stage were 2.45 and 0.36 percent, respectively, while the corresponding values at one-half bloom were 2.01 and 0.27 percent. The alfalfa data are not sufficiently numerous to illustrate the change in composition throughout the life of the plant, but other examples include the compilation of data on Kentucky bluegrass and on rape plant. In the latter case single analyses of the plant grown in Iowa show a maximum in the calcium content during the life of the plant and a steadily diminishing phosphorus content.

Variations in the nutritive value of plants at different stages of growth and the stage of maturity at which most plants have the highest nutritive value are well-known. In general, the nutritive elements attain maximum values during the life of the plant although some differences in observations of how the mineral composition of

plants varies with the age of the plant are not explainable at the present time, but a very likely cause is the availability of these elements or the rate at which they become available in the soil. Although it has long been known that forage and other crops must be harvested at certain stages of development so that they will be palatable and so as to obtain maximum yields of dry matter, comparatively little attention has been given to obtaining maximum yields of nutrients. As the plant matures, particularly beyond the bloom stage, the percentage of digestible nutrients and of available mineral matter may drop so low as to nullify completely the effects of increased yield of total dry matter.

CLIMATE

Climate was shown by LeClerc and coworkers (336, 338, 339) to be of greater influence on the protein content and slightly more influence on the phosphorus and potassium contents of wheat grain than was the soil. Unfortunately, their work did not include the composition of the straw nor of any part of the plant that would respond more accurately to soil deficiencies than does the seed. The work is of interest, however, because of the procedure, which consisted of transporting fairly large amounts of soils from two of three localities (California, Kansas, and Maryland) to the third locality for plot tests.

In similar experiments with sugarcane, Borden (83) failed to demonstrate the predominance of climate over that of soil as an influence on the mineral composition of the plant. The two localities for this experiment, although only 3 miles apart in Hawaii, differed markedly in climatic conditions, for, although the temperature range is about the same for both places, Manoa receives nearly four times as much rainfall and 40 percent less sunshine than does Makiki. Tubs of soil from both localities were planted with three different varieties of sugarcane, which were allowed to grow at each locality. The results, classified as to soils and as to localities, are given in table 18. The climate at Makiki was apparently quite favorable for the production of sugar, but while the cane grown at Manoa has a slightly higher potassium and phosphorus content than has that grown at Makiki, the dominating influence of soils is evident, for the average content of these minerals in the cane grown in the Makiki soil at both localities is much higher than that of cane grown in the Manoa soil.

TABLE 18.—*Influence of soil and of climate on the sugar, phosphorus, and potassium content of sugarcane*

[Borden (83)]

Items	Sugar produced	P in 100 cc. of juice	K in 100 cc. of juice
Averages by soils:	<i>Pounds</i>	<i>Percent</i>	<i>Percent</i>
Makiki soil.....	5.41	0.044	0.21
Manoa soil.....	6.26	.014	.10
Averages by localities:			
Makiki soil.....	8.93	.027	.134
Manoa soil.....	2.73	.031	.172

Kunze (322) in 1914 determined the phosphorus, potassium, and calcium content of oats grown in normal and dry years in Germany. No significant differences were observed in the phosphorus content of the grain, but about half as much phosphorus was found in the

straw and chaff in dry years as in good years. Both the calcium and potassium contents were much higher in the dry years, that in the straw and chaff often being double the amount found in these parts in normal years.

The data of Daniel and Harper (135), reported in 1935 from their study of the relation between effective rainfall in Oklahoma and total calcium and phosphorus in alfalfa and prairie hay, support these findings. They state that

Although the mathematical correlation between the actual amount of rain and the phosphorus in the alfalfa hay is not high, the figures show that the phosphorus content of alfalfa fluctuates with the rainfall, and that the calcium content of the hay is inversely proportional to the quantity of effective rainfall which occurs. The moisture requirement of a crop is affected by the temperature and humidity of the atmosphere; consequently, the total rainfall could influence plant growth quite differently, depending upon whether high rainfall occurs during the spring or summer months. A greater variation in the calcium and phosphorus content of alfalfa occurred between the rainfall and the different cuttings of hay harvested in 1932 than in the hay harvested in 1933. During the season of 1932, the period of high rainfall was in May and June, while in 1933 the period of high rainfall was in March, April, and May.

Salgues (501), working with two soils in France, reported in 1938 that the phosphorus content of plants grown in calcareous soils was increased with high rainfall, but that this was not true of plants growing in siliceous soils. With lower precipitations, the phosphorus-calcium ratio was lowered for all species grown in calcareous soils, but the ratio was actually higher for the same species grown in siliceous soils.

The effect of rainfall on leaching of minerals from plants was investigated by LeClerc and Breazeale (337) in 1909. They found that mineral nutrients, exuded upon the surface of the leaves of plants, are washed off by rain and dew. They concluded, therefore, that the analysis of plants gathered at maturity may give erroneous or misleading results, especially for comparative purposes. Guilbert, Mead, and Jackson (233), found that leached plants contained considerably less phosphorus than did normal plants, although the calcium contents of the leached and normal plants did not materially differ.

While plants are dependent upon the soil for their mineral nutrients, climatic conditions so affect respiration, assimilation, photosynthesis, metabolism, and other physiological processes that the composition of both the mineral and the organic matter of crops may be greatly modified even though the crops are grown upon identical soils. Indirect influences of climate are also reflected in the composition of the plant, for the significant differences in the characteristics of our great soil groups are largely differences due to climate. Thus, in the Southeast we have well-leached acid soils low in the basic constituents, while in the Northwest the soils are more nearly neutral and high in bases. Differences within any soil group have, furthermore, been effected by the particular vegetation growing in the soil, which in turn was influenced by climatic factors. Thus, a soil developed under grass will be quite different from one developed under forest.

Investigations that show clearly the effect of climate on the mineral metabolism and composition of the plant are not available, however because differences in soils, species of plants, time of rainfall in relation to the growing season, and other factors have not always been

considered. It seems quite safe to conclude, however, that the plant composition is modified by both the climate and the soil in which it is growing, and that both of these factors are closely inter-related, often one modifying the effect of the other. Thus the assimilation of phosphorus by the plant growing in calcareous soils may be less in dry than in wet years, whereas entirely contrary results may be obtained in plants growing in siliceous soils. The period of plant growth at which high rainfall occurs is of paramount importance in the assimilation of nutrients by the plant as well as in the growth of the plant, and, for this reason, crops growing in the same soil type separated by a sufficient distance so as not to receive the same amount of rainfall at the same time in any one year may be quite different in their chemical composition.

IRRIGATION

The discussion under climate indicates that the composition of the plant, under some conditions at least, is closely related to the moisture or rainfall available during certain times of the growing period. Irrigation is a climatic factor, inasmuch as moisture is supplied artificially instead of naturally.

In 1900, von Daszewski and Tollens (137) studied the effect of irrigation in Germany on the composition of potatoes. They concluded that although yields were increased with irrigation, no change in the phosphorus and potassium content of the plant occurred. Stahl-Schroder (536) reported that increasing the water in a soil from 35 up to 95 percent of its water capacity resulted in a large increase in ash, silica, and phosphorus in the grain of oats. The nitrogen content decreased to a minimum at 90-percent water capacity, but increased markedly at 95 percent.

Irrigation in varying amounts had no effect on the nitrogen content of the corn grain, according to Harris and Pittman (241), in Utah. The same observation held for phosphorus, calcium, and magnesium, although a large variation in the mineral nutrients was observed in the crops grown in different years, but with the same amount of irrigation. The opposite effect was reported by Greaves and Carter (221) in 1923, in their experiments with wheat, oats, and barley. These authors found that phosphorus, potassium, calcium, and magnesium in the grain were all increased as a result of irrigation. Greaves and Nelson (224) showed that while the iron, chlorine, and sulfur contents of wheat were increased by irrigation, these elements in oats and barley either underwent no change or actually decreased. Further work of Greaves and Nelson (223), in 1925, using corn, confirms the work of Harris and Pittman, who used the same crop.

The addition of irrigation water to potatoes grown in Georgia had no influence on the iodine content of that crop, according to the experiments of Holley, Pickett, and Brown (274), reported in 1935.

The effect of irrigation, like climate, varies with other factors such as the species of plant, the soil, or the amount of water added. The data are too few and too narrow in scope to justify any general conclusions regarding this factor.

VARIETY

Whether different varieties of the same species differ in mineral content when grown in the same soil is still an unsolved problem. It would seem reasonable that there should be differences, for the differences in yields among varieties indicate variations in the feeding power of the plant that might be reflected in their mineral composition. Considerable work has been done by Fort (10x) on the mineral variations in different varieties of the juice of sugarcane, and his data indicate that this factor may be of considerable importance. His studies did not include careful soil surveys, but the same varieties of cane occupied the same relative positions in regard to mineral content in each area in which they were grown. Ayres (38) in 1936 found little difference in the phosphorus content but significant variations in the potassium content of two varieties of sugarcane grown on the same soil. Similar results were obtained by Deomano (151) with four varieties of sugarcane, and by Maxwell (401), who analyzed 13 varieties of cane grown under uniform conditions.

Remy (483), in 1922, failed to find any variation in the phosphorus content of different varieties of corn grain, and Maume and Bouat (399) reported, in 1937, similar findings on the sulfur content of different varieties of wheat. Greaves and Anderson (220), on the other hand, reported in 1936 that they obtained about one-third as much copper in Kofod \times Turkey variety of wheat as in Montana 36, and many varieties of wheat varied significantly in copper when grown in the same soil. Likewise, Bertrand and Levy (59) stated that while aluminum is present in all phanerogams, the amount present varies significantly with the species, exceptional species containing as much as 1 gm. per kilogram of dry matter.

Peterson and coworkers (456, 459, 460) found that different varieties of cabbage were remarkably uniform in their contents of calcium, phosphorus, iron, and nitrogen, and similar results were reported in 1938 by Vogel and von Hosslin (577).

Delbet (145) examined potatoes grown in the same soil in Egypt and found only normal variations in calcium, magnesium, potassium, sodium, and phosphorus content of the several different varieties. Poehlman (469), in some comprehensive experiments on two varieties of soybeans, concluded in 1935 that

Variance due to varieties is not significantly different from the variance due to experimental error. This signifies that we cannot measure any significant differences between Morse and Virginia varieties of soybeans by chemical analyses of the expressed plant juice for nitrogen, phosphorus or potassium.

Weathers (580) apparently obtained no differences due to variety in the calcium, magnesium, potassium, iron, manganese, or phosphorus content of four varieties of lespedeza, all varieties having been grown in three different localities.

Some differences in potassium and phosphorus contents of corn stover due to varietal differences were reported by Wimer (596). These variations were not always consistent under different conditions, however, but they seemed to be affected greatly in some cases by grain and stover yields.

Ashton (18) reported in 1938 that different varieties of oats showed only very slight differences in chemical composition. Van Itallie (293) working with soils in the Netherlands, in 1937, found that

there was a fairly consistent relationship between the variety of sugar beets and their potassium and sodium contents. A variety consistently high in potassium was usually found to be low in sodium.

Smirnova and Lavrova (529), working with the soybean plant in Russia, concluded that variations of the ash content of different varieties grown on one soil are not significant. Conditions of growth were associated with greater differences in mineral content than was varietal difference.

Definite conclusions as to the effect of the variety of the plant on the chemical composition cannot be made at this time. It is entirely possible that in some species one variety is a better feeder on certain nutrients than are other varieties of the same species. Such a supposition would seem to be at least a reasonable one. There are few data to support this assumption, however, because experiments of this nature have not included careful investigations that would assure uniform soil conditions.

MINERAL CONTENT OF DIFFERENT PARTS OF THE PLANT

Many examples of variations in the mineral content of the different parts of the plant have been cited in the compilation of analyses given in the appendix. Thus, the average value for calcium in the stem of the alfalfa plant is nearly 40 percent less than that in the leaf, and the data for the corn plant show that the leaf contains about 14 times as much calcium and less than one-third the phosphorus as does the grain. Similar differences are always found among the seed, stalk, or leaves of plants. As an example of the care that must be taken in sampling a crop, the data in table 19 are presented, showing the enormous differences in the calcium content of the leaves of the same head of cabbage.

TABLE 19.—*Calcium content of leaves of cabbage*

[Cowell (125)]

Sample No.	Description of leaf	Ca per 100 gm. moist leaf
		<i>Mg.</i>
1	Outer dark-green leaf	476
	Inner pale-green leaf	35
2	Outer dark-green leaf	910
	Inner yellowish-heart leaf	34
3	Outer dark-green leaf	998
	Inner pale-green leaf	53
4	Outermost leaf, dark-green	708
	Third leaf from outside, pale-green	96
	Yellowish white-heart leaf	26
5	Outermost leaf, dark-green	1,058
	Third leaf from outside, dark-green	216
	Inner leaf, pale-greenish yellow	71
	Heart leaf, yellowish-white	32

Sarata (502), on the contrary, reported in 1937 only slight differences in the copper content of different leaves of the Pe-tsai cabbage. The structure of this plant is, however, quite different from the compact head of common cabbage. Darkis and coworkers (136) have recently determined the difference in composition of the leaves from different parts of the tobacco plant, and they report that both calcium and potassium vary considerably according to the position of the leaf on the stalk. This factor is also of importance in the studies of Thomas and of Lagatu and Maume already cited.

LITERATURE CITED

- (1) ABBOTT, O. D., BRYAN, O. C., and NEAL, W. M.
1937. STUDIES OF NUTRITIONAL ANEMIA OF RURAL SCHOOL CHILDREN IN CERTAIN AREAS OF FLORIDA. *Assoc. South. Agr. Workers Proc.* 37-38: 257-258.
- (2) ———, NEAL, W. M., and BRYAN, O. C.
1934. HUMAN DIETARY DEFICIENCIES IN ALACHUA COUNTY, FLORIDA, WITH SPECIAL REFERENCE TO NUTRITIONAL ANEMIA IN RELATION TO THE COMPOSITION OF HOME GROWN FOODS. *Fla. Agr. Expt. Sta. Ann. Rpt.* 58-59.
- (3) ADOLPH, W. H., and CH'EN, S. C.
1930. IODINE IN NUTRITION IN NORTH CHINA. *Chinese Jour. Physiol.* 4: 437-447.
- (4) ——— and CH'EN, S. C.
1932. IODINE IN NUTRITION IN COASTAL MID CHINA. *Chinese Jour. Physiol.* 6: 345-352.
- (5) ——— and PROCHASKA, F. J.
1929. AN IODINE SURVEY OF NEBRASKA. *Amer. Med. Assoc. Jour.* 92: 2158-2160.
- (6) ALWAY, FREDERICK J., SHAW, WILLIAM N., and METHLEY, WILLIAM J.
1926. PHOSPHORIC ACID CONTENT OF CROPS GROWN UPON PEAT SOILS AS AN INDEX OF THE FERTILIZATION RECEIVED OR REQUIRED. *Jour. Agr. Res.* 33: 701-740, illus.
- (7) AMES, J. W.
1910. THE COMPOSITION OF WHEAT. *Ohio Agr. Expt. Sta. Bul.* 221, 37 pp., illus.
- (8) ——— and SCHOLLENBERGER, C. J.
1924. THE AVAILABILITY OF PHOSPHORUS IN CALCAREOUS AND NON-CALCAREOUS SOILS. *Ohio Agr. Expt. Sta. Bul.* 380; pp. 215-242.
- (9) ANDERSON, J. T.
1894. FERTILIZERS AS REQUIRED BY COTTON AS DETERMINED BY THE ANALYSIS OF THE PLANT. *Ala. Agr. Expt. Sta. Bul.* 57, 16 pp.
- (10) ANDERSON, P. J., and SWANBACK, T. R.
1929. REPORT OF THE TOBACCO SUBSTATION AT WINDSOR FOR 1928. *Conn. Agr. Expt. Sta. Bul.* 299, pp. 145-198.
- (11) ARCHIBALD, J. G., and BENNETT, E.
1935. YIELD AND CHEMICAL COMPOSITION OF CERTAIN SPECIES OF GRASS. *Jour. Agr. Res.* 50: 711-715.
- (12) ——— and KUCINSKI, K. J.
1938. NUTRITIONAL ANEMIA IN CATTLE IN SOUTHEASTERN MASSACHUSETTS. *Mass. Agr. Expt. Sta. Bul.* 347 (Ann. Rpt. 1937): 24.
- (13) ——— KUCINSKI, K. J., BROOKE, R. O., and FREEMAN, S. L.
1938. NUTRITIONAL ANEMIA IN CATTLE IN SOUTHEASTERN MASSACHUSETTS. *Jour. Dairy Sci.* 21: 59-68.
- (14) ARMSTRONG, S. F.
1907-8. THE BOTANICAL AND CHEMICAL COMPOSITION OF THE HERBAGE OF PASTURES AND MEADOWS. *Jour. Agr. Sci. [England]* 2: 283-304.
- (15) ARND, TH., and HOFFMAN, W.
1937. SPURENELEMENTE UND IHRE WIRKUNG AUF DAS PFLANZENWACHSTUM UNTER BESONDERER BERUECKSICHTIGUNG VON VERSUCHSERGEBNISSEN MIT KUPFER. *Landw. Vers. Sta.* 129: 71-99.
- (16) ARTIS, BONNIBEL, and MAXWELL, HAROLD L.
1916. BARIUM IN TOBACCO AND OTHER PLANTS. *Chem. News* 114: 62-63.
- (17) ASHTON, W. M.
1937. THE CHEMICAL COMPOSITION OF THE STRAW OF STRONG AND WEAK STRAWED VARIETIES OF OATS. *Welsh Jour. Agr.* 13: 144-151.
- (18) ———
1938. THE CHEMICAL COMPOSITION OF THE GRAIN AND STRAW OF VARIETIES OF OATS BRED AT THE WELSH PLANT BREEDING STATION: COMPARISON WITH SOME OLDER VARIETIES. *Empire Jour. Expt. Agr.* 6: 69-78.
- (19) ASKEW, H. O.
1933. INFLUENCE OF SEASON AND OF AMMONIUM SULFATE ON THE CHEMICAL COMPOSITION OF PERENNIAL RYE-GRASS AND OF WHITE CLOVER. *New Zeal. Jour. Sci. and Technol.* 15: 143-154.

- (20) ASKEW, H. O.
1935. THE BORON STATUS OF FRUIT AND LEAVES IN RELATION TO INTERNAL CORK OF APPLES IN THE NELSON DISTRICT. PRELIMINARY REPORT. *New Zeal. Jour. Sci. and Technol.* 17: 388-391.
- (21) ———
1936. INTERNAL CORK OF APPLES. *Orchardist of New Zeal.* 9: 185-188.
- (22) ——— and DIXON, J. K.
1936. THE IMPORTANCE OF COBALT IN THE TREATMENT OF CERTAIN STOCK AILMENTS IN THE SOUTH ISLAND, NEW ZEALAND. *New Zeal. Jour. Sci. and Technol.* 18: 73-92.
- (23) ——— and DIXON, J. K.
1937. INFLUENCE OF COBALT TOP-DRESSING ON THE COBALT STATUS OF PASTURE PLANTS. *New Zeal. Jour. Sci. and Technol.* 18: 688-693.
- (24) ——— and DIXON, J. K.
1937. THE VALUE OF COBALT SALTS FOR PASTURE TOP-DRESSING IN THE TREATMENT OF STOCK. *New Zeal. Jour. Sci. and Technol.* 19: 317-325.
- (25) ——— and MAUNSELL, P. W.
1937. THE COBALT CONTENT OF SOME NELSON PASTURES. *New Zeal. Jour. Sci. and Technol.* 19: 337-342.
- (26) ——— and RIGG, T.
1932. OCCURRENCE OF BUSH SICKNESS AT GLENHOPE, NELSON, NEW ZEALAND. *New Zeal. Dept. Sci. and Indus. Res. Bul.* 32: 1-20.
- (27) ——— and RIGG, T.
1934. SHEEP-SICKNESS IN THE PAKAWAU DISTRICT OF COLLINGWOOD COUNTY, NELSON. *New Zeal. Jour. Sci. and Technol.* 15: 305-308.
- (28) ——— and THOMSON, R. H. K.
1937. OCCURRENCE OF INTERNAL CORK OF APPLES IN CENTRAL OTAGO, NEW ZEALAND. *New Zeal. Jour. Sci. and Technol.* 18: 661-664.
- (29) ——— THOMSON, R. H. K., and CHITTENDEN, E.
1938. EFFECT OF BORAX TOP-DRESSING ON BORON STATUS OF SOIL AND FRUIT. *New Zeal. Jour. Sci. and Technol.* 20A: 74-78.
- (30) ASTON, B. C.
1912. THE CHEMISTRY OF BUSH SICKNESS. *New Zeal. Dept. Agr. Ann. Rpt.* 20: 75-78.
- (31) ———
1928. MINERAL CONTENT OF PASTURES. *New Zeal. Jour. Agr.* 36: 22-27, 75-82.
- (32) ———
1928. MINERAL CONTENT OF PASTURES. LIME-DEFICIENT AREAS AND A HITHERTO UNSUSPECTED MALNUTRITION DISEASE IN SHEEP. *New Zeal. Jour. Agr.* 37: 145-151.
- (33) ———
1930. MINERAL CONTENT OF PASTURES RESEARCH. *New Zeal. Jour. Agr.* 41: 90-99.
- (34) ——— GRIMMETT, R. E. R., BROGAN, F. J. A., and SYKES, P. H.
1931. MINERAL CONTENT OF PASTURES. THE WAIRARAPA DISTRICT. *New Zeal. Jour. Sci. and Technol.* 12: 304-320.
- (35) ATKINSON, H. J., and WOODWARD, J. C.
1938. PASTURE RESEARCH. CENTRAL EXPERIMENT FARM, OTTAWA. *Canad. Chem. and Proc. Indus.* 22: 67-71.
- (36) AUCHTER, E. C.
1939. THE INTERRELATION OF SOILS AND PLANT, ANIMAL AND HUMAN NUTRITION. *Science* 89: 421-427.
- (37) AUSTIN, RUSSELL H.
1928. THE EFFECT OF SOIL TYPE AND FERTILIZER TREATMENT ON THE COMPOSITION OF THE SOYBEAN PLANT. 30 pp. [Thesis Mich. State Col. of Agr. and Appl. Sci. For an abstract of this thesis under the same title refer to *Amer. Soc. Agron. Jour.* 22: 135-156 (1930).]
- (38) AYRES, ARTHUR.
1936. FACTORS INFLUENCING THE MINERAL COMPOSITION OF SUGAR CANE. *Assoc. Hawaii. Sugar Technol. Rpts.* 15: 29-41.

- (39) BACHOFEN, F.
1900. A COCOANUT ANALYSIS. *Queensland Agr. Jour.* 6: 297.
- (40) BADGER, C. J., and SNIDER, H. J.
1933. THE COMPOSITION OF THE SPRING GROWTH OF SWEET CLOVER AS INFLUENCED BY PREVIOUS FALL TREATMENT. *Amer. Soc. Agron. Jour.* 25: 105-108.
- (41) BAILEY, E. H. S., and SAYRE, L. E.
1911. ON THE PRESENCE OF BARIUM IN THE ASH AND EXTRACT OF CERTAIN KANSAS WEEDS. *Kans. Acad. Sci. Trans.* 23-24: 194-198.
- (42) BAILEY, E. M., and ANDERSON, P. J.
1928. CHEMICAL ANALYSES OF TOBACCO FROM THE NITROGEN PLOTS. *Conn. Agr. Expt. Sta. Tobacco Sta. Bul.* 10: 35T-50T.
- (43) ——— and ANDERSON, P. J.
1930. CHEMICAL COMPOSITION OF A POOR BURNING TOBACCO CROP COMPARED WITH A GOOD BURNING CROP. *Conn. Agr. Expt. Sta. Bul.* 311: 228-233.
- (44) BAKER, G. O., and VANDECAVEYE, S. C.
1935. EFFECT OF FERTILIZERS, SOIL TYPE, AND CERTAIN CLIMATIC FACTORS ON THE YIELD AND COMPOSITION OF OATS AND VETCH. *Jour. Agr. Res.* 50: 961-974, illus.
- (45) BALKS, RUDOLPH.
1935. UNTERSUCHUNGEN ZUR JODFRAGE IN WESTFALEN. *Landw. Jahrb.* 81: 939-1002. [The same data appear under the title, *Über den Jodgehalt von Lebensmitteln. Ztschr. f. Untersuch. der Lebensmitl.* 71: 76-93 (1936).]
- ✓(46) BARBIER, G.
1937. ACTION DES CHLORURES ET DES SULFATES SUR LA NUTRITION MINÉRALE DE LA PLANTE. *Acad. d'Agr. de France Compt. Rend.* 23: 700-706.
- (47) BARNES, EARL E.
1924. EFFECT ON PERMANENT PASTURES OF TREATMENTS WITH LIMESTONE AND ACID PHOSPHATE AS MEASURED BY THE QUANTITY AND QUALITY OF THE VEGETATION PRODUCED. *Amer. Soc. Agron. Jour.* 16: 241-251.
- (48) BEATH, O. A., EPPSON, H. F., and GILBERT, C. S.
1935. SELENIUM AND OTHER TOXIC MINERALS IN SOILS AND VEGETATION. *Wyo. Agr. Expt. Sta. Bul.* 206, 56 pp.
- (49) BEAUMONT, A. B., and KARNS, GEO. M.
1932. EFFECT OF AN IODIDE FERTILIZER ON IODINE CONTENT OF A FOOD PLANT. *Science* 76: 567.
- (50) ——— and SNELL, M. E.
1935. THE EFFECT OF MAGNESIUM DEFICIENCY ON CROP PLANTS. *Jour. Agr. Res.* 50: 553-562., illus.
- (51) BECKER, R. B., NEAL, W. M., and SHEALY, A. L.
1931. I. SALT SICK: ITS CAUSE AND PREVENTION. II. MINERAL SUPPLEMENTS FOR CATTLE. *Fla. Agr. Expt. Sta. Bul.* 231, 23 pp.
- (52) ——— NEAL, W. M., SHEALY, A. L., and YORK, GUS.
1933. STIFFS OR SWEENY (PHOSPHORUS DEFICIENCY) IN CATTLE. *Fla. Agr. Expt. Sta. Bul.* 264, 27 pp., illus.
- (53) BEESON, K. C., and LECLERC, J. A.
1937. THE SIGNIFICANCE OF THE SOIL TO HUMAN AND ANIMAL NUTRITION. *Soil Sci. Soc. Amer. Proc.* 2: 335-341.
- (54) BENNETTS, H. W.
1937. COPPER DEFICIENCY—THE CAUSE OF ENZOÏTIC ATAXIA (RICKETS) OF LAMBS. *Jour. Dept. Agr. W. Australia* 14: 247-250.
- (55) BERTRAND, GABRIEL.
1920. DE L'IMPORTANCE DES INFINIMENT PETITS CHIMIQUES DANS L'ALIMENTATION. *Soc. Sci. d'Hyg. Aliment. Bul.* 8: 49-66.
- (56) ——— and ANDREITSCHEVA.
1934. SUR LA TENEUR COMPARÉE EN ZINC DES FEUILLES VERTES ET DES FEUILLES ÉTIOLÉES. *Inst. Pasteur [Paris] Ann.* 52: 249-251.
- (57) ——— and BENZON, BOJE.
1929. SUR LA TENEUR EN ZINC DES PRINCIPAUX ALIMENTS D'ORIGINE VÉGÉTALE. *Inst. Pasteur [Paris] Ann.* 43: 386-393. [The same data appear under the same title in *Soc. Sci. d'Hyg. Aliment. Bul.* 16: 457-463 (1928).]

- (58) BERTRAND, GABRIEL, and GHITESCU, VIRGIL.
1934. SUR LA COMPOSITION ÉLÉMENTAIRE DE QUELQUES PLANTES CULTIVÉES. [Paris] Acad. des Sci. Compt. Rend. 199: 1269-1273. [The same data appear under the title, La Composition Élémentaire de Quelques Plantes de Grand Culture. Acad. d'Agr. de France Compt. Rend. 20: 1052-1056 (1934).]
- (59) ——— and LEVY, GEORGETTE.
1931. L'ALUMINUM DANS LES PLANTES. Acad. d'Agr. de France Compt. Rend. 17: 235-238.
- (60) ——— and MOKRAGNATZ, M.
1930. RÉPARTITION DU NICKEL ET DU COBALT DANS LES PLANTES. [Paris] Acad. des Sci. Compt. Rend. 190: 21-25.
- (61) ——— and MOKRAGNATZ, M.
1930. SUR LA RÉPARTITION DU NICKEL ET DU COBALT DANS LES PLANTES. Soc. Chim. de France Bul. (4) 47: 326-331.
- (62) ——— and PERIETZEANU, J.
1927. SUR LA PRÉSENCE DU SODIUM CHEZ LES PLANTES. Soc. Chim. de France Bul. (4) 41: 709-712.
- (63) ——— and ROSENBLATT, M.
1921. RÉCHERCHES SUR LA PRÉSENCE DU MANGANÈSE DANS LE RÈGNE VÉGÉTAL. Inst. Pasteur [Paris] Ann. 35: 815-819. [The same data appear under the same title in [Paris] Acad. des Sci. Compt. Rend. 173: 333-336 (1921).]
- (64) ——— and ROSENBLATT, M.
1928. SUR LA PRÉSENCE GÉNÉRALE DU SODIUM CHEZ LES PLANTES. [Paris] Acad. des Sci. Compt. Rend. 186: 200-202.
- (65) ——— and ROSENBLATT, M.
1932. SUR LA TENEUR INÉGALE EN MANGANÈSE DES FEUILLES VERTES ET DES FEUILLES ÉTIOLÉES. Soc. Chim. de France Bul. (4) 51: 862-864.
- (66) ——— SILBERSTEIN, LAZARE, and WAAL, HERMANUS L. DE
1936. NOUVELLES RÉCHERCHES SUR LES TENEURS COMPARATIVES EN BORE, SOUFRE, PHOSPHORE ET EN AZOTE DE PLANTES CULTIVÉES SUR LE MÊME SOL. Ann. Agron. [Paris] (n. s.) 6: 183-186, 537-541; 7: 333-335, 505-507.
- (67) ——— and VORONCA-SPIRT.
1929. RECHERCHES SUR LA PRÉSENCE ET LA RÉPARTITION DU TITANE DANS LES PLANTES PHANÉROGAMES. Soc. Chim. de France Bull. (4) 45: 1044-1052.
- (68) ——— and WAAL, HERMANUS L. DE.
1936. SUR LA TENEUR COMPARATIVE EN BORE DE PLANTES CULTIVÉES SUR LE MÊME SOL. [Paris] Acad. des Sci. Compt. Rend. 202: 605-609.
- (69) BETTINI, T. M.
1935. SULLA COMPOSIZIONE CHIMICA E SUL VALORE NUTRITIVO DELL'ERBA MEDICA A DIVERSI STADI DI VEGETAZIONE. Riv. di Zootec. 12: 425-438.
- (70) BIRCKNER, VICTOR.
1919. THE ZINC CONTENT OF SOME FOOD PRODUCTS. Jour. Biol. Chem. 38: 191-203.
- (71) BISHOP, EDNA.
1934. THE CALCIUM AND PHOSPHORUS CONTENT OF SOME ALABAMA VEGETABLES. Jour. Nutr. 8: 239-245.
- (72) ——— and LAWRENZ, MARGARET.
1932. COBALT IN PLANT ASH. Science 75: 264-265.
- (73) BISHOP, WILFRID B. S.
1928. THE DISTRIBUTION OF MANGANESE IN PLANTS AND ITS IMPORTANCE IN PLANT METABOLISM. Austral. Jour. Expt. Biol. and Med. Sci. 5: 125-141.
- (74) BISSET, NORMAN.
1934. CHRONIC COPPER POISONING IN SHEEP. Vet. Jour. 90: 405-407.
- (75) BLAIR, A. W., and PRINCE, A. L.
1932. THE INFLUENCE OF PHOSPHATES ON THE PHOSPHORIC ACID CONTENT OF THE PLANT. Jour. Agr. Res. 44: 579-590.
- (76) BLEYER, B.
1926. ZUR KENNTNIS DES JODS ALS BIOGENES ELEMENT. Biochem. Ztschr. 170: [265]-276.

- (77) **BLOM, I. J. B.**
1934. STUDIES IN MINERAL METABOLISM. XXIX. THE IODINE CONTENT OF FOODSTUFFS IN RELATION TO THE OCCURRENCE OF ENDEMIC GOITER IN THE LANGKLOOF VALLEY. XXX. VARIATIONS IN THE IODINE CONTENT OF GRASSES AT DIFFERENT STAGES OF GROWTH AND A NOTE ON THE IODINE CONTENT OF MILK. Onderstepoort Jour. Vet. Sci. and Anim. Indus. 2: 131-150.
- (78) **BOBKO, E. V., and ZERLING, V. V.**
1938. INFLUENCE DU BORE SUR LE DÉVELOPPEMENT REPRODUCTIF DES PLANTES. Ann. Agron. [Paris] (n. s.) 8: 174-184.
- (79) **BODE, G., and HEMBD, K.**
1921. ÜBER DEN MANGANGEHALT VON KORTOFFELN. Biochem. Ztschr. 124: 84-89.
- (80) **BOLIN, DONALD W.**
1934. THE MANGANESE CONTENT OF GRASSES AND ALFALFA FROM GRAZED PLOTS. Jour. Agr. Res. 48: 657-663.
- (81) **BONAME, P.**
1896. COMPOSITION MINÉRALE D'UNE RÉCOLTE DE CANNES. Mauritius Sta. Agron. Rap. Ann. 1896: 63-73.
- (82) **BONE, T.**
1930. INVESTIGATION OF RANCHING PROBLEMS OF THE WARANAMA SAVANNAH, BERBICE RIVER. Brit. Guiana Agr. Jour. 3: 160-163.
- (83) **BORDEN, R. J.**
1936. CANE GROWTH STUDIES. Hawaii. Planters Rec. 40: 143-156.
- (84) **BORST, H. L., and THATCHER, L. E.**
1931. LIFE HISTORY AND COMPOSITION OF THE SOYBEAN PLANT. Ohio Agr. Expt. Sta. Bul. 494, 96 pp.
- (85) **BOWMAN, HOWARD H. M., and YEE, MARTIN A.**
1926. THE NUTRITIVE VALUE OF THE MUNG BEAN. Mich. Acad. Sci., Arts, and Letters 6: 181-188.
- (86) **BRADLEY, W. B., BEATH, O. A., and EPPSON, H. F.**
1939. OAT HAY POISONING. Science 89: 365.
- (87) **BRENCHLEY, WINIFRED E.**
1936. THE ESSENTIAL NATURE OF CERTAIN MINOR ELEMENTS FOR PLANT NUTRITION. Bot. Rev. 2: 173-196.
- (88) _____
1937. SOME DEFICIENCY DISEASES OF CROP PLANTS. [Gt. Brit.] Min. Agr. and Fisheries Jour. 44: 116-122.
- (89) **BROWN, B. A.**
1932. THE EFFECTS OF FERTILIZATION ON THE CHEMICAL COMPOSITION OF VEGETATION IN PASTURES. Amer. Soc. Agron. Jour. 24: 129-145.
- (90) _____
1933. PASTURE INVESTIGATIONS (3RD REPORT). EFFECT OF FERTILIZERS ON THE BOTANICAL AND CHEMICAL COMPOSITION OF THE VEGETATION IN PERMANENT PASTURES. Conn. (Storrs) Agr. Expt. Sta. Bul. 187, 54 pp.
- (91) **BROWN, JANET W.**
1926. CHEMICAL STUDIES IN THE PHYSIOLOGY OF APPLES. V. METHODS OF ASH ANALYSIS, AND THE EFFECT OF ENVIRONMENT ON THE MINERAL CONSTITUTION OF THE APPLE. Ann. Bot. [London] 40: 129-147.
- (92) _____
1929. CHEMICAL STUDIES IN THE PHYSIOLOGY OF APPLES. XI. THE RELATION BETWEEN THE MINERAL CONSTITUTION OF APPLES AND THE SOIL ON WHICH THEY ARE GROWN. Ann. Bot. [London] 43: 817-831.
- (93) **BROWNE, C. A.**
1938. SOME RELATIONSHIPS OF SOIL TO PLANT AND ANIMAL NUTRITION. THE MAJOR ELEMENTS. SOILS AND MEN, U. S. Dept. Agr. Year-book, 1938: 777-806.
- (94) **BRYAN, O. C., and BECKER, R. B.**
1935. THE MINERAL CONTENT OF SOIL TYPES AS RELATED TO "SALT SICK" OF CATTLE. Amer. Soc. Agron. Jour. 27: 120-127.

- (95) BRYAN, O. C., and NEAL, W. M.
1936. THE INFLUENCE OF VARYING AMOUNTS OF WATER-SOLUBLE PHOSPHORUS IN DIFFERENT SOIL TYPES ON THE RESPONSE OF CULTIVATED CROPS. Jour. Agr. Res. 52: 459-466, illus.
- (96) BUCKNER, G. DAVIS.
1919. THE COMPOSITION OF THE ASH OF CRAB GRASS (*DIGITARIA SANGUINALIS*) AS AFFECTED BY THE SOIL IN WHICH IT IS GROWN. Amer. Chem. Soc. Jour. 41: 1384-1385.
- (97) BURGESS, PAUL S., and PEMBER, F. R.
1923. "ACTIVE" ALUMINUM AS A FACTOR DETRIMENTAL TO CROP PRODUCTION IN MANY ACID SOILS. R. I. Agr. Expt. Sta. Bul. 194, 40 pp.
- (98) BURKE, EDMUND, NYGARD, IVER J., and MARTIN, W. M.
1933. EXPERIMENTS WITH PHOSPHATE FERTILIZERS ON MONTANA SOILS. Mont. Agr. Expt. Sta. Bul. 280, 52 pp.
- (99) BYCZKOWSKI, ALEKSY.
1936. MECHANISM OF THE ACTION OF POTASH FERTILIZERS ON THE YIELD OF PLANTS (BARLEY). Polish Agr. Forest Ann. 37: 37-134.
- (100) BYERS, HORACE G.
1935. SELENIUM OCCURRENCE IN CERTAIN SOILS IN THE UNITED STATES WITH A DISCUSSION OF RELATED TOPICS. U. S. Dept. Agr. Tech. Bul. 482, 48 pp., illus.
- (101) ———
1936. SELENIUM OCCURRENCE IN CERTAIN SOILS IN THE UNITED STATES WITH A DISCUSSION OF RELATED TOPICS. Second report. U. S. Dept. Agr. Tech. Bul. 530, 79 pp., illus.
- (102) ———
1937. SELENIUM IN MEXICO. Indus. and Engin. Chem. 29: 1200-1202.
- (103) ——— and KNIGHT, HENRY G.
1935. SELENIUM IN SOILS IN RELATION TO ITS PRESENCE IN VEGETATION. Indus. and Engin. Chem. 27: 902-904.
- (104) ——— MILLER, JOHN T., WILLIAMS, K. T., and LAKIN, H. W.
1938. SELENIUM OCCURRENCE IN CERTAIN SOILS IN THE UNITED STATES WITH A DISCUSSION OF RELATED TOPICS. Third report. U. S. Dept. Agr. Tech. Bul. 601, 75 pp., illus.
- (105) CARLSTROM, B., and HJÄRRE, A.
1938. DIE BEDEUTUNG DER FÜTTERUNG ALS PRÄDISPONIERENDEN FAKTORS BEI DER INFEKTIOSEN ANÄMIE. 13th Internatl. Tierärztl. Kong., Zürich-Interlaken, Switzerland, 1: 287-305.
- (106) CARLYLE, E. C.
1931. MANGANESE IN TEXAS SOILS AND ITS RELATION TO CROPS. Tex. Agr. Expt. Sta. Bul. 432, 37 pp.
- (107) CAROLUS, R. L.
1933. TOMATO FERTILIZATION. II. EFFECT OF DIFFERENT FERTILIZER RATIOS ON THE CHEMICAL COMPOSITION OF TOMATOES. Va. Truck Expt. Sta. Bul. 81, pp. 1085-1117.
- (108) ———
1933. SOME SIGNIFICANT VARIATIONS IN THE CHEMICAL COMPOSITION OF THE PLANT ASSOCIATED WITH A MALNUTRITION TROUBLE OF POTATOES. Am. Potato Jour. 10: 147-164.
- (109) ———
1935. EFFECTS OF MAGNESIUM DEFICIENCY IN THE SOIL ON THE YIELD, APPEARANCE AND COMPOSITION OF THE VEGETABLE CROPS. Amer. Soc. Hort. Sci. Proc. (1934) 31: 610-614.
- (110) CARPENTER, F. B.
1895. TYPES OF TOBACCO AND THEIR ANALYSES. N. C. Agr. Expt. Sta. Bul. 122, pp. 331-366.
- (111) CARY, C. A.
1920. DEFICIENCY DISEASES. Amer. Vet. Med. Assoc. Jour. 56: 609-614.
- (112) CHATIN.
1851. [Paris] Acad. des Sci. Compt. Rend. 33: 529-531, 584-585.
- (113) CHUNG, H. L., and RIPPERTON, J. C.
1929. UTILIZATION AND COMPOSITION OF ORIENTAL VEGETABLES IN HAWAII. Hawaii Agr. Expt. Sta. Bull. 60, 64 pp.

- (114) CLAYTON, JOHN.
1688. AN ACCOUNT OF SEVERAL OBSERVABLES IN VIRGINIA. Roy Soc. London, Phil. Trans. 17: 943.
- (115) CLEVELAND, M. M., and FELLERS, C. R.
1932. MINERAL COMPOSITION OF DATES. Indus. and Engin. Chem., Analyt. Ed. 4: 267-268.
- (116) COHEN, N. H.
1914. TOBAKSGRONDEN EN DE DAAROP GEGROEIDE TABAK. Proefstat. v. Vorstenland. Tabak Meded., No. 11, 18 pp.
- (117) COLBY, GEORGE E.
1894. ANALYSES OF BANANAS AND BANANA SOILS FROM THE SANDWICH ISLANDS. Calif. Agr. Expt. Sta. Rpt. 1892-93: 275-279.
- (118) COLE, VERSA V., CURTIS, GEO. M., and BONE, MARY L.
1934. THE IODINE CONTENT OF HOSPITAL FOODS. Amer. Dietet. Assoc. Jour. 10: 200-207.
- (119) COLEMAN, J. M., and RUPRECHT, R. W.
1935. THE EFFECT OF FERTILIZERS AND SOIL TYPES ON THE MINERAL CONTENT OF PLANTS. Jour. Nutr. 9: 51-62.
- (120) COOK, F. C.
1916. BORON: ITS ABSORPTION AND DISTRIBUTION IN PLANTS AND ITS EFFECT ON GROWTH. Jour. Agr. Res. 5 (19): 877-890.
- (121) ——— and WILSON, J. B.
1917. EFFECT OF THREE ANNUAL APPLICATIONS OF BORON ON WHEAT. Jour. Agr. Res. 10: 591-597.
- (122) COOPER, H. P., and WILSON, J. K.
1930. RELATION OF ASH CONSTITUENTS OF PASTURE PLANTS TO THE OXIDATION-REDUCTION POTENTIALS OF NUTRIENTS. Soil Sci. 30: 421-430.
- (123) CORLETTE, C. E.
1928. OBSERVATIONS ON CALCIUM DEFICIENCY AS A CAUSE OF CERTAIN DISEASES OF ANIMALS AND MAN. Austral. Med. Jour. 1: 198-215.
- (124) CORNER, H., and SMITH, ALEXANDER M.
1938. THE INFLUENCE OF COBALT ON PINE DISEASE IN SHEEP. Biochem. Jour. 32: 1800-1805.
- (125) COWELL, STUART JASPER.
1932. A NOTE ON THE CALCIUM CONTENT OF CABBAGE. Biochem. Jour. 26: 1422-1423.
- (126) CRAMPTON, E. W., and FINLAYSON, D. A.
1935. PASTURE STUDIES VII. THE EFFECT OF FERTILIZATION ON THE NUTRITIVE VALUE OF PASTURE GRASS. Empire Jour. Expt. Agr. 3: 331-345.
- (127) CRAWFORD, M.
1927. THE INFLUENCE OF THE VARIOUS MINERAL CONSTITUENTS ON ANIMAL NUTRITION, AND THE EFFECTS OF DEFICIENCIES AND EVIDENCE OF SUCH IN CEYLON. Trop. Agr. [Ceylon] 68: 272-281.
- (128) CUNDY, A. T.
1924. WYOMING FORAGE PLANTS AND THEIR CHEMICAL COMPOSITION. Wyo. Agr. Expt. Sta. Bul. 137, 16 pp.
- (129) CUNNINGHAM, IRA JAMES.
1931. SOME BIOCHEMICAL AND PHYSIOLOGICAL ASPECTS OF COPPER IN ANIMAL NUTRITION. Biochem. Jour. 25: 1267-1293.
- (130) DAMIENS, A., and BLAIGNAN, S.
1931. SUR LE BROME NORMAL (RÈGNE VÉGÉTAL): GRAINES COMESTIBLES, BLÉ, PAIN. [Paris] Acad. des Sci. Compt. Rend. 193: 1460-1462.
- (131) ——— and BLAIGNAN, S.
1932. SUR LE BROME NORMAL (RÈGNE VÉGÉTAL): PLANTES ET FRUITS COMESTIBLES. [Paris] Acad. des Sci. Compt. Rend. 194: 2077-2080.
- (132) DANIEL, HARLEY A.
1934. THE CALCIUM, PHOSPHORUS AND NITROGEN CONTENT OF GRASSES AND LEGUMES, AND THE RELATION OF THESE ELEMENTS IN THE PLANT. Amer. Soc. Agron. Jour. 26: 496-503.

- (133) DANIEL, HARLEY A.
1935. THE TOTAL CALCIUM, PHOSPHORUS, AND NITROGEN CONTENT OF NATIVE AND CULTIVATED PLANTS IN THE HIGH PLAINS OF OKLAHOMA AND A STUDY OF THE MINERAL DEFICIENCIES THAT MAY DEVELOP IN LIVESTOCK WHEN EMERGENCY FEEDS ARE FED. [Okla.] Agr. Expt. Sta., Panhandle Bul. 56, 18 pp.
- (134) ——— and HARPER, HORACE J.
1934. THE RELATION BETWEEN TOTAL CALCIUM AND PHOSPHORUS IN MATURE PRAIRIE GRASS AND AVAILABLE PLANT FOOD IN THE SOIL. Amer. Soc. Agron. Jour. 26: 986-992.
- (135) ——— and HARPER, HORACE J.
1935. THE RELATION BETWEEN EFFECTIVE RAINFALL AND TOTAL CALCIUM AND PHOSPHORUS IN ALFALFA AND PRAIRIE HAY. Amer. Soc. Agron. Jour. 27: 644-652.
- (136) DARKIS, F. R., DIXON, L. F., WOLF, F. A., and GROSS, P. M.
1936. FLUE-CURED TOBACCO. CORRELATION BETWEEN CHEMICAL COMPOSITION AND STALK POSITION OF TOBACCOS PRODUCED UNDER VARYING WEATHER CONDITIONS. Indus. and Engin. Chem. 28: 1214-1223.
- (137) DASZEWSKI, A. VON, and TOLLENS, B.
1900. DER EINFLUSS DES WASSERS UND DER DÜNGUNG AUF DIE ZUSAMMENSETZUNG DER ASCHEN DER KARTOFFELPFLANZE. Jour. f. Landw. 48: 223-249.
- (138) DAVIDSON, JEHIEL.
1929. MANGANESE IN CEREALS AND CEREAL MILL PRODUCT. Cereal Chem. 6: 128-133.
- (139) ——— and CHAMBLISS, C. E.
1932. CHEMICAL COMPOSITION OF RICE AND ITS RELATION TO SOIL FERTILITY IN CHINA AND JAPAN. Science 75: 294.
- (140) ——— and LÉCLERC, J. A.
1936. THE VARIATION IN THE MINERAL CONTENT OF VEGETABLES. Jour. Nutr. 11: 55-66.
- (141) DAVIDSON, R. J.
1895. ANALYSES OF PARTS OF TOBACCO PLANT AT DIFFERENT STAGES OF GROWTH. Va. Agr. Expt. Sta. Bul. 50, pp. 35-52.
- (142) ———
1898. THE INFLUENCE OF COMMERCIAL FERTILIZERS UPON THE QUALITY OF IRISH POTATOES. Va. Agr. Expt. Sta. Bul. 92, pp. 97-108.
- (143) DAVIS, WILLIAM A.
1917. THE PHOSPHATE DEPLETION OF SOILS OF BIHAR AND ITS CONTINGENT RISKS OF MALNUTRITION AND ENDEMIC DISEASE: A WARNING. Agr. Jour. India 12: 181-184.
- (144) DAWSON, CHARLES F.
1906. BOVINE UNCINARIASIS (SALT SICK). Fla. Agr. Expt. Sta. Bul. 86, 14 pp.
- (145) DELBET, PIERRE.
1934. AGRICULTURE, CUISINE ET MAGNÉSIUM. Acad. de Med. [Paris] Bul. 111: 393-415.
- (146) ——— and ROBINET, L.
1934. TERRAINS MAGNÉSIENS ET CANCER. Acad. de Med. [Paris] Bul. 111: 415-429.
- (147) DELONG, W. A.
1937. CALCIUM AND BORON CONTENTS OF THE APPLE FRUIT AS RELATED TO THE INCIDENCE OF BLOTCHY CORK. Plant Physiol. 12: 553-556.
- (148) DENHAM, H. G.
1937. COBALT INVESTIGATIONS IN NEW ZEALAND. Science 85: 383.
- (149) DENSCH, A., and HUNNIUS, T.
1924. VERSUCHE MIT KUPFERSULFAT. Ztschr. f. Pflanzenernähr., Düngung u. Bodenk. 3A: 369-386.
- (150) ——— STEINFATT and GUNTHER.
1929. EIN DÜNGUNGSVERSUCH MIT NATRONALSALPETER, CHILESALPETER UND JOD ZU MÖHREN. Ztschr. f. Pflanzenernähr., Düngung u. Bodenk. (B)8: 205-211.
- (151) DEOMANO, FRANCISCO V.
1931. A STUDY OF THE CHEMICAL COMPOSITION OF FOUR SUGAR CANE VARIETIES OF THE SAME AGE AND GROWN UNDER SIMILAR CONDITIONS. Philippine Agr. 20: 139-153.

- (152) DEPARDON.
1938. INFLUENCE DU SOL SUR LA COMPOSITION DES RÉCOLTES. Assoc. Franc. pour l'Etude du Sol, Bul. 4: 104, 132-140.
- (153) DIETZ, CARL.
1938. IODINE CONTENT OF SOME OHIO VEGETABLES. Ohio Veg. Growers Assoc. Proc. 23: 57-63. [The same data appear under the same title in: Food Res. 3: 359-365 (1938).]
- (154) DINGWALL, ANDREW, and BEANS, H. T.
1934. STUDIES ON CHROMIUM. III. THE OCCURRENCE OF CHROMIUM IN CERTAIN PLANTS IN THE PROVINCE OF QUEBEC. Amer. Chem. Soc. Jour. 56: 1666-1667.
- (155) ——— McKIBBIN, R. R., and BEANS, H. T.
1934. STUDIES ON THE DISTRIBUTION OF MOLYBDENUM IN BIOLOGICAL MATERIAL. I. A SPECTROGRAPHIC STUDY OF THE OCCURRENCE OF MOLYBDENUM IN PLANTS GROWN IN THE PROVINCE OF QUEBEC. Canad. Jour. Res. 11: 32-39.
- (156) DODD, A. S.
1927. THE NATURAL OCCURRENCE OF BORON COMPOUNDS IN CACAO AND CACAO PRODUCTS. Analyst 52: 459-466.
- (157) ———
1929. THE NATURAL OCCURRENCE OF BORON COMPOUNDS IN FRUITS AND VEGETABLE PRODUCTS. Analyst 54: 15-22.
- (158) DOHERTY, W. M.
1892. THE ANALYSIS OF THE CAVENDISH BANANA (*MUSA CAVENDISHII*) IN RELATION TO ITS VALUE AS A FOOD. Chem. News 66: 187-188.
- (159) DUDLEY, H. C.
1939. BROMIDE CONTENT OF FRUITS AND VEGETABLES FOLLOWING FUMIGATION WITH METHYL BROMIDE. Indus. and Engin. Chem., Analyt. Ed. 11: 259-261.
- (160) EATON, FRANK M.
1935. BORON IN SOILS AND IRRIGATION WATERS AND ITS EFFECT ON PLANTS, WITH PARTICULAR REFERENCE TO THE SAN JOAQUIN VALLEY OF CALIFORNIA. U. S. Dept. Agr. Tech. Bul. 448, 132 pp., illus.
- (161) ECKLES, C. H., BECKER, R. B., and PALMER, L. S.
1926. A MINERAL DEFICIENCY IN THE RATIONS OF CATTLE. Minn. Agr. Expt. Sta. Bul. 229, 49 pp., illus.
- (162) ——— GULLICKSON, T. W., and PALMER, L. S.
1932. PHOSPHORUS DEFICIENCY IN THE RATIONS OF CATTLE. Minn. Agr. Expt. Sta. Tech. Bul. 91, 118 pp.
- (163) EDDY, WALTER H.
1933. THE NUTRITIVE VALUE OF THE BANANA. 37 pp. Bur. of Pub. Teachers Col., Columbia Univ.
- (164) ELLIOT, WALTER, ORR, J. B., and WOOD, T. B.
1926. INVESTIGATION ON THE MINERAL CONTENT OF PASTURE GRASS AND ITS EFFECT ON HERBIVORA. Jour. Agr. Sci. [England] 16: 59-64.
- (165) ——— ORR, J. B., and WOOD, T. B.
1926. INVESTIGATION ON THE MINERAL CONTENT OF PASTURE GRASS AND ITS EFFECT ON HERBIVORA. Jour. Agr.; Sci. [England] 16: 78-88.
- (166) ELVEHJEM, C. A., and HART, E. B.
1929. THE COPPER CONTENT OF FEEDING STUFFS. Jour. Biol. Chem. 82: 473-477.
- (167) EMMERT, E. M.
1931. THE EFFECT OF SOIL REACTION ON THE GROWTH OF TOMATOES AND LETTUCE AND ON THE NITROGEN, PHOSPHORUS AND MANGANESE CONTENT OF THE SOIL AND PLANT. Ky. Agr. Expt. Sta. Bul. 314, 83 pp.
- (168) ENGELS, O.
1938. SPURENELEMENTE UND IHRE BEDEUTUNG FÜR DAS PFLANZENWACHSTUM. Chem. Ztg. 62: 165-167.
- (169) EVANS, R. J., and GREAVES, J. E.
1937. SOME FACTORS INFLUENCING THE SULFUR CONTENT OF ALFALFA. Utah Acad. Sci., Arts and Letters Proc. 14: 17-22.

- (170) FAGAN, T. W.
1928. FACTORS THAT INFLUENCE THE CHEMICAL COMPOSITION OF HAY. Welsh Jour. Agr. 4: 92-102.
- (171) ——— and MILTON, W. E. J.
1931. THE CHEMICAL COMPOSITION OF ELEVEN SPECIES AND STRAINS OF GRASSES AT DIFFERENT STAGES OF MATURITY. Welsh Jour. Agr. 7: 246-255.
- (172) ——— MILTON, W. E. J., and PROVAN, A. L.
1928. THE EFFECT OF NITRATE OF SODA ON THE YIELD AND CHEMICAL COMPOSITION OF A SIMPLE SEEDS MIXTURE IN THE FIRST HARVEST YEAR UNDER DIFFERENT SYSTEMS OF MANAGEMENT. Welsh Plant Breeding Sta., Aberystwyth, Ser. H., No. 9 (1926-27) 27 pp.
- (173) FAVILLE, GEO. C.
1896. OSTEOPOROSIS. Amer. Vet. Rev. 20: 28-39.
- (174) FELLEBERG, THEO. VON.
1923. UNTERSUCHUNGEN UEBER DAS VORKOMMEN VON JOD IN DER NATUR. I. Biochem. Ztschr. 139: 371-451.
- (175) ———.
1924. UNTERSUCHUNGEN UEBER DAS VORKOMMEN VON JOD IN DER NATUR. Mitt. aus dem Geb. der Lebensmtl. Untersuch. u. Hyg. 15: 245-248.
- (176) ———.
1924. UNTERSUCHUNGEN UEBER DAS VORKOMMEN VON JOD IN DER NATUR. VI. UEBER DEN ZUSAMMENHANG ZWISCHEN DER HÄUFIGKEIT DES AUFTRETENS VON KROPF UND DEM JODEGHALT DER UNWELT. Biochem. Ztschr. 152: 141-152.
- (177) ———.
1926. DAS VORKOMMEN, DER KREISLAUF UND DER STOFFWECHSEL DES JODS. Ergeb. der Physiol. 25: 176-363.
- (178) ———.
1929. UEBER DEN ARSENGEHALT NATÜRLICHER UND MIT ARSENPRÄPARATEN BEHANDELTER LEBENSMITTEL. Mitt. aus dem Geb. der Lebensmtl. Untersuch. u. Hyg. 20: 338-354.
- (179) FELLERS, C. R., and SMITH, E. G.
1936. CHEMICAL COMPOSITION AND FERMENTATION STUDIES OF CITRON. Jour. Agr. Res. 53: 859-867.
- (180) FERGUSON, W. S., LEWIS, A. H., and WATSON, S. J.
1938. ACTION OF MOLYBDENUM IN NUTRITION OF MILKING CATTLE Nature [London] 141: 553.
- (181) FIELD, ADA M., PEACOCK, M. T., COX, EDITH, and EARLE, I. P.
1928. CHEMICAL COMPOSITION OF MUSTARD AND TURNIP GREENS AND LOSSES OF IRON IN COOKING. West. Hosp. and Nurses Rev. 11: 26-28.
- (182) FILMER, J. F.
1933. ENZOÏTIC MARASMUS OF CATTLE AND SHEEP. Austral. Vet. Jour. 9: 163-179.
- (183) ——— and UNDERWOOD, E. J.
1934. TREATMENT WITH LIMONITE FRACTIONS. Austral. Vet. Jour. 10: 83-92.
- (184) ——— and UNDERWOOD, E. J.
1936. WASTING DISEASE. DIAGNOSIS, PREVENTION AND TREATMENT. West Austral. Dept. Agr. Jour. 13: 199-201.
- (185) FLEURENT, E., and LEVI, LUCIEN.
1920. SUR LA PRÉSENCE DU CUIVRE DANS L'ORGANISME VÉGÉTAL ET ANIMAL. Soc. Chim. de France Bul. 27: 441-442.
- (186) FLICHE, P., and GRANDEAU, L.
1873. DE L'INFLUENCE DE LA COMPOSITION CHIMIQUE DU SOL SUR LA VÉGÉTATION DU PIN MARITIME (PINUS PINASTER, SOLAND). Ann. de Chim. et de Phys. (4) 29: 383.
- (187) FONDARD, LOUIS, and MARGAILLAN, LOUIS.
1928. RÉCHERCHES SUR LA COMPOSITION DE L'ÉPINARD. Acad. d'Agr. de France Compt. Rend. 14: 1265-1270.
- (188) FONDER, JOHN F.
1929. A CRITICAL STUDY OF THE INFLUENCE OF SOIL TYPE ON THE CALCIUM AND MAGNESIUM CONTENT AND OTHER PHYSIOLOGICAL CHARACTERS OF THE ALFALFA PLANT. Soil Sci. 27: 205-232, illus.

- (189) FONDER, JOHN F.
1929. THE RELATIONSHIP OF SOIL TYPE TO THE CALCIUM AND MAGNESIUM CONTENT OF GREEN BEAN STEMS AND LEAVES AND OF THEIR EXPRESSED JUICE. *Soil Sci.* 27: 415-431.
- (190) ———
1929. VARIATIONS IN THE CALCIUM AND MAGNESIUM CONTENTS OF PEA PLANTS ON DIFFERENT SOIL TYPES. *Soil Sci.* 28: 15-26.
- (191) ———
1929. VARIATIONS IN POTASSIUM CONTENT OF ALFALFA DUE TO STAGE OF GROWTH AND SOIL TYPE AND THE RELATIONSHIP OF POTASSIUM AND CALCIUM IN PLANTS GROWN UPON DIFFERENT SOIL TYPES. *Amer. Soc. Agron. Jour.* 21: 732-750.
- (192) FORBES, E. B., BEEGLE, F. M., and MENSCHING, J. E.
1913. MINERAL AND ORGANIC ANALYSES OF FOODS. *Ohio Agr. Expt. Sta. Bul.* 255, pp. 211-231.
- (193) ——— and JOHNSON, S. R.
1939. PHOSPHORUS DEFICIENCY AMONG CATTLE IN PENNSYLVANIA. *Pa. Agr. Expt. Sta. Bul.* 371, 25 pp.
- (194) ——— and KEITH, M. HELEN
1914. A REVIEW OF THE LITERATURE OF PHOSPHORUS COMPOUNDS IN ANIMAL METABOLISM. *Ohio Agr. Expt. Sta. Tech. Bul.* 5, 583 pp.
- (195) ——— WHITTIER, A. C., and COLLISON, R. C.
1910. THE MINERAL NUTRIENTS IN BLUEGRASS. *Ohio Agr. Expt. Sta. Bul.* 222, pp. 39-53.
- (196) FRANKE, KURT W., and PAINTER, E. PAGE.
1937. EFFECT OF SULFUR ADDITIONS ON SELENIFEROUS SOILS. BINDING OF SELENIUM BY SOIL. *Indus. and Engin. Chem.* 29: 591-595.
- (197) ——— RICE, T. D., JOHNSON, A. G., and SCHOENING, H. W.
1934. REPORT ON A PRELIMINARY FIELD SURVEY OF THE SO-CALLED "ALKALI DISEASE" OF LIVESTOCK. *U. S. Dept. Agr. Cir.* 320, 10 pp., illus.
- (198) FREAR, DONALD E. H.
1934. A STUDY OF THE IODINE CONTENT OF PENNSYLVANIA POTATOES. *Jour. Agr. Res.* 48: 171-182, illus.
- (199) FUDGE, B. R.
1938. MAGNESIUM DEFICIENCY IN RELATION TO YIELD AND CHEMICAL COMPOSITION OF SEEDY AND COMMERCIALY SEEDLESS VARIETIES OF GRAPEFRUIT. *Fla. Proc. State Hort. Soc.* 51: 34-43.
- (200) FULMER, H. L.
1916. SWEET CLOVER. *Ontario Dept. Agr. Bul.* 235, 32 pp.
- (201) GARNER, W. W., McMURTREY, J. E., Jr., BOWLING, J. D. Jr., and MOSS, E. G.
1930. MAGNESIUM AND CALCIUM REQUIREMENTS OF THE TOBACCO CROP. *Jour. Agr. Res.* 40: 145-168.
- (202) ——— McMURTREY, J. E., Jr., BOWLING, J. D., and MOSS, E. G.
1930. RÔLE OF CHLORINE IN NUTRITION AND GROWTH OF THE TOBACCO PLANT AND ITS EFFECT ON THE QUALITY OF THE CURED LEAF. *Jour. Agr. Res.* 40: 627-648, illus.
- (203) GAUTIER, ARMAND, and CLAUSMAN, PAUL.
1916. LE FLUOR DANS LE RÈGNE VÉGÉTAL. [*Paris*] *Acad. des Sci. Compt. Rend.* 162: 105-112.
- (204) GEILMAN, W.
1920. ÜBER DIE VERBREITUNG DES TITANS IN BÖDEN UND PFLANZEN. *Jour. f. Landw.* 68: 107-124.
- (205) ——— and BRUNGER, K.
1935. ÜBER DIE AUFNAHME VON GERMANIUM DURCH PFLANZEN. *Biochem. Ztschr.* 275: 387-395.
- (206) GIESECKE, F., SCHMALFUSS, K., and RATHJE, W.
1938. VERSUCHE ÜBER DIE SPURENELEMENTWIRKUNG DES CHILESALPETERS IN GEFÄSSKULTUREN. *Bodenk. w. Pflanzenernähr.* 9-10: 580-587.
- (207) GILBERT, B. E., McLEAN, F. T., and HARDIN, LEO J.
1926. THE RELATION OF MANGANESE AND IRON TO LIME INDUCED CHLOROSIS. *Soil Sci.* 22: 437-446.

- (208) GILE, P. L.
1911. RELATION OF CALCAREOUS SOILS TO PINEAPPLE CHLOROSIS. P. R. Agr. Expt. Sta. Bul. 11, 45 pp.
- (209) ——— and AGETON, C. N.
1914. THE EFFECT OF STRONGLY CALCAREOUS SOILS ON THE GROWTH AND ASH COMPOSITION OF CERTAIN PLANTS. P. R. Agr. Expt. Sta. Bul. 16, 45 pp.
- (210) GILRUTH, J. A.
1900. "BUSH" OR "TAURANGA" DISEASE IN CATTLE AND SHEEP. New Zeal. Dept. Agr. Ann. Rpt. 8: 187.
- (211) GODDEN, WILLIAM.
1926. INVESTIGATION ON THE MINERAL CONTENT OF PASTURE GRASS AND ITS EFFECT ON HERBIVORA. III. REPORT ON THE CHEMICAL ANALYSES OF SAMPLES OF PASTURE FROM VARIOUS AREAS IN THE BRITISH ISLES. Jour. Agr. Sci. [England] 16: 78-88.
- (212) ——— and GRIMMETT, R. E. R.
1928. FACTORS AFFECTING THE IRON AND MANGANESE CONTENT OF PLANTS WITH SPECIAL REFERENCE TO HERBAGE CAUSING "PINING" AND "BUSH-SICKNESS." Jour. Agr. Sci. [England] 18: 363-368.
- (213) GODLEWSKI, E.
1901. NÄHRSTOFFBEDURFNIS EINIGER KULTURPFLANZEN. Ztschr. Landw. Versuchsw. Österr. 4: 479-536.
- (214) GOESSMANN, C. A.
1891. COMPOSITION OF CROPS. Mass. Agr. Expt. Sta. Ann. Rpt. (1890) 8: 159-186.
- (215) ———
1894. NEW FORAGE CROPS. Mass. Agr. Expt. Sta. Ann. Rpt. (1893) 11: 212-219.
- (216) ———
1895. FORAGE AND VEGETABLE CROPS. Mass. Agr. Expt. Sta. Ann. Rpt. (1894) 12: 231-255; 398-399.
- (217) GOHREN, THEODOR VON.
1861. UEBER EINE BEIGABE VON KNOCHENERDE ZUM FUTTER DER THEIRE NEBST EINEM PHYSIOLOGISCH-CHEMISCHEN VERSUCH UEBER DEREN VERDAULICHKEIT. Landw. Vers. Sta. 3: 161-178.
- (218) GRANGE, L. I., TAYLOR, N. H., RIGG, T., and HODGSON, L.
1932. THE OCCURRENCE OF BUSH SICKNESS ON THE VOLCANIC SOILS OF THE NORTH ISLAND. New Zeal. Dept. Sci. and Indus. Res. Bul. 32: 21-51.
- (219) GREAVES, ETHELYN O., and GREAVES, J. E.
1933. NUTRITIVE VALUE OF HIGH AND LOW CALCIUM-CARRYING WHEAT. Jour. Nutr. 6: 113-125.
- (220) GREAVES, J. E., and ANDERSON, ARIEL.
1936. THE INFLUENCE OF SOIL AND VARIETY ON THE COPPER CONTENT OF GRAINS. Jour. Nutr. 11: 111-118.
- (221) ——— and CARTER, E. G.
1923. THE INFLUENCE OF IRRIGATION WATER ON THE COMPOSITION OF GRAINS AND THE RELATIONSHIP TO NUTRITION. Jour. Biol. Chem. 58: 531-541.
- (222) ——— and HIRST, C. T.
1929. THE MINERAL CONTENT OF GRAINS. Utah Agr. Expt. Sta. Bul. 210, 1-38.
- (223) ——— and NELSON, D. H.
1925. THE INFLUENCE OF IRRIGATION WATER AND MANURE ON THE COMPOSITION OF THE CORN KERNEL. Jour. Agr. Res. 31: 183-189.
- (224) ——— and NELSON, D. H.
1925. THE IRON, CHLORINE AND SULFUR CONTENTS OF GRAINS AND THE INFLUENCE OF IRRIGATION WATER UPON THEM. Soil Sci. 19: 325-330.
- (225) GREEN, H. H.
1935. MINERALS IN RELATION TO DISEASE OF THE LARGER DOMESTICATED ANIMALS. Empire Jour. Expt. Agr. 3: 363-378.
- (226) GREIG, J. RUSSELL, DRYERRE, HENRY, GODDEN, W., CRICHTON, A., and OGG W. G.
1933. "PINE": A DISEASE AFFECTING SHEEP AND YOUNG CATTLE. Vet. Jour. 89: 99-110.

- (227) GRENDEL, F.
1930. OVER HET KOPERGEHALTE VAN EENIGE VOEDINGSMIDDELEN. Pharm. Weekbl. 67: 913-921.
- (228) GREW, N.
1682. THE ANATOMY OF PLANTS WITH AN IDEA OF A PHILOSOPHICAL HISTORY OF PLANTS AND SEVERAL OTHER LECTURES READ BEFORE THE ROYAL SOCIETY. 304 pp., illus. London.
- (229) GRIZZARD, A. L.
1935. EFFECTS OF SOIL TYPE AND SOIL TREATMENT ON THE CHEMICAL COMPOSITION OF ALFALFA PLANTS. Amer. Soc. Agron. Jour. 27: 81-99.
- (230) GROUVEN, H.
1868-69. Jahresber. Agr. Chem. 11-12: 544.
- (231) GRUNDER, MAYNARD S.
1933. YIELD AND CHEMICAL COMPOSITION OF CERTAIN PASTURE CROPS, FERTILIZED AND UNFERTILIZED. Jour. Agr. Res. 46: 375-386.
- (232) GUERITHAULT, B.
1920. SUR LA PRÉSENCE DU CUIVRE DANS LES PLANTES ET PARTICULIÈREMENT DANS LES MATIÈRES ALIMENTAIRES D'ORIGINE VÉGÉTALE. [Paris] Acad. des Sci. Compt. Rend. 171: 196-198.
- (233) GUILBERT, H. R., MEAD, S. W., and JACKSON, H. G.
1931. THE EFFECT OF LEACHING ON THE NUTRITIVE VALUE OF FORAGE PLANTS. Hilgardia 6: 13-26.
- (234) GUTHRIE, F. B., HENRY, MAX, JENSEN, H. I., and RAMSAY, A. A.
1914. NOTES ON OSTEOMALACIA. N. S. Wales Dept. Agr. Sci. Bul. 12, 23 pp.
- (235) HAAS, A. R. C., and BATCHELOR, L. D.
1928. RELATION OF PHOSPHORUS CONTENT TO SHRIVELLING OF WALNUT KERNELS. Bot. Gaz. 86: 448-455.
- (236) ——— and QUAYLE, H. J.
1935. COPPER CONTENT OF CITRUS LEAVES AND FRUIT IN RELATION TO EXANTHEMA AND FUMIGATION INJURY. Hilgardia 9: 143-177.
- (237) HALL, E. H.
1922. SULFUR AND NITROGEN CONTENT OF ALFALFA GROWN UNDER VARIOUS CONDITIONS. Bot. Gaz. 73: 401-411.
- (238) HARDING, E. P., and DYSTERHEFT, GEORGE.
1927. A DETERMINATION OF IRON, CALCIUM, MAGNESIUM, PHOSPHORUS, ASH AND PROTEIN IN HARD SPRING WHEAT AND IN THE FLOUR STREAMS REPRESENTING THE LARGEST VOLUME OF FLOUR PRODUCED IN ITS MILLING. Cereal Chem. 4: 47-57.
- (239) HARPER, H. J., DANIEL, H. A., and VOLK, G. W.
1938. FERTILIZING CONSTITUENTS OF COTTON BURS OR COTTON BUR ASHES AND THEIR EFFECT ON CROP YIELDS. Amer. Soc. Agron. Jour. 30: 827-832.
- (240) HARRINGTON, H. H.
1890. MISCELLANEOUS ANALYSES. Tex. Agr. Expt. Sta. Bul. 13: 29-40.
- (241) HARRIS, F. S., and PITTMAN, D. W.
1917. IRRIGATION AND MANURING STUDIES. II. THE EFFECT OF VARYING QUANTITIES OF IRRIGATION WATER AND MANURE ON THE GROWTH AND YIELD OF CORN. Utah Agr. Expt. Sta. Bul. 154, 29 pp.
- (242) HART, E. B., BEACH, B. A., DELWICHE, E. J., and BAILEY, E. G.
1927. PHOSPHORUS DEFICIENCY AND A DAIRY CATTLE "DISEASE." Wis. Agr. Expt. Sta. Bul. 389, 10 pp., illus.
- (243) ——— and ELVEHJEM, C. A.
1936. MINERAL METABOLISM. Ann. Rev. Biochem. 5: 271-294.
- (244) ——— PHILLIPS, P. H., and BOHSTEDT, G.
1934. RELATION OF SOIL FERTILIZATION WITH SUPERPHOSPHATES AND ROCK PHOSPHATES TO FLUORINE CONTENT OF PLANTS AND DRAINAGE WATERS. Amer. Jour. Pub. Health 24: 936-940.
- (245) HART, G. H., and GUILBERT, H. R.
1928. FACTORS INFLUENCING PERCENTAGE CALF CROP IN RANGE HERDS. Calif. Agr. Expt. Sta. Bul. 458, 43 pp., illus.
- (246) ——— GUILBERT, H. R., and GOSS, H.
1932. SEASONAL CHANGES IN THE CHEMICAL COMPOSITION OF RANGE FORAGE AND THEIR RELATION TO NUTRITION OF ANIMALS. Calif. Agr. Expt. Sta. Bul. 543, 62 pp., illus.

- (247) HARTWELL, B. L.
1932. THE INFLUENCE OF FERTILIZERS ON CROP QUALITY. *Nat. Fert. Assoc., Ann. Conv. (Sect. 2):* 5-60. Proc. 8.
- (248) ———
1913. THE PERCENTAGE OF TOTAL PHOSPHORUS IN FLAT TURNIPS AS INFLUENCED BY THE AMOUNT AVAILABLE IN SOILS. *R. I. Agr. Expt. Sta. Bul.* 154, pp. 121-148.
- (249) HASELHOFF, E.
1893. VERSUCHE ÜBER DEN ERSATZ DES KALKES DURCH STRONTIAN BEI DER PFLANZENERNÄHRUNG. *Landw. Jahrb.* 22: 851-861.
- (250) ——— and ST. WERNER.
1913. ÜBER DIE VERÄNDERUNGEN IN DER ZUSAMMENSETZUNG DER ROTKLEEPFLANZE IN VERSCHIEDENEN WACHSTUMS-STADIEN. *Landw. Jahrb.* 44: 651-680.
- (251) HASKINS, H. D.
1906. COMPILATION OF ANALYSES OF FRUITS, GARDEN CROPS AND INSECTICIDES. *Mass. Agr. Expt. Sta. Ann. Rpt. (1905)* 18: 184-199.
- (252) HAUT, I. C., WEBSTER, J. E., and COCHRAN, G. W.
1936. THE INFLUENCE OF COMMERCIAL FERTILIZERS UPON THE FIRMNESS AND CHEMICAL COMPOSITION OF STRAWBERRIES AND TOMATOES. *Amer. Soc. Hort. Sci. Proc. (1935)* 33: 405-410.
- (253) HEADDEN, W. P.
1897. A STUDY OF ALFALFA AND SOME OTHER HAYS. *Colo. Agr. Expt. Sta. Bul.* 39, 34 pp.
- (254) ———
1907. COLORADO FODDERS. AN EXAMINATION INTO THEIR COMPOSITIONS AND COMPARATIVE VALUES. *Colo. Agr. Expt. Sta. Bul.* 124, 99 pp.
- (255) ———
1915. OCCURRENCE OF MANGANESE IN WHEAT. *Jour. Agr. Res.* 5: 349-355.
- (256) ———
1921. TITANIUM, BARIUM, STRONTIUM AND LITHIUM IN CERTAIN PLANTS. *Colo. Agr. Expt. Sta. Bul.* 267, 20 pp.
- (257) ———
1924. THE EFFECTS OF NITRATES ON THE COMPOSITION OF THE POTATO. *Colo. Agr. Expt. Sta. Bul.* 291, 32 pp.
- (258) HELLER, V. G., JONES, MAURINE, and PURSELL, LEE.
1935. IODINE CONTENT OF OKLAHOMA VEGETABLES. *Okla. Agr. Expt. Sta. Bul.* 229, 8 pp.
- (259) HENGL, FRANZ, and RECKENDORFER, PAUL.
1928. DER SCHWEFELGEHALT DES BODENS UND SEINE BEZIEHUNG ZUR PFLANZE. *Fortschr. der Landw.* 3: 598-600.
- (260) HENRY, MAX.
1912. OSTEO-MALACIA IN CATTLE IN NEW SOUTH WALES. *Agr. Gaz. N. S. Wales* 23: 885-888.
- (261) ———
1915. MORTALITY AMONGST CATTLE IN THE BEGA DISTRICT OF NEW SOUTH WALES. *Vet. Jour.* 71: 62-72.
- (262) ———
1925. THE INFLUENCE OF THE MINERAL CONSTITUENTS OF FOOD ON ANIMAL HEALTH. *Agr. Gaz. N. S. Wales* 36: 888-894.
- (263) HERCUS, C. E., AITKEN, H. A. A., THOMPSON, H. M. S., and COX, G. H.
1931. THE OCCURRENCE OF IODINE IN RELATION TO ENDEMIC GOITRE IN NEW ZEALAND AND ON IODINE METABOLISM. *Jour. Hyg. [London]* 31: 493-522.
- (264) ——— BENSON, W. NOEL, and CARTER, CHAS. L.
1925. ENDEMIC GOITRE IN NEW ZEALAND AND ITS RELATION TO THE SOIL-IODINE. *Jour. Hyg. [London]* 24: 321-402, illus.
- (265) ——— and ROBERTS, K. C.
1927. THE IODINE CONTENT OF FOODS, MANURES, AND ANIMAL PRODUCTS IN RELATION TO THE PROPHYLAXIS OF ENDEMIC GOITRE IN NEW ZEALAND. *Jour. Hyg. [London]* 26: 49-83, illus.
- (266) HESTER, JACKSON B.
1935. THE INFLUENCE OF SOIL ACIDITY AND SOIL TYPE UPON THE GROWTH AND COMPOSITION OF THE LIMA BEAN PLANT. *Amer. Soc. Hort. Sci. Proc. (1934)* 32: 600-603.

- (267) HESTER, JACKSON B.
1935. THE AMPHOTERIC NATURE OF THREE COASTAL PLAIN SOILS: II. IN RELATION TO THE LEACHING AND ABSORPTION OF SOIL CONSTITUENTS BY PLANTS. *Soil Sci.* 39: 247-254.
- (268) ——— and SHELTON, FLORENCE A.
1934. COMPARATIVE DATA FOR THREE COASTAL PLAIN SOILS FOR SOIL CHARACTERISTICS AND PLANT GROWTH. *Va. Truck Expt. Sta. Bul.* 84, pp. 1157-1193.
- (269) ——— and SHELTON, FLORENCE A.
1936. THE INFLUENCE OF CERTAIN REPLACEABLE BASES IN THE SOIL UPON THE ELEMENTAL COMPOSITION OF VEGETABLE CROPS. *Soil Sci.* 42: 335-340.
- (270) HILLS, J. L.
1893. THE PROPER TIME FOR HARVESTING FODDER CORN. *Vt. Agr. Expt. Sta. Ann. Rpt.* 6 (1892): 157-162.
- (271) ——— WHITE, B. O., and JONES, C. H.
1898. CHEMICAL COMPOSITION OF CORN AS AFFECTED BY FERTILIZATION. *Vt. Agr. Expt. Sta. Ann. Rpt.* 11: 146-188.
- (272) ——— WHITE, B. O., and JONES, C. H.
1898. CHEMICAL COMPOSITION OF POTATOES AS AFFECTED BY FERTILIZATION. *Vt. Agr. Expt. Sta. Ann. Rpt.* 11: 185-188.
- (273) HJORTH-HANSEN, SVERRER.
1931. ZUR RELATION ZWISCHEN BODENREAKTION UND CHEMISCHER ZUSAMMENSETZUNG DES HAFERS (*AVENA SATIVA*). *Biochem. Ztschr.* 235: 359-366.
- (274) HOLLEY, K. T., PICKETT, T. A., and BROWN, W. L.
1935. THE IODINE CONTENT OF SOME GEORGIA VEGETABLES AND WATER AS A FACTOR IN ITS VARIATION. *Ga. Expt. Sta. Bul.* 190, 11 pp.
- (275) HOLTZ, HENRY F.
1930. EFFECT OF CALCIUM AND PHOSPHORUS CONTENT OF VARIOUS SOIL SERIES IN WESTERN WASHINGTON UPON THE CALCIUM AND PHOSPHORUS COMPOSITION OF OATS, RED CLOVER AND WHITE CLOVER. *Wash. Agr. Expt. Sta. Bul.* 243, 45 pp.
- (276) HOPKINS, E. F., and GOURLEY, J. H.
1933. A STUDY OF THE ASH CONSTITUENTS OF APPLE FRUITS DURING THE GROWING SEASON. *Ohio Agr. Expt. Sta. Bul.* 519, 30 pp.
- (277) HOPKIRK, C. S. M., DAYUS, C. V., SIMPSON, B. W., and GRIMMETT, R. E. R.
1930. IODINE DEFICIENCY AND LIVESTOCK, A PRELIMINARY INVESTIGATION IN THE WANAKA AREA. (1) GOITRE IN LAMBS. *New Zeal. Jour. Agr.* 40: 226-232.
- (278) HOPPER, T. H., and NESBITT, L. L.
1930. THE CHEMICAL COMPOSITION OF SOME NORTH DAKOTA PASTURE AND HAY GRASSES. *N. Dak. Agr. Expt. Sta. Bul.* 236, 38 pp.
- (279) HOSKING, J. S., and GREAVES, G. A.
1935-36. A SOIL SURVEY OF AN AREA AT GINGIN, WESTERN AUSTRALIA. *Roy. Soc. West Austral. Jour.* 22: 71-112.
- (280) HUDIG, J., and LEHR, J. J.
1938. EINIGE BEMERKUNGEN ÜBER CHILESALPETER IN BEZIEHUNG ZUR BOFRAGE. *Bodenk. u. Pflanzenernähr.* 9-10: 552-579.
- (281) HUFFMAN, C. F., and TAYLOR, GEORGE E.
1926. DEPRAVED APPETITE IN DAIRY CATTLE . . . *Mich. Agr. Expt. Sta. Quart. Bul.* 8: 174-177, illus.
- (282) HURD-KARRER, ANNIE M.
1935. FACTORS AFFECTING THE ABSORPTION OF SELENIUM FROM SOILS BY PLANTS. *Jour. Agr. Res.* 50: 413-427, illus.
- (283) ———
1936. INHIBITION OF ARSENIC INJURY TO PLANTS BY PHOSPHORUS. *Wash. Acad. Sci. Jour.* 26: 180-181.
- (284) ———
1937. RUBIDIUM AND STRONTIUM TOXICITY TO PLANTS INHIBITED BY POTASSIUM AND CALCIUM RESPECTIVELY. *Wash. Acad. Sci. Jour.* 27: 351-353.
- (285) ——— and KENNEDY, MARY H.
1936. INHIBITING EFFECT OF SULFUR IN SELENIZED SOIL ON TOXICITY OF WHEAT TO RATS. *Jour. Agr. Res.* 52: 933-942, illus.
- (286) IBAÑEZ, R. P.
1937. BROMINE IN FLOURS AND WHEATS. *Abstract in Analyst* 62: 394.

- (287) IBELE, J.
1916. VORLÄUFIGER BERICHT ÜBER DIE CHEMISCHE ZUSAMMENSETZUNG DES HEUES IN BEZUG ZUR LECKSUCHT UND KNOCHENBRÜCHIGKEIT DES RINDES. Landw. Jahrb. f. Bayern 6: 183-204.
- (288) ———
1916. DER MINERALSTOFFGEHALT DES FUTTERS UND DIE LECKSUCHT DES RINDES. München. Tierärztl. Wehnschr. 67: 829-836, 849-856.
- (289) ———
1918. LECKSUCHT, PSEUDOLECKSUCHT UND "SUCHT ZU LECKEN" BEIM RINDE UND DEREN BEKÄMPFUNG. Ver. z. Förd. der Moorkult. im Deut. Reiche, Mitt. 36: 14-17.
- (290) ILJIN, W. S.
1938. CALCIUM CONTENT OF DIFFERENT PLANTS AND ITS INFLUENCE ON PRODUCTION OF ORGANIC ACIDS. Bul. Assoc. Russ. pour les Recherches Scientifiques à Prague 7 (41): 43-76.
- (291) INGLE, HERBERT.
1907. THE COMPOSITION OF SOME TRANSVAAL FODDERS. Transvaal Agr. Jour. 6: 51-65.
- (292) ITALLIE, T. B. VAN.
1934. DIE CHEMISCHE SAMENSTELLING VAN EEN AANTAL AFZONDERLIJKE GRASSOORTEN IN VERSCHILLENDE GROEISTADIA. Verslag. Landbouwk. Onderzoek. 40A: 639-693.
- (293) ———
1937. DE INVLOED VAN VERSCHILLENDE FACTOREN (MAAITIJD, BOTANISCHE SAMENSTELLING, BEMESTING EN GRONDSOORT) OP DE CHEMISCHE SAMENSTELLING VAN GRAS. Landbouwk. Tijdschr. [Wageningen] 49: 155-170.
- (294) ———
1938. DE NATRIM—EN KALIUMOPNAME BIJ VERSCHILLENDE BIETENRASSEN. VERSLAG. Landbouwk. Onderzoek. Rijkslandbouw proefsta. Groningen 1937, 43 (15) A: 721-766.
- (295) JACKS, G. V., and SCHERBATOFF, H.
1934. SOIL DEFICIENCIES AND PLANT DISEASES. Imp. Bur. Soil Sci. Tech. Commun, 31, 66 pp.
- (296) JACKSON, C. T.
1859. CHEMICAL ANALYSES OF TOBACCO SOILS—ANALYSES OF THE ASH OF TOBACCO PLANTS. U. S. Pat. Off. Rpt. (Agr.) for 1858, pp. 290-307.
- (297) JADIN, F., and ASTRUC, A.
1912. QUELQUES DÉTERMINATIONS QUANTITATIVES DU MANGANÈSE DANS LE RÈGNE VÉGÉTAL. [Paris] Acad. des Sci. Compt. Rend. 155: 406-408.
- (298) ——— and ASTRUC, A.
1912. LA PRÉSENCE DE L'ARSENIC DANS LE RÈGNE VÉGÉTAL. Jour. de Pharm. et de Chim. (7) 6: 529-535.
- (299) ——— and ASTRUC, A.
1913. LA RÉPARTITION DU MANGANÈSE DANS LE RÉGNE VÉGÉTAL. Jour. de Pharm. et de Chim. (7) 7: 155-161.
- (300) ——— and ASTRUC, A.
1914. L'ARSENIC ET LE MANGANÈSE DANS QUELQUES PRODUITS VÉGÉTAUX SERVANT D'ALIMENTS AUX ANIMAUX. [Paris] Acad. des Sci. Compt. Rend. 159: 268-272
- (301) JAVILLIER, MAURICE.
1908. LE ZINC CHEZ LES PLANTES. Inst. Pasteur [Paris] Ann. 22: 720-727.
- (302) JOACHIM, A. W. R.
1938. THE ANALYSIS OF CEYLON FOODSTUFFS I-V. Trop. Agr. [Ceylon] 90: 3-29.
- (303) JOHNSON, J. C., and DeLONG, W. A.
1937. BORON CONTENT OF APPLES AT DIFFERENT STAGES OF DEVELOPMENT. Plant Physiol. 12: 219-220.
- (304) JOHNSTONE-WALLACE, D. B.
1937. THE INFLUENCE OF WILD WHITE CLOVER ON THE SEASONAL PRODUCTION AND CHEMICAL COMPOSITION OF PASTURE HERBAGE, AND UPON SOIL TEMPERATURE, SOIL MOISTURE, AND EROSION CONTROL. 4th Internatl. Grassland Cong., Gt. Brit., 1937, Rpt., pp. 188-196.

- (305) JONES, D. BREESE.
1936. MINERAL ELEMENTS IN NUTRITION. Amer. Pub. Health Assoc. Yearbook 1936-37: 63-71.
- (306) JONES, J. S., and BULLIS, D. E.
1921. MANGANESE IN COMMONLY GROWN LEGUMES. Jour. Indus. and Engin. Chem. 13: 524-526.
- ✓(307) ——— and HATCH, M. B.
1937. THE SIGNIFICANCE OF INORGANIC SPRAY RESIDUE ACCUMULATIONS IN ORCHARD SOILS. Soil Sci. 44: 37-62.
- (308) JOSLAND, S. W.
1937. THE EFFECT OF FEEDING EXCESSES OF COBALT TO HEALTHY SHEEP. New Zeal. Jour. Sci. and Technol. 19: 31-37.
- (309) KALKUS, J. W.
1920. A STUDY OF GOITRE AND ASSOCIATED CONDITIONS IN DOMESTIC ANIMALS. Wash. Agr. Expt. Sta. Bul. 156, 48 pp.
- (310) KARMRODT, C.
1868. Jahresber. f. Agr. Chem. 12: 545.
- (311) KEATON, CLARK M.
1937. THE INFLUENCE OF LEAD COMPOUNDS ON THE GROWTH OF BARLEY. Soil Sci. 43: 401-412.
- (312) KELLEY, W. P., and BROWN, S. M.
1928. BORON IN THE SOILS AND IRRIGATION WATERS OF SOUTHERN CALIFORNIA AND ITS RELATION TO CITRUS AND WALNUT CULTURE. Hilgardia 3: [445]-458.
- (313) ——— and THOMPSON, ALICE R.
1910. A STUDY OF THE COMPOSITION OF THE RICE PLANT. Hawaii Agr. Expt. Sta. Bul. 21, 51 pp.
- (314) KELLNER, O., KOHLER, A., and BARNSTEIN, F.
1894. UNTERSUCHEN VERSCHIEDENER RAUHFUTTERARTEN AUS WIRTSCHAFTEN IN DENEN KNOCHENBRÜCHIGKEIT AUFTRITT. Sächs. Landw. Ztschr. (1895). 42: 167-171. Abstract in Biedermann's Zentbl. f. Agr. Chem. 24: 441-443.
- (315) KIDSON, E. B.
1937. COBALT STATUS OF NEW ZEALAND SOILS. New Zeal. Jour. Sci. and Technol. 18: 694-707.
- (316) ———
1938. SOME FACTORS INFLUENCING THE COBALT CONTENT OF SOILS. Soc. Chem. Indus. Jour. 57: 95T-96T.
- (317) KLIMMER and SCHMIDT.
1906. BEITRAG ZUR AETIOLOGIE DER HALISTERESIS OSSIUM NEBST THERAPEUTISCHEN BEMERKUNGEN. Monatsh. f. Prakt. Tierheilk. 17: 481-517.
- (318) KÖNIG, J., and KARST, H.
1923. DER EINFLUSS DES BODENS UND DER DÜNGUNG AUF DIE ZUSAMMENSETZUNG DER PFLANZEN. EIN BEITRAG ZUR GESCHICHTE DER LECKSUCHT. Landw. Vers. Sta. 100: 269-413.
- (319) KOHLER, G. O., ELVEHJEM, C. A., and HART, E. B.
1936. MODIFICATIONS OF THE BIPYRIDINE METHOD FOR AVAILABLE IRON. Jour. Biol. Chem. 113: 49-53.
- (320) KOSSOWITSCH, P.
1909. THE ASSIMILATION OF PHOSPHORUS BY PLANTS GROWN IN DIFFERENT SOILS. Zhur. Opytn. Agron. 10: 782-842.
- (321) KRÜGEL, C., DFEHSPRING, E., and LOTTHAMMER, R.
1938. WIRD DER PHOSPHORSAUREGHEALT VON WILDFUTTERPFLANZEN DURCH SUPERPHOSPHAT-DÜNGUNG ERHÖHT? Superphosphat 14: 103-105.
- (322) KUNZE, ERICH.
1914. ÜBER DEN MINERALSTOFFGEHALT EINIGER LANDWIRTSCHAFTLICHER FUTTERMITTEL (HAFER UND GRUMMET) IM ZUSAMMENHANGE MIT DEM WETTER UND DER MINERALSTOFFZUFUHR BEI DER ERNÄHRUNG DER RINDER. Landw. Inst. Leipzig Mitt. 12: 63-96.
- (323) KÜSTER, W., and UMBRECHT, J.
1928. ÜBER DEN GEHALT DER LINSEN UND ERBSEN AN NATRIUM UND KALIUM. Hoppe-Seylers Ztschr. f. Physiol. Chem. 179: 139-148.

- (324) LAGATU, H.
1921. SUR LE RÔLE RESPECTIF DES TROIS BASES: POTASSE, CHAUX, MAGNÉSIE, DANS LES PLANTES CULTIVÉES. [Paris] Acad. des Sci. Compt. Rend. 172: 129-131.
- (325) ——— and MAUME, L.
1924. ÉVOLUTION RÉMARQUABLEMENT RÉGULIÈRE DE CERTAINS RAPPORTS PHYSIOLOGIQUES (CHAUX, MAGNÉSIE, POTASSE) DANS LES FEUILLES DE LA VIGNE BIEN ALIMENTÉE. [Paris] Acad. des Sci. Compt. Rend. 179: 782-785.
- (326) ——— and MAUME, L.
1925. RÉLATION LINÉAIRE ENTRE LES QUANTITÉS SUCCESSIVES D'ACIDE PHOSPHORIQUE ET D'AZOTE CONTENUES DANS LA FEUILLE DE LA VIGNE BIEN ALIMENTÉE. [Paris] Acad. des Sci. Compt. Rend. 180: 1179-1181.
- (327) ——— and MAUME, L.
1927. MÉTHODE DE CONTRÔLE CHIMIQUE DU MODE D'ALIMENTATION DE LA VIGNE. Acad. d'Agr. de France Compt. Rend. 13: 437-455.
- (328) ——— and MAUME, L.
1931. VARIATION DES RAPPORTS PHYSIOLOGIQUES ENTRE LES CONSTITUANTS MINÉRAUX D'UNE ESPÈCE VÉGÉTALE. [Paris] Acad. des Sci. Compt. Rend. 193: 103-116.
- (329) ——— and MAUME, L.
1934. EXAMEN CRITIQUE DU DIAGNOSTIC FOLIAIRE. Acad. d'Agr. de France Compt. Rend. 20: 246-256.
- (330) ——— and MAUME, L.
1938. SUR LA SENSIBILITÉ DU DIAGNOSTIC FOLIAIRE POUR DÉCÉLER UNE ABSORPTION D'ENGRAIS PAR LA VIGNE. Acad. d'Agr. de France Compt. Rend. 24: 615-624.
- (331) LAKIN, H. W., WILLIAMS, K. T., and BYERS, H. G.
1938. "NON-TOXIC" SELENIFEROUS SOILS. Indus. and Engin. Chem. 30: 599-600.
- (332) LAWES, J. B., and GILBERT, J. H.
1884. ON THE COMPOSITION OF THE ASH OF WHEAT GRAIN, AND WHEAT STRAW, GROWN AT ROTHAMSTED, IN DIFFERENT SEASONS, AND BY DIFFERENT MANURES. [London] Chem. Soc. Jour. 45: 305-407.
- (333) ——— and GILBERT, J. H.
1900. AGRICULTURAL, BOTANICAL AND CHEMICAL RESULTS OF EXPERIMENTS ON THE MIXED HERBAGE OF PERMANENT GRASS-LAND. III. THE CHEMICAL RESULTS. [London] Roy. Soc. Phil. Trans. (B) 192: 139-210.
- (334) LECHARTIER, G.
1892. DES VARIATIONS DE COMPOSITION DU TOPINAMBOUR. Ann. Agron. [Paris] 18: 68-91.
- (335) ———
1903. ÉTUDE SUR LE SOJA HISPIDA. CULTURE ET COMPOSITION. Ann. de la Sci. Agron. Franç. et Étrang. (2) 8: 380-396.
- (336) LECLERC, J. A.
1907. EFFECT OF CLIMATIC CONDITIONS ON THE COMPOSITION OF DURUM WHEAT. U. S. Dept. Agr. Yearbook 1906: 199-212.
- (337) ——— and BREAZEALE, J. F.
1910. PLANT FOOD REMOVED FROM GROWING PLANTS BY RAIN OR DEW. 7th Internat. Cong. Appl. Chem. 1909, Sec. 7 (Agr.), pp. 55-78. London.
- (338) ——— and LEAVITT, SHERMAN.
1910. THE INFLUENCE OF ENVIRONMENT ON THE COMPOSITION OF WHEAT. 7th Internat. Cong. Appl. Chem. 1909, Sect. 7 (Agr.), pp. 136-147. London.
- (339) ——— and YODER, P. A.
1912. TRI-LOCAL SOIL EXCHANGE EXPERIMENTS WITH WHEAT. 8th Internat. Cong. Appl. Chem. 26: 137-150.
- (340) LEEPER, G. W.
1934. RELATIONSHIP OF SOILS TO MANGANESE DEFICIENCY OF PLANTS. Nature [London] 134: 972-973.
- (341) ———
1935. MANGANESE DEFICIENCY OF CEREALS: PLOT EXPERIMENTS AND A NEW HYPOTHESIS. Roy. Soc. Victoria Proc., 47: 225-261.

- (342) LEEUWEN, MARIA VAN.
1930. RÉCHERCHES SUR LA TENEUR EN CUIVRE ET EN MANGANÈSE DE QUELQUES TISSUS VÉGÉTAUX. *Ann. Physiol., Physicochim., Biol.* 6: 178-181.
- (343) LEHMANN, K. B.
1896. HYGIENISCHE STUDIEN ÜBER KUPFER. IV. DER KUPFERGEGHALT VON PFLANZEN UND TIEREN IN KUPFERREICHEN GEGENDEN. *Arch. f. Hyg.* 27: 1-17.
- (344) LEMMERMANN, O., EINECKE, A., and FISCHER, H.
1911. UNTERSUCHUNGEN ÜBER DIE WIRKUNG EINES VERSCHIEDENEN VERHÄLTNISSSES VON KALK UND MAGNESIA IN EINIGEN BÖDEN AUF HÖHERE PFLANZEN UND MIKROORGANISMEN. *Landw. Jahrb.* 40: 173-244.
- (345) LEPAPE, ADOLPHE, and TRANNOY, RENÉ.
1934. FIXATION, PAR LES PLANTES, DU RADIUM MIS À LEUR DISPOSITION DANS LE SOL. [Paris] *Acad. des Sci. Compt. Rend.* 199: 803-805.
- (346) LESKIEN, J.
1937. STAND DER KENNTNISSE ÜBER DIE LECKSUCHT DES KINDES. *For- schungsdienst.* 4: 542-548.
- (347) LEVINE, HAROLD, CULP, F. B., and ANDERSON, C. B.
1932. THE VALUE OF SOME VEGETABLES IN NUTRITIONAL ANEMIA. *Jour. Nutr.* 5: 295-306.
- (348) LEWITE, MAXIM.
1907. ZUR FRAGE DES ZUSAMMENHANGES ZWISCHEN WETTER, MINERAL- STOFFGEGHALT DER FUTTERPFLANZEN UND KNOCHENBRÜCHIGKEIT DES RINDES. *Landw. Inst. Leipzig Mitt.* 8: 69-112.
- (349) LIBERI, G., CUSMANO, A., MARSIGLIA, T., and ZAY, C.
1916. SULLA PRESENZA DEL RAME NEI FRUTTI E NELLE CONSERVE DI POMODORO. *Roma R. Staz. Chim. Agr. Sper. Ann.* (2) 8: 163-303.
- (350) LICHTENSTEIN, HENRY.
1930. TRAVELS IN SOUTH AFRICA. Transl. by Anne Plumptre. 2v, Cape Town. 1930.
- (351) LICHTIN, AARON.
1923. THE IRON CONTENT OF LETTUCE. *Amer. Jour. Pharm.* 95: 154-159.
- (352) LIEBSCHER, KRETSCHMER, VON SEELHORST, and WILMS, J.
1898. ERMITTELUNG DER DÜNGERBEDÜRFTIGKEIT DES BODENS AUS DER ZUSAMMENSETZUNG DER ERNTELOCKENSUBSTANZ. *Jour. f. Landw.* 46: 367-412.
- (353) LIGON, W. S.
1935. THE SOLUBILITY OF APPLIED NUTRIENTS IN MUCK SOILS AND THE COMPOSITION AND QUALITY OF CERTAIN MUCK CROPS AS IN- FLUENCED BY SOIL REACTION CHANGES AND MOISTURE CONDI- TIONS. *Mich. Agr. Expt. Sta. Tech. Bul.* 147, 51 pp.
- (354) LINCOLN, R.
1937. THE MINERAL CONTENT OF MAURITIUS PASTURES. *Mauritius Dept. Agr. Bul.* 26 (Sci. Ser.), 14 pp.
- (355) LINDOW, C. W., ELVEHJEM, C. A., and PETERSON, W. H.
1929. THE COPPER CONTENT OF PLANT AND ANIMAL FOODS. *Jour. Biol. Chem.* 82: 465-471.
- (356) ——— and PETERSON, W. H.
1927. THE MANGANESE CONTENT OF PLANT AND ANIMAL MATERIALS. *Jour. Biol. Chem.* 75: 169-175.
- (357) LÖHNIS, MARIE P.
1936. WAT VEROORZAAKT KWAVE HARTEN IN ERWTEN. *Tijdschr. over Plantenziekten* 42: 159-167.
- (358) LOTT, WREAL L.
1939. THE RELATION OF HYDROGEN-ION CONCENTRATION TO THE AVAIL- ABILITY OF ZINC IN SOIL. *Soil Sci. Soc. Amer. Proc.* (1938) 3: 115-121.
- (359) MCBRYDE, J. B.
1890. THE COWPEA AS A FORAGE CROP. *S. C. Agr. Expt. Sta. Ann. Rpt.* (1889) 2: 169-180.
- (360) MCCANCE, R. A.
1936. MEDICAL PROBLEMS IN MINERAL METABOLISM. LEGACIES OF EVO- LUTION. *Lancet* 230: 643-650.

- (361) McCARRISON, R.
1927. GOITRE. *Indian Jour. Med. Res.* 14: 655-658.
- (362) ——— NEWCOMB, C., VISWANATH, B., and NORRIS, R. V.
1927. THE RELATION OF ENDEMIC GOITER TO THE IODINE CONTENT OF SOIL AND DRINKING WATER. *Indian Jour. Med. Res.* 15: 207-246.
- (363) McCLENDON, J. F.
1924. INVERSE RELATION BETWEEN IODINE IN FOOD AND DRINK AND GOITER, SIMPLE AND EXOPHTHALMIC. *Amer. Med. Assoc. Jour.* 82: 1668-1672.
- (364) ——— and HOLDRIDGE, C. E.
1935. IODINE IN CABBAGE. *Biochem. Jour.* 29: 272-274.
- (365) ——— and REMINGTON, ROE E.
1929. THE DETERMINATION OF TRACES OF IODINE. II. IODINE IN VEGETABLES. *Amer. Chem. Soc. Jour.* 51: 394-399, illus.
- (366) McCOOL, M. M., and WELDON, M. D.
1928. THE EFFECT OF SOIL TYPE AND FERTILIZATION ON THE COMPOSITION OF THE EXPRESSED SAP OF PLANTS. *Amer. Soc. Agron. Jour.* 20: 778-792.
- (367) McGOWAN, J. P., and SMITH W. G.
1922. ON PINING, VINQUISH OR DAISING IN SHEEP. *Scot. Jour. Agr.* 5: 274-286.
- (368) MACH, F., and HERRMANN, R.
1934. NÄHRSTOFF- UND ASCHENANALYSEN VON WIRTSCHAFTSEIGENEN FUTTERMITTELN, AUSGEFÜHRT VON MITGLIEDERN DES VERBANDES DEUTSCHER LANDWIRTSCHAFTLICHER VERSUCHSSTATIONEN. *Landw. Vers. Sta.* 119: 1-173.
- (369) McHARGUE, J. S.
1913. THE OCCURRENCE OF BARIUM IN TOBACCO AND OTHER PLANTS. *Amer. Chem. Soc. Jour.* 35: 826-834.
- (370) ———
1914. THE OCCURRENCE AND SIGNIFICANCE OF MANGANESE IN THE SEED COAT OF VARIOUS SEEDS. *Amer. Chem. Soc. Jour.* 36: 2532-2536.
- (371) ———
1923. IRON AND MANGANESE CONTENT OF CERTAIN SPECIES OF SEEDS. *Jour. Agr. Res.* 23: 395-399.
- (372) ———
1925. THE OCCURRENCE OF COPPER, MANGANESE, ZINC, NICKEL AND COBALT IN SOILS, PLANTS, AND ANIMALS, AND THEIR POSSIBLE FUNCTION AS VITAL FACTORS. *Jour. Agr. Res.* 30: 193-196.
- (373) ———
1925. THE SIGNIFICANCE OF THE OCCURRENCE OF COPPER, MANGANESE AND ZINC IN FORAGE CROPS AND FOODS. *Amer. Soc. Agron. Jour.* 17: 368-372.
- (374) ———
1926. MINERAL CONSTITUENTS OF THE COTTON PLANT. *Amer. Soc. Agron. Jour.* 18: 1076-1083.
- (375) ———
1927. SIGNIFICANCE OF THE OCCURRENCE OF MANGANESE, COPPER, ZINC, NICKEL AND COBALT IN KENTUCKY BLUEGRASS. *Indus. and Engin. Chem.* 19: 274-276.
- (376) ———
1931. EFFECT OF MANGANESE DEFICIENT LETTUCE ON THE GROWTH OF RATS. *Ky. Agr. Expt. Sta. Ann. Rpt.* 44: 24-25.
- (377) ———
1933. IODINE CONTENT OF CORN PRODUCED IN NINETEEN EASTERN KENTUCKY COUNTIES. *Ky. Agr. Expt. Sta. Ann. Rpt.* 46: 44-45.
- (378) ——— and CALFEE, R. K.
1937. THE NECESSITY OF MINOR ELEMENTS FOR THE GROWTH OF TOMATOES IN A POOR SOIL. *Amer. Soc. Agron. Jour.* 29: 385-391.
- (379) ——— and ROY, W. R.
1931. MINERAL, NITROGEN AND FAT CONTENT OF SOME VARIETIES OF MATURE BEAN SEED AND OF STRING BEANS. *Jour. Nutr.* 3: 479-481.

- (380) McHARGUE, J. S., and ROY, W. R.
1931. THE EFFECT OF A DIET OF POLISHED RICE ON THE MINERAL CONTENT OF THE CARCASSES OF PIGEONS. *Amer. Jour. Physiol.* 99: 221-226.
- (381) ——— ROY, W. R., and PELPHREY, J. G.
1932. THE IRON, MANGANESE, COPPER, ZINC AND IODINE CONTENT OF SOME IMPORTANT FORAGE CROPS. *Amer. Soc. Agron. Jour.* 24: 562-565.
- (382) ——— and SHEDD, O. M.
1930. THE EFFECT OF MANGANESE, COPPER, ZINC, BORON, AND ARSENIC ON THE GROWTH OF OATS. *Amer. Soc. Agron. Jour.* 22: 739-746.
- (383) ——— YOUNG, D. W., and CALFEE, R. K.
1935. THE EFFECT OF CERTAIN FERTILIZER MATERIALS ON THE IODINE CONTENT OF IMPORTANT FOODS. *Amer. Soc. Agron. Jour.* 27: 559-565.
- (384) McINTOSH, R. A.
1934. REPORT OF THE COMMITTEE MAKING A SURVEY OF MINERAL DEFICIENCY DISEASE IN CATTLE FOR THE YEAR 1933. *Ontario Vet. Col. Rpt.* 1934: 13-24.
- (385) MACK, WARREN B., and BRASHER, EUGENE P.
1936. THE INFLUENCE OF COMMERCIAL FERTILIZERS, POTASSIUM IODIDE AND SOIL ACIDITY ON THE IODINE CONTENT OF CERTAIN VEGETABLES. *Jour. Agr. Res.* 53: 789-800.
- (386) McKENNA, C. T.
1929. SOME DISEASES OF STOCK IN SOUTH AUSTRALIA CONSIDERED TO BE DUE TO MINERAL DEFICIENCY. *So. Austral. Dept. Agr. Jour.* 32: 997-1002.
- (387) McLARTY, H. R., WILCOX, J. C., and WOODBRIDGE, C. G.
1937. A YELLOWING OF ALFALFA DUE TO BORON DEFICIENCY. *Sci. Agr.* 17: 515-517.
- (388) McMURTREY, J. E., Jr., and ROBINSON, W. O.
1938. NEGLECTED SOIL CONSTITUENTS THAT AFFECT PLANT AND ANIMAL DEVELOPMENT. *Soils and Men (U. S. Dept. Agr. Yearbook, 1938):* 807-829.
- (389) MADER, E. O., and MADER, M. T.
1937. EFFECT OF BORDEAUX MIXTURE ON THREE VARIETIES OF POTATOES WITH RESPECT TO YIELDS, COMPOSITION OF TUBERS, AND CONTROL OF SCAB. *Phytopathology* 27: 1032-1045.
- (390) ——— and MADER, M. T.
1937. THE COMPOSITION OF TUBERS OF SPRAYED AND UNSPRAYED POTATO PLANTS IN RELATION TO COOKING QUALITY. *Amer. Potato Jour.* 14: 56-59.
- (391) MAQUENNE, L., and DEMOUSSY, E.
1920. LE CUIVRE DANS LA TERRE ET LES PLANTES. *Soc. Chim. de France Bul. Doc.* 27: 266-278.
- (392) ——— and DEMOUSSY, E.
1920. SUR LA DISTRIBUTION ET LA MIGRATION DU CUIVRE DANS LES TISSUS DES PLANTES VERTES. [Paris] *Acad. des Sci. Compt. Rend.* 170: 87-93.
- (393) MARAÑON, JOAQUIN.
1935. NUTRITIVE MINERAL VALUE OF PHILIPPINE FOOD PLANTS [CALCIUM, PHOSPHORUS, AND IRON CONTENTS]. *Philippine Jour. Sci.* 58: 317-358.
- (394) ——— and LASERNA, G.
1937. CHEMICAL ANALYSIS OF SOME PHILIPPINE FORAGE PLANTS. *Philippine Jour. Sci.* 64: 189-197.
- (395) MARINE, DAVID.
1933. THE IMPORTANCE OF RELATIVE IODINE DEFICIENCIES IN CERTAIN FORMS OF GOITER. *Amer. Dietet. Assoc. Jour.* 9: 1-5.
- (396) MARSH, C. DWIGHT, ALSBERG, C. L., and BLACK, O. F.
1912. THE RELATION OF BARIUM TO THE LOCO-WEED DISEASE. *U. S. Dept. Agr., Bur. Plant Indus. Bul.* 246, 67 pp.
- (397) MASCHHAUPT, J. G.
1918. DE INVLOED VAN GRONDSOORT EN BEMESTING OP HET GEHALTE ONZER CULTURGEWASSEN AAN STIKSTOF EN ASCHBESTANDDEELLEN. [Netherlands] *Verslag. van Landbouwk. Onderzoek.* 22: 25-114, 1918; 23: 40-56, 1919; 25: 115-129, 1921; 27: 114-124, 1922.

- (398) MATHER, THOMAS H.
1929. THE EFFECT OF FERTILIZERS UPON THE FORMS OF PHOSPHORUS AND THE AMOUNTS OF PHOSPHORUS, NITROGEN, AND SILICA IN HAYS. *Sci. Agr.* 10: 35-63, illus.
- (399) MAUME, L., and BOUAT, A.
1937. INFLUENCES DE LA VARIÉTÉ ET DU MILIEU SUR L'ABSORPTION DU SOUFRE PAR LE BLÉ. *Acad. d'Agr. de France Compt. Rend.* 23: 426-430.
- (400) ——— and MONTEIL, L.
1938. DÉFICIENCE MINÉRALE DES FOURRAGES EN COMBRAILLE BOURBONNAISE. *Acad. d'Agr. de France Compt. Rend.* 24: 45-52.
- (401) MAXWELL, WALTER.
1900. SUGAR CANE. *Hawaii. Sugar Planters' Assoc. Expt. Sta. Rpt.* 1900, 51 pp.
- (402) MAYNARD, L. A.
1929. THE ROLE OF PASTURE IN THE MINERAL NUTRITION OF FARM ANIMALS. *Amer. Soc. Agron. Jour.* 21: 700-708.
- (403) MAYNARD, LEONARD A.
1937. ANIMAL NUTRITION. 483 pp., illus., New York and London.
- (404) MAYRHOFER, A., SCHNEIDER, C., and WASITZKY, A.
1932. BIOCHEMISCHE STUDIEN ÜBER DAS VORKOMMEN KLEINER MENGEN VON JOD UND FLUOR IN ORGANISMUS. II. *Biochem. Ztschr.* 251: 70-86.
- (405) MAZÉ, P.
1936. THE ROLE OF SPECIAL ELEMENTS (BORON, COPPER, ZINC, MANGANESE, ETC.) IN PLANT NUTRITION. *Ann. Rev. Biochem.* 5: 525-538.
- (406) MEINCK, F.
1927. BEITRAG ZUM JODGEHALT DES BODENS, DES WASSERS UND EINIGER NAHRUNGSMITTEL IN KROPPBEHAFTETEN GEGENDEN. *Kleine Mitt. Mitglieder Ver. Wasser-, Boden- u. Luft Hyg.* 3:10-11.
- (407) MEUNIER, PAUL.
1935. NOUVELLE MÉTHODE D'IDENTIFICATION ET DE DOSAGE DE TRÈS PETITES QUANTITÉS D'ALUMINIUM DANS LES VÉGÉTAUX. *Soc. de Chim. Biol. Bul.* 17: 548-560.
- (408) MILLER, CAREY D., BAZORE, KATHERINE, and ROBBINS, RUTH C.
1937. SOME FRUITS OF HAWAII. *Hawaii Agr. Expt. Sta. Bul.* 77, ed. 2, 133 pp.
- (409) ——— and ROBBINS, RUTH C.
1934. COMPOSITION AND VITAMIN STUDIES OF GREEN SOYBEANS. *Hawaii Agr. Expt. Sta. Rpt.* 1933, 24-25.
- (410) MILLER, EDWIN C.
1938. PLANT PHYSIOLOGY. Ed. 2, 1201 pp., illus. New York.
- (411) MILLER, LILA, and MITCHELL, HELEN S.
1931. CORRELATION OF COPPER AND MANGANESE CONTENT OF PLANTS AND MINERAL ADDITIONS TO THE SOIL. *Amer. Dietet. Assoc. Jour.* 7: 252-7.
- (412) MITCHELL, H. H., and McCLURE, F. J.
1937. MINERAL NUTRITION OF FARM ANIMALS. *Natl. Res. Council Bul.* 99, 135 pp.
- (413) MITCHELL, J. H., and MATTISON, W. T.
1933. A STUDY OF THE RELATIONSHIP BETWEEN THE MINERAL CONTENT OF THE SOIL AND THE PLANTS GROWN ON THE SOIL. *S. C. Agr. Expt. Sta. Ann. Rpt.* 46: 51-52.
- (414) ——— WARNER, J. D., and MORROW, K. S.
1928. THE MINERAL CONTENT OF FEEDS, SOILS AND WATERS OF SOUTH CAROLINA. *S. C. Agr. Expt. Sta. Bul.* 252, 32 pp.
- (415) MOORE, C. C.
1903. PRELIMINARY CROP AND SOIL DATA FOR THE COOPERATIVE STUDY OF AVAILABLE PLANT FOOD. *U. S. Dept. Agr., Bur. Chem. Cir.* 11, 9 pp.
- (416) MORSE, FRED. W.
1928. THE IODINE CONTENT OF CAPE COD CRANBERRIES. *Jour. Biol. Chem.* 79: 409-411.
- (417) ———
1929. THE MINERAL CONSTITUENTS OF CRANBERRIES. *Jour. Biol. Chem.* 81: 77-79.

- (418) MORTIMER, G. B., and AHLGREN, H. L.
1936. INFLUENCE OF FERTILIZATION, IRRIGATION AND STAGE AND HEIGHT OF CUTTING ON YIELD AND COMPOSITION OF KENTUCKY BLUEGRASS (*POA PRATENSIS* L.). *Amer. Soc. Agron. Jour.* 28: 515-533.
- (419) MOXON, ALVIN L.
1937. ALKALI DISEASE OR SELENIUM POISONING. *S. Dak. Agr. Expt. Sta. Bul.* 311, 87 pp.
- (420) MÜLLER, H.
1873. ÜBER DIE RESULTATE DER BOTANISCHEN UNTERSUCHUNG DER BOKER HAIDE BEHUFES ERMITTELUNG DER URSACHE DE KNOCHENBRÜCHIGKEIT. *Landw. Jahrb.* 2: 1-107.
- (421) MUNGER, STELLA, and PETERSON, W. H.
1928. THE MANGANESE CONTENT OF RAW AND COOKED VEGETABLES. *Jour. Home Econ.* 20: 194-200.
- (422) MURPHY, H. F.
1930. EFFECT OF FERTILIZERS ON THE YIELD AND COMPOSITION OF WHEAT. *Amer. Soc. Agron. Jour.* 22: 765-770.
- (423) ———
1936. THE NITROGEN, PHOSPHORUS AND CALCIUM CONTENT OF THE COTTON PLANT AT PRE-BLOOMING TO EARLY BOLL STAGES OF GROWTH. *Amer. Soc. Agron. Jour.* 28: 52-57.
- (424) MYERS, C. N., and VOEGTLIN, CARL.
1914. SOLUBLE ALUMINUM COMPOUNDS. THEIR OCCURRENCE IN CERTAIN VEGETABLE PRODUCTS. [U. S.] *Pub. Health Rpts.* 29: 1625-1629.
- (425) MYERS, H. E., and METZGER, W. H.
1936. THE INFLUENCE OF SUPERPHOSPHATE AND LIGHT LIME APPLICATIONS ALONE AND IN COMBINATION ON THE COMPOSITION OF SWEET CLOVER. *Amer. Soc. Agron. Jour.* 28: 976-984.
- (426) NAFTEL, JAMES A.
1937. SOIL LIMING INVESTIGATIONS: IV. THE INFLUENCE OF LIME ON YIELDS AND ON THE CHEMICAL COMPOSITION OF PLANTS. *Amer. Soc. Agron. Jour.* 29: 537-547.
- (427) NEAL, W. M., and AHMANN, C. F.
1937. THE ESSENTIALITY OF COBALT IN BOVINE NUTRITION. (Abstract.) *Jour. Dairy Sci.* 20: 406-407.
- (428) ———, and BECKER, R. B.
1933. THE COMPOSITION OF FEEDSTUFFS IN RELATION TO NUTRITIONAL ANEMIA IN CATTLE. *Jour. Agr. Res.* 47: 249-255.
- (429) ———, BECKER, R. B., and SHEALY, A. L.
1931. A NATURAL COPPER DEFICIENCY IN CATTLE RATIONS. *Science* 74: 418-419.
- (430) NEIDIG, RAY E., McDOLE, G. R., and MAGNUSON, H. P.
1923. EFFECT OF SULFUR, CALCIUM AND PHOSPHORUS ON THE YIELD AND COMPOSITION OF ALFALFA ON SIX TYPES OF IDAHO SOILS. *Soil Sci.* 16: 127-136.
- (431) ———, and SNYDER, ROBERT S.
1925. MINERAL COMPOSITION OF SUNFLOWERS GROWN FOR SILAGE. *Jour. Agr. Res.* 31: 1165-1171.
- (432) NELLER, J. R.
1925. THE INFLUENCE OF SULFUR AND GYPSUM UPON THE COMPOSITION AND YIELD OF LEGUMES. *Wash. Agr. Expt. Sta. Bul.* 190, 47 pp.
- (433) NĚMEC, B., BABIČKA, J., and OBORSKÝ, A.
1936. ÜBER DAS VORKOMMEN VON GOLD IN DEN SHACHTELHALMEN (*EQUISETUM PALUSTRE* UND *ARVENSE*). *Bul. Internatl. de l'Acad. des Sci. de Bohême.* (Contribution from the Pflanzenphysiologisches Institut der Karlsuniversität in Prag.) 7 pp.
- (434) NĚMEC, ANTONIN, and KÁŠ, VÁCLAV.
1923. STUDIEN ÜBER DIE PHYSIOLOGISCHE BEDEUTUNG DES TITANS IM PFLANZENORGANISMUS. *Biochem. Ztschr.* 140: 583.
- (435) NESSLER, J.
1873. UNTERSUCHUNGEN DER KNOCHEN VON KNOCHENBRÜCHIGEM RINDVIEH. *Landw. Vers. Sta.* 16: 187-192.
- (436) NEYEN, J. A.
1863. LA CACHEXIE OSSIFRAGE. *Rec. de Méd. Vét.* (4) 10: 340-351.

- (437) NICOLAISEN, W., and SEELBACH, W.
1938. UNTERSUCHUNGEN ÜBER DIE KUPFERSULFATDÜNGUNG GEGEN
URBARMACHUNGSKRANKHEIT UND LECKSUCHT. Forschungs-
dienst 5: 383-387.
- (438) NOLTE, O.
1923. UEBER DIE WIRKUNG STEIGENDER KALIGABEN AUF DEN ERTRAG
UND DEN PHOSPHORSAÜREGEHALT VON KULTURPFLANZEN BEI
GEGENWART VON BODEN- UND DUNGERPHOSPHORSÄURE NACH
VERSUCHEN IM JAHRE 1921. Ztschr. f. Pflanzenernähr. Düngung
u. Bodenk. (B) 2: 23-33.
- (439) NORMAN, MARION, and NORMAN, A. G.
1937. NUTRITION AND THE HALOGENS. Food Mfr. 12: 5-7.
- (440) NYGARD, IVER J.
1931. PHOSPHATE DEFICIENCY IN THE SOILS OF MONTANA. Mont. Agr.
Expt. Sta. Bul. 240, 32 pp.
- (441) OLSEN, CARSTEN.
1934. UEBER DIE MANGANAUFNAHME DER PFLANZEN. Biochem. Ztschr.
269: 329-348.
- (442) OPITZ, K., ROTHSACK, K., and MORGENROTH, E.
1937. UEBER DIE BEDEUTUNG DES NÄHRSTOFFVERHÄLTNISSSES BEI DER
DÜNGUNG MIT MINERALISCHEN DUNGEMITTELN. Bodenk. u.
Pflanzenernähr. 4: 210-232.
- (443) ORR, J. B., with the assistance of SCHERBATOFF, HELENE.
1929. MINERALS IN PASTURES & THEIR RELATION TO ANIMAL NUTRITION.
150 pp., illus. London.
- (444) ———
1929. THE RELATION OF THE CHEMICAL COMPOSITION OF PASTURE TO ITS
FEEDING VALUE. Highland and Agr. Soc. Scot. Trans. 41:
99-114.
- (445) ———
1931. IODINE SUPPLY AND THE INCIDENCE OF ENDEMIC GOITRE. [Gt.
Brit.] Med. Res. Council, Spec. Rpt. Ser. 154, 18 pp.
- (446) ——— and HOLM, ALEXANDER.
1931. THE MINERAL DEFICIENCIES OF PASTURES IN KENYA COLONY AND
THEIR EFFECTS ON GRAZING ANIMALS. Econ. Advisory Council
Com. on Min. Content of Nat. Pastures [Gt. Brit.] Rpt. 6: 14-55.
- (447) ——— KELLY, F. C., and STUART, G. L.
1928. THE EFFECT OF IODINE MANURING ON THE IODINE CONTENT OF
PLANTS. Jour. Agr. Sci. [England] 18: [159]-161.
- (448) OSTERTAG, R., and ZUNTZ, N.
1907. STUDIEN UEBER DIE LECKSUCHT DER RINDER. Ztschr. f. Infektions-
krank., Parasitäre Krank. u. Hyg. der Haustiere 2: 409-424.
- (449) PARKER, J. M.
1904. CREEPS, AN OSTEOMALACIAL DISEASE OF CATTLE. (Paper read
before the 41st meeting of the Amer. Vet. Med. Assoc., St. Louis,
Mo., Aug. 16-19. [Abstract in Expt. Sta. Rec. 16: 124.]
- (450) PARKER, M. M., HESTER, J. B., and CAROLUS, R. L.
1933. THE EFFECT OF SOIL CONDITIONS ON THE GROWTH AND COMPOSITION
OF CERTAIN VEGETABLE CROP PLANTS AS INFLUENCED BY SOIL
REACTIONS. Amer. Soc. Hort. Sci. Proc. (1933) 30: 452-457.
- (451) PATTERSON, J. B. E.
1937. COBALT, AND SHEEP DISEASES. Nature [London] 140: 363.
- (452) PETER, A. M.
1902. ANALYSES OF FORAGE PLANTS FROM THE PLOTS. Ky. Agr. Expt.
Sta. Bul. 98: 51-61.
- (453) ——— BEATTY, L. O., and AVERITT, S. D.
1902. MISCELLANEOUS CHEMICAL ANALYSES MADE IN 1902. Ky. Agr.
Expt. Sta. Ann. Rpt. 15: 283-335.
- (454) PETER, ROBERT.
1884. CHEMICAL EXAMINATION OF THE HEMP AND BUCKWHEAT PLANTS.
Ky. Geol. Survey Chem. Anal., 435-463.
- (455) PETERSON, W. H., and ELVEHJEM, C. A.
1928. THE IRON CONTENT OF PLANT AND ANIMAL FOODS. Jour. Biol.
Chem. 78: 215-223.
- (456) ——— ELVEHJEM, C. A., and JAMISON, LIDA A.
1925. VARIATIONS IN THE MINERAL CONTENT OF CABBAGE AND SAUER-
KRAUT. Soil Sci. 20: 451-457.

- (457) PETERSON, W. H., and HOPPERT, C. A.
1925. THE LOSS OF MINERAL AND OTHER CONSTITUENTS FROM VEGETABLES BY VARIOUS METHODS OF COOKING. *Jour. Home Econ.* 17: 265-280.
- (458) ——— and LINDOW, C. W.
1928. VARIATIONS IN THE MANGANESE CONTENT OF CERTAIN VEGETABLES. *Soil Sci.* 26: 149-153.
- (459) ——— PARMELE, H. B., and FRED, E. B.
1927. SOME OF THE FACTORS WHICH INFLUENCE THE COMPOSITION OF CABBAGE AND THEIR RELATION TO THE QUALITY OF SAUERKRAUT. *Soil Sci.* 24: 299-307.
- (460) ——— and PETERSON, CLARA B.
1926. THE WATER-SOLUBLE CONTENT OF CALCIUM AND PHOSPHORUS IN CABBAGE. *Jour. Agr. Res.* 33: 695-699.
- (461) ——— and SKINNER, J. T.
1931. DISTRIBUTION OF MANGANESE IN FOODS. *Jour. Nutr.* 4: 419-426.
- (462) PETIT, P.
1892. DISTRIBUTION ET ÉTAT DU FER DANS L'ORGE. [Paris] *Acad. des Sci. Compt. Rend.* 115: 246-248.
- (463) PFEIFFER, G., and COURTH, H.
1929. DER TRANSPORT UND DIE TRANSFORMATION ORGANISCH GEBUNDEN PFLANZENJODS IM TIERKÖRPER. *Biochem. Ztschr.* 213: 74-85.
- (464) PFUTZER, G., and PFAFF, C.
1938. ZUSAMMENHÄNGE ZWISCHEN MINERALSTOFF AUFNAHME UND VITAMINGEHALTEN VON PFLANZEN. *Forschungsdienst. Sonderh.* 7: 104-115.
- (465) PIERRE, W. H., and STUART, A. D.
1933. SOLUBLE ALUMINUM STUDIES: IV. THE EFFECTS OF PHOSPHORUS IN REDUCING THE DETRIMENTAL EFFECTS OF SOIL ACIDITY ON PLANT GROWTH. *Soil Sci.* 36: 211-227.
- (466) PINGREE, M. H.
1906. THE INFLUENCE OF NITROGENOUS, PHOSPHATIC AND POTASSIC FERTILIZERS UPON THE PERCENTAGE OF NITROGEN AND MINERAL CONSTITUENTS OF THE OAT PLANT. *Pa. Agr. Expt. Sta. Ann. Rpt.* 1906: 43-53.
- (467) PIPER, C. S.
1936. THE BORON STATUS OF SOUTH AUSTRALIAN APPLES. *Austral. Council Sci. and Indus. Res. Jour.* 9: 245-248.
- (468) PIROTTA, R.
1906. SULLA POSSIBILITA DI ACCUMULARE ARSENICO NEI FRUTTI DI TALUNE PIANTE. *R. Accad. Naz. dei Lincei. Att. Rend. Cl. di Sci. Fis., Mate. Nat.* (5) 15: 730-731.
- (469) POEHLMAN, J. M.
1935. SOME LIMITATIONS OF PLANT JUICE ANALYSES AS INDICATORS OF THE NUTRIENT NEEDS OF PLANTS. *Amer. Soc. Agron. Jour.* 27: 195-207.
- (470) POPP, MAX.
1922. DIE SOGENANNT "THOMASMEHLSEUCHE" UND IHRE BEKÄMPFUNG. *Landw. Jahrb.* 56: 647-670.
- (471) POTTS, H. W.
1896. DISEASE AMONGST DAIRY CATTLE. *Austral. Farm and Home* 5: 57.
- (472) PRICE, O. B.
1923. FERTILIZERS AFFECT QUALITY OF ALFALFA. *Mich. Agr. Expt. Sta. Quart. Bul.* 5: 158-159.
- (473) RADEMACHER, BERNHARD.
1937. KUPFERGEHALT, KUPFERBEDARF UND KUPFERANEIGNUNGSVERMÖGEN VERSCHIEDENER HAFERSORTEN ALS GRUNDLAGE FÜR DIE ZÜCHTUNG GEGEN DIE HIEDEMOORKRANKHEIT WIDERSTANDSFÄHIGER SORTEN. *Ztschr. f. Pflanzenkrank.* 47: 545-560.
- (474) RADU, I. F.
1936. DER VERLAUF DER QUANTITAVEN AUFNAHME VON N, P₂O₅, K₂O, CaO UND MgO DURCH DIE LUZERNE. *Ztschr. f. Pflanzenernähr., Düngung u. Bodenk.* 45: 189-205.
- (475) REAKES, C. J.
1913-14. BUSH SICKNESS. *New Zeal. Dept. Agr. Ann. Rpts.* 21: 43-46, 1913; 22: 35-37, 1914.

- (476) REAKES, C. J., and ASTON, B. C.
1913. THE BUSH SICKNESS INVESTIGATION. *New Zeal. Jour. Agr.* 6: 399-401, 616-624.
- (477) ——— and ASTON, B. C.
1915. BUSH SICKNESS. WORK AT THE MAMAKU FARM. *New Zeal. Jour. Agr.* 11: 377-383.
- (478) REED, O. E., and HUFFMAN, C. F.
1928. DAIRY CATTLE NEED PHOSPHORUS IN RATION. *Mich. Agr. Expt. Sta. Quart. Bul.* 10: 151-156.
- (479) REID, H. A., and ASTON, B. C.
1910. A MALIGNANT BONE DISEASE OF SHEEP (OSSEOUS CACHEXIA). *New Zeal. Dept. Agr. Ann. Rpt.* 18: 47-52.
- (480) REIMER, F. C., and TARTAR, H. V.
1919. SULFUR AS A FERTILIZER FOR ALFALFA IN SOUTHERN OREGON. *Oreg. Agr. Expt. Sta. Bul.* 163, 40 pp.
- (481) REMINGTON, ROE E., CULP, F. BARTOW, and von KOLNITZ, HARRY.
1929. THE POTATO AS AN INDEX OF IODINE DISTRIBUTION. *Amer. Chem. Soc. Jour.* 51: 2942-2947, illus.
- (482) ——— and SHIVER, H. E.
1930. IRON, COPPER, AND MANGANESE CONTENT OF SOME COMMON FOODS. *Assoc. Off. Agr. Chem. Jour.* 13: 129-132.
- (483) REMY, E.
1922. VERGLEICHENDE UNTERSUCHUNGEN UEBER WEISSEN, GELBEN, ROTEN UND VIOLETEN MAIS. *Ztschr. f. Untersuch. der Nahr. u. Genussmtl.* 44: 209-213.
- (484) RICEMAN, D. S., DONALD, C. M., and PIPER, C. S.
1938. RESPONSE TO COPPER ON A SOUTH AUSTRALIAN SOIL. *Austral. Inst. Agr. Sci. Jour.* 4: 41.
- (485) RICHARDS, MARION BROCK.
1930. MANGANESE IN RELATION TO NUTRITION. *Biochem. Jour.* 24: 1572-1590.
- (486) RIDDLE, G. H.
1938. THE "MINOR ELEMENTS," THEIR OCCURRENCE AND FUNCTION IN PLANT LIFE WITH REFERENCE ABSTRACT BIBLIOGRAPHY. New York, F. A. Russo, Inc., photostaters.
- (487) RIGG, T., and ASKEW, H. O.
1934. SOIL AND MINERAL SUPPLEMENTS IN THE TREATMENT OF BUSH SICKNESS. *Empire Jour. Expt. Agr.* 2: 1-8.
- (488) ——— ASKEW, H. O., and CHITTENDEN, E.
1937. BROWN-HEART OF SWEDES AND TURNIPS IN NELSON DISTRICT: A BORON DEFICIENCY AILMENT. *New Zeal. Jour. Sci. and Technol.* 18: 750-755.
- (489) RIPPERTON, J. C.
1922. MINERAL CONSTITUENTS OF HAWAIIAN VEGETABLES. *Hawaii Agr. Expt. Sta. Ann. Rpt.* 1922: 16.
- (490) ——— GOFF, R. A., EDWARDS, D. W., and DAVIS, W. C.
1933. RANGE GRASSES IN HAWAII. *Hawaii Agr. Expt. Sta. Bul.* 65, 58 pp.
- (491) ROBINET, L.
1934. SUICIDES, TUBERCULOSE, SENILITÉ ET TERRAINS MAGNÉSIENS. *Acad. de Med. [Paris] Bul.* 111: 501.
- (492) ROBINSON, R. R.
1937. SOIL PROPERTIES DETERMINING THE BOTANICAL COMPOSITION OF PASTURES IN WEST VIRGINIA. *Jour. Agr. Res.* 54: 877-897.
- (493) ROBINSON, W. O.
1936. SELENIUM CONTENT OF WHEATS FROM VARIOUS PARTS OF THE WORLD. *Indus. and Engin. Chem., Indus. Ed.* 28: 736-738, illus.
- (494) ———
1938. THE AGRICULTURAL SIGNIFICANCE OF THE MINOR ELEMENTS. *Amer. Fert.* 89: 5-8, 24, 26.
- (495) ——— STEINKOENIG, L. A., and MILLER, C. F.
1917. THE RELATION OF SOME OF THE RARER ELEMENTS IN SOILS AND PLANTS. *U. S. Dept. Agr. Bul.* 600, 25 pp.
- (496) ——— WHETSTONE, R. R. and BYERS, H. G.
1938. STUDIES ON INFERTILE SOILS. II. SOILS HIGH IN BARIUM. *Soil Sci. Soc. Proc.* 3: 87-91.

- (497) ROBINSON, W. O., WHETSTONE, R. R., and SCRIBNER, B. F.
1938. THE PRESENCE OF RARE EARTHS IN HICKORY LEAVES. *Science* 87: 470.
- (498) ROLOFF, F.
1869. UEBER OSTEOMALACIE. *Virchows Arch. f. Path. Anat. u. Physiol.* 46: 305-316.
- (499) RUBINO, MIGUEL C.
1935. NUESTROS CAMPOS SE EMPOBRECEN EN SALES MINERALES Y ESPECIALMENTE EN ACIDO FOSFORICO. *Asoc. Rural del Uruguay Rev.* 62 (6): 35-37.
- (500) RUSOFF, L. L., ROGERS, L. H., and GODDUM, L. W.
1937. QUANTITATIVE DETERMINATION OF COPPER AND ESTIMATION OF OTHER TRACE ELEMENTS BY SPECTROGRAPHIC METHODS IN WIRE GRASSES FROM "SALT SICK" AND HEALTHY AREAS. *Jour. Agr. Res.* 55: 731-738.
- (501) SALGUES, RENÉ.
1938. LE TAUX DU PHOSPHORE DANS LES FOURRAGES DE LÉGUMINEUSES ET DE GRAMINÉES EN ANNÉE SÈCHE ET EN ANNÉE HUMIDE EN SOL CALCAIRE ET EN SOL SILICEUX. *Ann. Agron. [Paris]* 8: 325-335.
- (502) SARATA, UICHIRO.
1937. BIOCHEMISTRY OF COPPER. 17. COPPER AND PIGMENTATION OF LEAVES AND FLOWERS. *Jap. Jour. Med. Sci. II. Biochem. Trans. and Abs.* 3: 197-205.
- (503) SATTERFIELD, GEORGE HOWARD, and JONES, S. O.
1932. COPPER CONTENTS OF SOME FOODS. *Elisha Mitchell Sci. Soc. Jour.* 48: 16.
- (504) SAUSSURE, THEODORE DE.
1804. RECHERCHES CHIMIQUES SUR LA VÉGÉTATION. 327 pp., illus. Paris.
- (505) SAYRE, C. B., WILLAMAN, J. J., and KERTESZ, Z. I.
1931. FACTORS AFFECTING THE QUALITY OF COMMERCIAL CANNING PEAS. *N. Y. State Agr. Expt. Sta. Tech. Bul.* 176, 76 pp.
- (506) SAYWELL, L. G., DIETZ, W. H., and ROBERTSON, P. D.
1934. THE IRON, COPPER, AND MANGANESE CONTENT OF CALIFORNIA PRUNES. *Assoc. Off. Agr. Chem. Jour.* 17: 290-293.
- (507) SCHAIBLE, P. J., BANDEMER, S. L., and DAVIDSON, J. A.
1938. THE MANGANESE CONTENT OF FEEDSTUFFS AND ITS RELATION TO POULTRY NUTRITION. *Mich. Agr. Expt. Sta. Tech. Bul.* 159, 27 pp.
- (508) SCHARRER, K., and SCHROPP, W.
1938. UEBER DIE WIRKUNG DES MAGNESIUM-IONS AUF DEN ERTRAG UND DIE ZUSAMMENSETZUNG DES HAFERS AUF BÖDEN VERSCHIEDENER REAKTION. *Bodenk. u. Pflanzenernähr.* 9-10: 321-336.
- (509) ——— and SCHWAIBOLD, J.
1927. ZUR KENNTNIS DES JODS ALS BIOGENES ELEMENT. *Biochem. Ztschr.* 185: 405-413.
- (510) ——— and STROBEL, A.
1927. DIE JODANREICHERUNG DER PFLANZEN DURCH JODZUFUHR. *Angew. Bot.* 9: 187-199.
- (511) SCHEUNERT, A., SCHATKE, A., and LÖTSCH, E.
1911. CaO-, MgO-, P₂O₅-GEHALT VON HEU UND HAFER, NACH DEREN VERFÜTERUNG PFERDE AN OSTEOMALACIE ERKRANKTEN. *Biochem. Ztschr.* 36: 240-244.
- (512) SCHMIDT, H.
1924. FIELD AND LABORATORY NOTES ON A FATAL DISEASE OF CATTLE OCCURRING ON THE COASTAL PLAINS OF TEXAS (LOIN DISEASE). *Tex. Agr. Expt. Sta. Bul.* 319, 32 pp., illus.
- (513) SCHRUMPF-PIERRON, P.
1932. CAUSES OF RARITY OF CANCER IN EGYPT. *Inst. d'Egypte Bul.* 14: 177-187.
- (514) SCHWEITZER, P.
1898. ASH ANALYSES OF SOME COMMON PLANTS. *Mo. Agr. Expt. Sta. Ann. Rpt.* 1898: 83-84.
- (515) SCOFIELD, CARL S., and WILCOX, L. V.
1930. THE BORON CONTENT OF ORANGES. *Science* 71: 542-543.

- (516) SCOTT, SAMUEL G.
1929. PHOSPHORUS DEFICIENCY IN FORAGE FEEDS OF RANGE CATTLE. *Jour. Agr. Res.* 38: 113-130.
- (517) SEABER, W. MACRO.
1933. BARIUM AS A NORMAL CONSTITUENT OF BRAZIL NUTS. *Analyst* 58: 575-580.
- (518) SHACKLETON, LESLIE, and McCANCE, ROBERT ALEXANDER.
1936. THE IONIZABLE IRON IN FOODS. *Biochem. Jour.* 30: 582-591.
- (519) SHEDD, O. M.
1925. THE RELATION OF SOME CHEMICAL CONSTITUENTS TO THE GRADES OF KENTUCKY TOBACCO. *Ky. Agr. Expt. Sta. Bul.* 258, pp. 33-58.
- (520) ———
1930. THE POTASSIUM, CHLORINE, AND SULFUR CONTENT OF KENTUCKY TOBACCO AS RELATED TO GRADE. *Ky. Agr. Expt. Sta. Bul.* 308, pp. 447-471.
- (521) SHEETS, OLIVE, FRAZIER, ERNESTINE, and DICKINS, DOROTHY.
1931. CONSERVATION OF IRON IN VEGETABLES BY METHODS OF PREPARATION AND COOKING. *Miss. Agr. Expt. Sta. Bul.* 291, 16 pp.
- (522) SHERMAN, W. C., ELVEHEM, C. A., and HART, E. B.
1934. FURTHER STUDIES ON THE AVAILABILITY OF IRON IN BIOLOGICAL MATERIAL. *Jour. Biol. Chem.* 107: 383-394.
- (523) SHOREY, EDMOND C.
1906. THE COMPOSITION OF SOME HAWAIIAN FEEDING STUFFS. *Hawaii Agr. Expt. Sta. Bul.* 13, 23 pp.
- (524) SHORLAND, F. B.
1934. THE ESTIMATION OF ALUMINUM IN PASTURES, WITH SPECIAL REFERENCE TO SOIL CONTAMINATION. *Roy. Soc. New Zeal. Trans. and Proc.* 64: 35-50.
- (525) SIMPSON, B. W.
1931. MINERAL FEEDING EXPERIMENTS. *New Zeal. Jour. Agr.* 42: 18-23.
- (526) SJOLLEMA, B.
1933. OVER HET WEZEN EN DE GENEZING VAN LIKZUCHT. *Tijdschr. v. Diergeneesk.* 60: 1137.
- (527) ———
1933. KUPFERMANGEL ALS URSACHE VON KRANKHEITEN BEI PFLANZEN UND TIEREN. *Biochem. Ztschr.* 267: 151-156.
- (528) SKINNER, J. T., and PETERSON, W. H.
1928. THE IRON AND MANGANESE CONTENT OF FEEDING STUFFS. *Jour. Biol. Chem.* 79: 679-687.
- (529) SMIENOVA, M. I., and LAVROVA, M. N.
1934. VARIABILITY IN THE CHEMICAL COMPOSITION OF DIFFERENT VARIETIES OF SOYBEANS. *Trudy Prikl. Bot., Genet. i Selekt. (Bull. Appl. Bot., Genet., and Plant Breeding)* (3) 5: 73-102.
- (530) SMITH, G. ENNIS.
1917. A STUDY OF THE IODINE REQUIREMENTS OF THE PREGNANT SOW. *Jour. Biol. Chem.* 29: 215-225.
- (531) SMITH, M. I., and WESTFALL, B. B.
1937. FURTHER FIELD STUDIES ON THE SELENIUM PROBLEM IN RELATION TO PUBLIC HEALTH. [U. S.] *Pub. Health Rpts.* 52: 1375-1384.
- (532) SOTOLA, JERRY.
1927. RELATION OF MATURITY TO THE NUTRITIVE VALUE OF FIRST, SECOND, AND THIRD CUTTINGS OF IRRIGATED ALFALFA. *Jour. Agr. Res.* 35: 361-383.
- (533) ———
1937. THE CHEMICAL COMPOSITION AND NUTRITIVE VALUE OF CERTAIN CEREAL HAYS AS AFFECTED BY PLANT MATURITY. *Jour. Agr. Res.* 54: 399-415.
- (534) SOXHLET, F. V.
1907. VERSUCHE ZUR BEKÄMPFUNG DER LECKSUCHT DER KÄLBER IM DONAUMOOS. *Vierteljahrsschr. d. Bayerischen Landw. Rats* 12: 283-294.
- (535) SPRENGEL, CARL.
1845. DIE LEHRE VOM DÜNGER. *Ed. 2*, 512 pp. Leipsig.

- (536) STAHL-SCHRODER, M.
1904. KANN DIE PFLANZENANALYSE UNS AUFSCHLUSS ÜBER DEN GEHALT AN ASSIMILIERBAREN NÄHRSTOFFEN IM BODEN GEBEN? Jour. f. Landw. 52: 31-95, 193-268.
- (537) STALÉ, J.
1937. CONTRIBUTION À L'ÉTUDE DE L'INFLUENCE DES ENGRAIS PHOSPHATÉS ET POTASSIQUES SUR LA COMPOSITION MINÉRALE DES FOINS DE PRAIRIES NATURELLES. Landw. Jahrb. des Schweiz 51: 418-430.
- (538) STANFIELD, KENNETH E.
1935. DETERMINATION OF MOLYBDENUM IN PLANTS AND SOILS. Indus. and Engin. Chem., Analyt., Ed. 7 pp. 273-274.
- (539) STEINKOENIG, L. A.
1919. RELATION OF FLUORINE IN SOILS, PLANTS, AND ANIMALS. Indus. and Engin. Chem. 11: 463-465.
- (540) STEWART, JOHN, and SMITH, EDWIN S.
1922. SOME RELATIONS OF ARSENIC TO PLANT GROWTH II. Soil Sci. 14: 119-126.
- (541) STIEBELING, HAZEL K.
1932. THE IRON CONTENT OF VEGETABLES AND FRUITS. U. S. Dept. Agr. Cir. 205, 19 pp.
- (542) STOKLASA, JULIUS.
1926. DIE PHYSIOLOGISCHE FUNKTION DES JODS BEIM BAU- UND BETRIEBS-STOFFWECHSEL IN DER CHLOROPHYLLHALTIGEN UND CHLOROPHYLLOSEN ZELLE. Biochem. Ztschr. 176: 38-61.
- (543) SULLIVAN, B.
1933. THE INORGANIC CONSTITUENTS OF WHEAT AND FLOUR. Cereal Chem. 10: 503-514.
- (544) SVANBERG, OLAF.
1934. OM DE LANTBRUKSKEMISKA FÖRÅTSÄTTNINGARNA FÖR UPPKOMSTEN AV LOKALBUNDA ANÄMISJUKDOMAR I NORRLAND. Lantbruks-Hogskol. Ann. 1: 209-250.
- (545) ———
1935. FORTGESetzte UNTERSUCHUNGEN UEBER DIE AGRIKULTURCHEMISCHEN VORAUSSETZUNGEN FÜR ÖRTLICH BEDINGTE ANÄMIEKRANKHEITEN BEIM PFERDE. Ann. Landw. Hochschule Schwedens 5: 355-375.
- (546) SVANBERG, OLAF, HANNERZ, ERIK, and WIJKSTROM, TORGIL.
1935. ANALYSER AV STRAFODERVEGETATION FRÅU SURA JORDAR I NORRBOTTEN. Lantbruks-Högelskol. Ann. 2: 31-50.
- (547) SYKES, P. H.
1934. AN IODINE SURVEY OF NEW ZEALAND LIVESTOCK. II. THE SHEEP OF THE WAIRARAPA DISTRICT. Roy. Soc. New Zeal. Trans. and Proc. 64: 17-34.
- (548) TAKEI, U.
1937. RELATION BETWEEN ENDEMIC GOITRE IN THE JEHOLO DISTRICT AND THE IODINE CONTENT IN PRINCIPAL FOOD PRODUCED IN THAT DISTRICT. 1. IODINE CONTENT IN VEGETABLES FROM GOITROUS AND NONGOITROUS REGIONS. 2. IODINE CONTENT IN GRAINS FROM GOITROUS AND NON-GOITROUS REGIONS. Jour. Orient. Med. 27: 134-135.
- (549) TAYLOR, N. H.
1930. THE RELATION OF GEOLOGY TO SHEEP SICKNESS IN MAIROA DISTRICT. New Zeal. Jour. Sci. and Technol. 12: 1-10.
- (550) TELLER, G. L.
1893. FODDER. Ark. Agr. Expt. Sta. Bul. 24 pp. 111-142.
- (551) ———
1896. CONCERNING WHEAT AND ITS MILL PRODUCTS. PART II. Ark. Agr. Expt. Sta. Bul. 42, pp. 75-102.
- (552) TERLIKOWSKI, F., and GORNICKI, T.
1933. TITANIUM CONTENTS OF SOME CULTURAL PLANTS. Polish Agr. Forest Ann. 29: 289-298.
- (553) ——— and NOWICKI, B.
1932. BORON CONTENT OF CERTAIN SOILS, PLANTS AND POTASSIC FERTILIZERS. Polish Agr. Forest Ann. 28: 135-144.

- (554) THEILER, ARNOLD, GREEN, H. H., and DU TOIT, P. J.
1924. PHOSPHORUS IN THE LIVESTOCK INDUSTRY. Union So. Africa Dept. Agr. Jour. 8: 460-504, illus.
- (555) ——— VILJOEN, P. R., GREEN, H. H., DU TOIT, P. J., MEIER, HANS, and ROBINSON, E. M.
1927. So. Africa Dept. Agr. and Forestry, 11th and 12th Rpt. Dir. Vet. Ed. and Res., part II, January 1927: 821-1362.
- (556) THOMAS, R. P.
1935. EFFECT OF FERTILIZER TREATMENTS OF A SOIL ON THE QUALITY AND YIELD OF TOMATOES. Md. Agr. Expt. Sta. Bul. 386, pp. 369-390.
- (557) THOMAS, WALTER.
1929. BALANCED FERTILIZERS AND LIEBIG'S LAW OF THE MINIMUM. Science 70: 382-384.
- (558) ———
1932. THE RECIPROCAL EFFECTS OF NITROGEN, PHOSPHORUS, AND POTASSIUM AS RELATED TO THE ABSORPTION OF THESE ELEMENTS BY PLANTS. Soil Sci. 33: 1-20.
- (559) ———
1934. MISCONCEPTION RELATIVE TO THE MINERAL COMPOSITION OF PLANTS. Science 80: 587.
- (560) ———
1937. FOLIAR DIAGNOSIS: PRINCIPLES AND PRACTICE. Plant Physiol. 12: 571-600.
- (561) THOMPSON, ALICE R.
1908. CHEMICAL STUDIES OF RICE AND RICE PRODUCTS. Hawaii Agr. Expt. Sta. Ann. Rpt. 1908: 51-59.
- (562) TILT, JENNIE, and WINFIELD, MAY.
1928. CHEMICAL COMPOSITION OF WEST INDIAN SEEDLING AVOCADOS. Jour. Home Econ. 20: 43-46.
- (563) TOGASHI, KOGO.
1938. EINE BOTANISCHE UNTERSUCHUNG AUF DER YANAGAWAWEIDE AUS ANLASS EINER ENDEMISCHEN PFERDEKRANKHEIT. Bot. Mag. [Tokyo] 52: 627-635.
- (564) TRAUlsen, HERMANN.
1937. DIE LECKSUCHT DER RINDER IN SCHLESWIG-HOLSTEIN. Landw. Vers. Sta. 128: 89-125.
- (565) TROWBRIDGE, P. F., HAIGH, L. D., and MOULTON, C. R.
1915. STUDIES OF THE TIMOTHY PLANT. Mo. Agr. Expt. Sta. Res. Bul. 20, 67 pp.
- (566) TRUE, RODNEY H., BLACK, OTIS F., and KELLY, JAMES W.
1918. ASH CONTENT IN NORMAL AND IN BLIGHTED SPINACH. Jour. Agr. Res. 15: 371-375.
- (567) TUFF, PER.
1923. OSTEOMALACIA AND ITS OCCURRENCE IN CATTLE IN NORWAY. World's Dairy Cong. Proc. 2: 1494-1501. [The same material appears under the same title in Jour. Compar. Path. and Ther. 36: [143]-155, illus.]
- (568) TUORILA, PAULI.
1929. CHEMICAL COMPOSITION OF POTATOES IN FINLAND. Suomen Suoviljelysuhdis. [Finska Mosskulturför.] Tieteell. Julkaisu. 11: 1-73.
- (569) TYSON, JAMES.
1930. INFLUENCE OF SOIL CONDITIONS, FERTILIZER TREATMENTS AND LIGHT INTENSITY ON GROWTH, CHEMICAL COMPOSITION AND ENZYMIC ACTIVITIES OF SUGAR BEETS. Mich. Agr. Expt. Sta. Tech. Bul. 108, 44 pp.
- (570) UNDERWOOD, E. J., and FILMER, J. F.
1935. THE DETERMINATION OF THE BIOLOGICALLY POTENT ELEMENT (COBALT) IN LIMONITE. Austral. Vet. Jour. 11: 84-92.
- (571) [UNITED STATES] BUREAU OF ANIMAL INDUSTRY.
1898. OSTEOMALACIA OR "CREEPS." U. S. Bur. of Anim. Indus. Ann. Rpt. 15: 530.

- (572) VANDECAVEYE, S. C., and BOND, L. V.
1936. YIELD AND COMPOSITION OF ALFALFA AS AFFECTED BY VARIOUS FERTILIZERS AND SOIL TYPES. Amer. Soc. Agron. Jour. 28: 491-505.
- (573) VEDRODE, VICTOR.
1896. DAS KUPFER ALS BESTANDTHEIL UNSERER VEGETABILIIEN. Chem. Ztg. 20: 399-400.
- (574) VELU, H.
1931. UN CHAPITRE DÉLAISSÉ DE LA PATHOLOGIE MAROCAINE: LES MALADIES DE LA NUTRITION. Acad. Vét. de France Bul. 84: 71-81.
- (575) VILJOEN, P. R.
1918. INVESTIGATIONS INTO LAMZIEKTE IN CATTLE. So. Africa Dept. Agr. and Forestry, Rpt. Dir. of Vet. Res. 5 and 6: 257-320.
- (576) VIRTANEN, ARTTURI I.
1928. UEBER DIE EINWIRKUNG DER BODENOZIDITÄT AUF DAS WACHSTUM UND DIE ZUSAMMENSETZUNG DER LEGUMINOSEPFANZEN. Biochem. Ztschr. 193: 300-312.
- (577) VOGEL, F., and VON HOSSLIN, R.
1938. SORTEN- UND STANDORTSEINFLÜSSE AUF DIE GÜTE DES SAUERKRAUTES. Bodenk. u. Pflanzenernähr. 9-10: 665-696.
- (578) WAGNER, KURT.
1936. UEBER DEN BARIUMGEHALT DER PARA-NUSS. Jour. f. Prakt. Chem. 147: 110-112.
- (579) WATKINS, W. E.
1937. THE CALCIUM AND PHOSPHORUS CONTENTS OF IMPORTANT NEW MEXICO RANGE FORAGES. N. Mex. Agr. Expt. Sta. Tech. Bul. 246, 75 pp.
- (580) WEATHERS, E. K.
1938. MINERAL AND NITROGEN CONTENT OF LESPEDEZAS AND OTHER HAY CROPS IN TENNESSEE. Tenn. Agr. Expt. Sta. Bul. 166, 32 pp.
- (581) WEBSTER, JAMES E., and JANSMA, FRANK.
1929. A COMPARISON OF THE COPPER CONTENT OF OKLAHOMA WHEAT WITH THOSE OF OTHER STATES. Science 70: 174.
- (582) WELCH, HOWARD.
1924. BONE CHEWING BY CATTLE. Mont. Agr. Expt. Sta. Cir. 122, 8 pp., illus.
- (583) _____
1928. GOITER IN FARM ANIMALS. Mont. Agr. Expt. Sta. Bul. 214, 27 pp.
- (584) WERNER, ALBERT.
1934. UEBER DEN MINERALSTOFFGEHALT DEUTSCHER FUTTERMITTEL UND SEINE BEEINFLUSSUNG DURCH DÜNGUNG UND HERKUNST. Landw. Vers. Sta. 121: 223-280.
- ✓(585) WESTER, D. H.
1921. UEBER DEN MANGANGEHALT VON (HOLLÄNDISCHEN) SAMEN. Biochem. Ztschr. 118: 158-163.
- (586) _____
1923. UEBER DEN MANGANGEHALT EINIGER HOLLÄNDISCHER BODENARTEN UND EINIGE BEMERKUNGEN DAZU. Internatl. Mitt. f. Bodenk. 13: 1-5.
- (587) WESTGATE, J. M.
1933. Hawaii Agr. Expt. Sta. Rpt., 27 pp.
- (588) _____
1935. EGGPLANT. Hawaii Agr. Expt. Sta. Rpt., 1935: 26.
- (589) WHALLEY, MURIEL E.
1937. BIBLIOGRAPHY ON THE INFLUENCE OF MINERAL DEFICIENCIES ON GROWTH AND REPRODUCTION OF FARM ANIMALS. 89 pp. Nat. Res. Council of Canada, Ottawa.
- (590) WHEELER, H. J., HARTWELL, B. L., KELLOGG, J. W., and STEEL, MATHEW.
1906. THE FUNCTIONS OF SODIUM SALTS. R. I. Agr. Expt. Sta. Ann. Rpt. 19: 186-316.

- (591) WIADDROWSKA, JANINA.
1936. THE EFFECT OF THE ADDITION OF MAGNESIUM TO THE SOIL ON THE MAGNESIUM CONTENT OF PLANTS. Polish Agr. Forest Ann. 37: 307-331.
- (592) WILEY, HARVEY W.
1901. THE SUNFLOWER PLANT. U. S. Dept. Agr., Div. Chem. Bul. 60, 31 pp.
- (593) WILKINS, LOUIS K.
1917. THE HIGH CALCIUM CONTENT OF SOME CUCURBIT VINES. N. J. Agr. Expt. Sta. Bul. 310, 20 pp.
- (594) WILLIAMS, K. T., and WHETSTONE, R. R.
1940. ARSENIC DISTRIBUTION IN SOILS AND ITS PRESENCE IN CERTAIN PLANTS. U. S. Dept. Agr. Tech. Bul. 732, 20 pp., illus.
- (595) WILLIS, L. G.
1939. BIBLIOGRAPHY OF REFERENCES TO THE LITERATURE ON THE MINOR ELEMENTS AND THEIR RELATION TO PLANT AND ANIMAL NUTRITION. Ed. 3, 488 pp., New York.
- (596) WIMER, D. C.
1937. COMPOSITION OF MATURE CORN STOVER. Ill. Agr. Expt. Sta. Bul. 437, pp. 173-272.
- (597) WINTER, O. B., and BIRD, O. D.
1929. THE DETERMINATION OF ALUMINUM IN PLANTS. II. ALUMINUM IN PLANT MATERIALS. Amer. Chem. Soc. Jour. 51: 2964-2968.
- (598) WOODBRIDGE, C. G.
1937. THE BORON CONTENT OF APPLE TISSUES AS RELATED TO DROUGHT SPOT AND CORKY CORE. Sci. Agr. 18: 41-48.
- (599) WOODMAN, HERBERT ERNEST; BLUNT, DENZIL LAYTON; and STEWART, JAMES.
1926. NUTRITIVE VALUE OF PASTURE. I. SEASONAL VARIATIONS IN THE PRODUCTIVITY, BOTANICAL AND CHEMICAL COMPOSITION, AND NUTRITIVE VALUE OF MEDIUM PASTUREAGE ON A LIGHT SANDY SOIL. Jour. Agr. Sci. [England] 16: [205]-274, illus.
- (600) ——— EVANS, R. E., and EDEN, A.
1936. THE COMPOSITION AND NUTRITIVE VALUES OF MARROW STEM KALE AND THOUSAND-HEAD KALE. Jour. Agr. Sci. [England] 26: 212-238.
- (601) ——— EVANS, R. E., and NORMAN, D. B.
1933. NUTRITIVE VALUE OF LUCERNE. I. PRELIMINARY STUDIES OF YIELD, COMPOSITION AND NUTRITIVE VALUE. SEASON 1932. Jour. Agr. Sci. [England] 23: 419-458.
- (602) ——— EVANS, R. E., and NORMAN, D. B.
1934. NUTRITIVE VALUE OF LUCERNE. Jour. Agr. Sci. [England] 24: 283-311.
- (603) WOODRUFF, SYBIL, and KLAAS, HELEN.
1938. A STUDY OF SOYBEAN VARIETIES WITH REFERENCE TO THEIR USE AS FOOD. Ill. Agr. Expt. Sta. Bul. 443, pp. 425-467.
- (604) WRANGELL, M. V.
1927. DAS JOD ALS PFLANZENNÄHRSTOFF. Naturwissenschaften 15: 70-73.
- (605) YOUNG, R. S.
1935. CERTAIN RARER ELEMENTS IN SOILS AND FERTILIZERS AND THEIR ROLE IN PLANT GROWTH. N. Y. (Cornell) Agr. Expt. Sta. Mem. 174, 70 pp.
- (606) ZITKOWSKI, H. E., POTVLIET, M. P., and REED, I. W.
1916. COMPOSITION OF SUGAR BEETS AT VARIOUS STAGES OF THE GROWING SEASON. Sugar 18: 296-298.
- (607) ZUNTZ.
1912. PHYSIOLOGISCHE BEDEUTUNG DES NACH WITTERUNG UND BODEN WECHSELNDEN MINERALGEGHALTES DER FUTTERMITTEL. Deut. Landw. Gesell. Jahrb. 27: 570-594.

SOURCES OF UNPUBLISHED MATERIAL

In May of 1937 Dr. James T. Jardine, Director of Research, United States Department of Agriculture, invited the workers in the State agricultural experiment stations and in the Department to participate in a cooperative survey of unpublished data on the relationship of the chemical composition of plants to the soils in which they were grown. A large amount of material was received, and reference has been made in this report to this unpublished material as follows:

- (1x) BAILEY, C. H., and HUTCHINSON, WILLIS S. Minn. Agr. Exp. Sta. Data obtained through the U. S. Dept. Agr., Bur. of Home Econ.
- (2x) BIZZELL, J. A. N. Y. (Cornell) Agr. Expt. Sta.
- (3x) BRANNEN, C. O. Ark. Agr. Expt. Sta.
- (4x) BURNHAM, MARY C., MARKS, CATHERINE G., TAYLOR, DONA G., and GERHARDT, WARDA G.
 THE EFFECT OF VARIOUS FACTORS ON THE VARIATION IN THE CALCIUM AND PHOSPHORUS CONTENTS OF PEAS. Theses, Wis. Univ. Submitted for this survey by W. H. Peterson, Wis. Agr. Expt. Sta.
- (5x) COLEMAN, J. M., and RUPRECHT, R. W.
 THE EFFECT OF FERTILIZERS AND SOIL TYPES ON THE MINERAL CONTENT OF PLANTS. Thesis, Fla. Univ. Data obtained through the U. S. Dept. Agr., Bur. Home Econ.
- (6x) CULBERTSON, C. C. Iowa Agr. Expt. Sta.
- (7x) DUTCHER, R. ADAMS, and HALEY, D. C.
 Pa. Agr. Expt. Sta. Data obtained through the U. S. Dept. Agr., Bur. Home Econ.
- (8x) EISENMENGER, WALTER S. Mass. Agr. Expt. Sta.
- (9x) FLETCHER, S. W. Pa. Agr. Expt. Sta.
- (10x) FORT, C. A.
 EFFECT OF LOCATION AND VARIETY ON THE MINERAL CONTENT OF SUGARCANE. U. S. Dept. Agr., Bur. Chem. and Soils.
- (11x) HELLER, V. G. Okla. Agr. Expt. Sta.
- (12x) HOF, JOSEPHINE K., and SCHWOEGLER, JUNE K.
 THE MANGANESE AND COPPER CONTENTS OF WISCONSIN POTATOES. Theses, Wis. Univ. Data submitted by W. H. Peterson, Wis. Agr. Expt. Sta.
- (13x) JONES, H. D. Mimeograph 1934. Submitted by the U. S. Dept. Agr., Bur. Home Econ.
- (14x) MCHARGUE, J. S. Ky. Agr. Expt. Sta.
- (15x) MAGISTAD, O. C. Hawaii Agr. Expt. Sta.
- (16x) MOORE, C. C. U. S. Dept. Agr., Div. Chem.
- (17x) PIERRE, W. H. W. Va. Agr. Expt. Sta.
- (18x) ROBINSON, W. O. U. S. Dept. Agr., Bur. Plant Indus.
- (19x) SMITH, JOHN B. R. I. Agr. Expt. Sta.
- (20x) TOBISKA, J. W. Colo. Agr. Expt. Sta.
- (21x) VANDECAVEYE, S. C. Wash. Agr. Expt. Sta.
- (22x) WEATHERS, E. K. Tenn. Agr. Expt. Sta.
- (23x) WILEY, HARVEY W., MOORE, C. C., and EWELL, E. E.
 COMPARATIVE FERTILITY AND NITRIFYING POWER OF SOILS. Unpublished manuscript. U. S. Dept. Agr., Bur. Chem.

APPENDIX

TABLES OF CROP COMPOSITION

In the following tables there is tabulated the mineral composition, insofar as it is known, of the principal crops of interest in the United States. The data were compiled from several hundred published foreign and domestic reports and from unpublished data submitted by State agricultural experiment stations and by various bureaus of the United States Department of Agriculture. In every case in which individual analyses have been used, reference to the source of the data is given, and in addition the sources of data used in the summaries in the minor-element tables are given. Thus, the reader can quickly verify almost any item in which he is particularly interested.

The data have been chosen with every possible care, and particular attention was given to methods of handling the crops and to methods of chemical analysis. Thus, data on forages sampled from weathered materials are not included, and the results of pot tests or other artificial conditions have not been compiled. Data from other compilations have not been included, all data in this compilation having been obtained from original sources.

Furthermore, only analyses published on the moisture-free basis or those that could be calculated from given moisture contents to that basis were used. Moisture contents of forages and most vegetables vary considerably after harvest, and analyses on the fresh basis have little meaning unless the moisture content at the time of analysis is also given.

It was desirable in many instances to discard certain data that seemed to be entirely out of agreement with most of the known results. This was ordinarily done, however, only where a large number of values were available for comparison. Many extremes have been included where the lack of data did not justify their elimination, but occasionally in these instances the reader's attention is called in the tables to data that do not seem to be reasonable.

It is believed that the data in these tables should be of value to nutritionists, agronomists, and others interested in the composition of crops, and the descriptions of the crops and of the soils and the fertilizers used have been made with this in mind.

TABLE 20.—The principal chemical elements in some important crops

[Moisture-free basis]

Plant or plant part, soil type, and location ¹	Fertilizer used ²	Analyses	K	Ca	Mg	P	S	Reference No. ³
		Number	Percent ⁴	Percent ⁴	Percent ⁴	Percent ⁴	Percent ⁴	
Alfalfa, above-ground portion, in bud, all cuttings:	(None)	1						
	L	4						
	P	1						
	P.L	4						
	K	1						
	K.L	4						
	P.K	1						
	(P.K.L)	4						
Summary:								
Maximum			3.31	3.40		.45	0.96	
Minimum			1.10	1.56		.28	.24	
Mean			2.37(13)	2.45(33)	0.25(1)	.56(33)	.63(13)	
Alfalfa, above-ground portion, one-tenth to one-quarter bloom, all cuttings:								
Burgin silt clay loam, Kentucky	None	1	1.59	2.39		.34		14x
Maury silt loam, Kentucky	.do	1	1.75	2.32		.32		
Summary:								
Maximum			2.76	3.23		.43		
Minimum			1.59	1.10		.16		
Mean			1.93(4)	2.34(31)	.34(1)	.32(32)	.36(1)	
Alfalfa, above-ground portion, half bloom, all cuttings:								
Burgin silt clay loam, Kentucky	None	1	1.37	2.83		.31		14x
	do	1		2.94		.23		
	P	2		2.69		.28		
	P.K	2		2.85		.25		
	(None)	2		2.51		.18		
	P	2		2.73		.21		
	P.K	2		2.48		.20		
	(None)	4		2.60		.21		
	P	4		2.45		.24		
	K	2		2.57		.22		
	P.K	6		2.70		.23		
	(None)	1		4.15		.16		
	P	1		3.72		.18		
	P.K	1		3.39		.23		
Brookston clay loam, Michigan	None	1						
	do	1						
	P	2						
	P.K	2						
	(None)	2						
	P	2						
	P.K	2						
	(None)	4						
	P	4						
	K	2						
	P.K	2						
	(None)	1						
	P	1						
	P.K	1						
Brookston loam, Michigan	None	1						
	do	1						
	P	2						
	P.K	2						
	(None)	2						
	P	2						
	P.K	2						
	(None)	4						
	P	4						
	K	2						
	P.K	2						
	(None)	1						
	P	1						
	P.K	1						
Brookston silt loam, Michigan	None	1						
	do	1						
	P	2						
	P.K	2						
	(None)	2						
	P	2						
	P.K	2						
	(None)	4						
	P	4						
	K	2						
	P.K	2						
	(None)	1						
	P	1						
	P.K	1						
Gilford loam, Michigan	None	1						
	do	1						
	P	2						
	P.K	2						
	(None)	2						
	P	2						
	P.K	2						
	(None)	4						
	P	4						
	K	2						
	P.K	2						
	(None)	1						
	P	1						
	P.K	1						

¹ "Summary" items are not summaries of the immediately preceding data but represent independent summaries including other data.
² L=liming material; M=manure; N=nitrogen (mineral or organic); P=phosphate; K=potash; Mn=manganese salts; Cu=copper salts; Na=sodium salts; S=sulfur.
³ Numbers without letters refer to "Literature Cited," p. 59; numbers followed by x refer to "Sources of Unpublished Material," p. 91.
⁴ Figures in parentheses (after percentage of mean) denote number of analyses on which the "summary" data are based.

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
Alfalfa, above-ground portion, half bloom, all cuttings—Con.	None	4		2.68		0.40		229
	L	16		2.63		.31		
	P	4		2.51		.38		
	P.L.	16		2.75		.33		
Isabella sandy loam, Michigan	K	4		2.43		.43		229
	KL	16		2.26		.28		
	PK	4		2.55		.38		
	PKL	16		2.37		.29		
Miami silt loam, Michigan	None	1		3.43		.20		229
	P	1		3.30		.21		
	PK	1	1.96	2.29		.23		
	None	1	3.49	1.51		.23		
Dungeness silt loam, Washington	NP	1	3.32	1.20		.24		229
	PK	1	2.80	1.58		.36		
	NPK	1	2.80	1.45		.33		
	None	3	2.88	1.27		.32		
Folida silt loam, Washington	None	3	2.04	1.46		.42		229
	N	3	2.41	1.32		.36		
	NP	3	2.42	1.31		.33		
	PK	3	2.58	1.46		.40		
Lynden sandy loam, Washington	NPK	3	2.55	1.31		.35		229
	None	2	1.80	2.15		.32		
	N	2	1.92	2.21		.23		
	NP	2	1.62	1.96		.24		
Palouse silt loam, Washington	PK	2	1.96	1.94		.27		572
	NPK	2	1.54	2.35		.30		
	None	1	1.76	1.77		.32		
	do	5	2.30	1.42		.24		
Puget silt loam, Washington	NP	5	1.88	1.38		.26		572
	PK	5	1.78	1.68		.26		
	NPK	5	1.73	1.46		.24		
	None	2	2.43	1.25		.20		
Puget sandy loam, Washington	N	2	2.24	1.27		.21		572
	NP	2	2.52	1.20		.25		
	PK	2	2.57	1.21		.25		
	NPK	2	2.56	1.21		.26		
Ritzville sandy loam, Washington	None	2	2.44	1.94		.32		572
	N	2	2.58	1.78		.29		
	NP	2	2.68	1.64		.29		
	PK	2	2.50	1.75		.27		
NPK	2	2.14	1.60		.29			

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
		Number	Percent	Percent	Percent	Percent	Percent	
Apple, entire fruit; summary:								
Maximum			1.41	0.11	0.059	0.142	0.09	
Minimum			.49	.023	.018	.034	.034	
Mean			.78(34)	.04(26)	.029(14)	.067(35)	.06(2)	
Apple, edible portion; summary:								
Maximum			.90	.177	.113	.113		
Minimum			.62	.021	.055	.055		
Mean			.74(25)	.077(24)	.071(27)	.071(27)		
Barley, grain; summary:								
Maximum			.76	.150	.22	.62	.28	
Minimum			.30	.015	.07	.15	.08	
Mean			.55(133)	.064(252)	.14(70)	.41(316)	.17(165)	
Standard deviation			±.03	±.022	±.03	±.08	±.03	
Mode			.57-.60	.050-.059	.12-.13	.45-.48	.16-.18	
Barley, straw; summary:								
Maximum			1.95	.60	.13	.56	.23	
Minimum			1.08	.15	.04	.04	.08	
Mean			1.48(10)	.35(18)	.08(10)	.12(53)	.15(9)	
Beans, field (<i>Phaseolus vulgaris</i> L.); mature seed:								
Bridgelyhampton very fine sandy loam, Rhode Island								
Summary:		1	1.87	.100	.145	.319	.552	19x
Maximum		1	1.62	.28	.25	.78	.55	
Minimum			1.07	.07	.11	.26	.06	
Mean			1.41(16)	.17(16)	.19(17)	.63(40)	.26(16)	
Beans, field (<i>P. vulgaris</i>); straw, without hulls, summary:								
Maximum			1.50	2.96	.20	.19	.12	
Minimum			.75	1.06	.08	.07	.11	
Mean			1.22(3)	1.71(3)	.13(3)	.11(27)	.11(3)	
Beans, field (<i>P. vulgaris</i>); hulls only, summary:								
Maximum			2.77	1.12	.23	.16	.06	
Minimum			1.34	.62	.16	.08	.04	
Mean			2.29(3)	.83(3)	.19(3)	.11(3)	.05(3)	
Beans, garden (<i>P. vulgaris</i>); edible pods and seeds:								
St. Lucie series, Florida		4	.97	.97	.67	.67	.67	5x
Summary:								
Maximum			3.23	1.21	.52	.84	.84	
Minimum			1.49	.41	.05	.20	.05	
Mean			2.85(6)	.64(53)	.34(11)	.48(60)	.37(1)	
Beans, garden (<i>P. vulgaris</i>); immature seed; grown in Kentucky		1	2.00	.271	.324	.744	.867	379
Beans, garden (<i>P. vulgaris</i>); edible hulls; grown in Kentucky		1	2.58	.671	.341	.563	.191	379
Beans, lima (<i>P. vulgaris</i>); mature seed; grown in Kentucky		1	1.89	.104	.200	.412	.227	379
Beans, mung; mature seed		1	.62	.61	.17	.39	.02	85

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analy- ses	K Percent	Ca Percent	Mg Percent	P Percent	S Percent	Refer- ence No.
Bur-clover (<i>M. hispida</i>), above-ground portion, advanced stage of maturity but still green; grown in California.		2		0.94		0.26		233
Cabbage, edible portion: Hartsell series, Alabama	NPK	1		1.50	0.032	.692		13x
	Na	1		1.22	.040	.645		
	Compost	1		.413	.024	.442		
Norfolk series, Alabama	NPK	1		1.776	.110	.692		5x
	NPK	1		1.206	.071			
	NPK	1		.58		.51		
Coxville series, Florida	NPK	8		.84	.69			495
	NPKMn	2		.76	.61	.61		
	NPKOu	2		.76	.57	.57		
Leon series, Florida	NPK	1	3.66	1.31	.56	.56	1.00	19x
Parkwood series, Florida	NPK	12	3.48	1.43	.233	.40		
Cecil clay, North Carolina	NPK	4	3.19			.18		
Bridgehampton very fine sandy loam, Rhode Island	NPK	1				.48		460
	NPK	1				.15		
	NPK	1				.50		
Clyde silt loam, Wisconsin	None	2	4.03	.63		.40		459, 460
Miami silt loam, Wisconsin	.60	4		.69		.46		
Cabbage, edible portion; summary:		9						
Maximum			9.02	1.78	.38	.77	1.91	
Minimum			1.03	.41	.02	.13	.82	
Mean			2.71 (34)	.73 (89)	.16 (13)	.38 (113)	1.16 (4)	
Standard deviation				±0.25		±.12		
Canada bluegrass (<i>Poa compressa</i> L.), above-ground portion, maturity unknown; summary:								
Maximum			4.19	.92	.33	.64		
Minimum			.82	.15	.07			
Mean			2.04 (19)	.39 (18)	.16 (11)	.25 (20)		
Carrots, roots:								
GloUCEster stony loam, New Hampshire		1	5.92	.376	.187	.35	.15	495
Bridgehampton very fine sandy loam, New Jersey	NPK	6	3.37	.32	.17	.32		19x
	NPK	2		.38	.16	.47		
Summary:								
Maximum			5.95	.56	.25	.65	.18	
Minimum			.46	.24	.12	.14	.13	
Mean			2.10 (27)	.40 (35)	.17 (24)	.33 (53)	.16 (5)	
Caniflower, edible portion; summary:								
Maximum			3.71	.71	.29	.88	1.13	
Minimum			3.45	.13	.24	.51	1.01	
Mean			3.58 (2)	.35 (5)	.26 (3)	.76 (7)	1.07 (2)	

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
		Number	Percent	Percent	Percent	Percent	Percent	
Clover, red (<i>T. pratense</i>), above-ground portion, in bloom; Hagers-town loam, Pennsylvania.		1	1.97	2.09	0.473	0.247	0.19	495
Summary:								
Maximum								
Minimum								
Mean								
Clover, red (<i>T. pratense</i>), above-ground portion, with blossoms and seeds; Gloucester stony loam, New Hampshire.		1	1.96(12)	1.67(11)	.51(9)	.25(12)	.19(1)	495
Clover, red (<i>T. pratense</i>), above-ground portion, end of bloom; grown in Germany.			1.53	1.45	.363	.271	.16	
Clover, red (<i>T. pratense</i>), above-ground portion, maturity unknown;		8	1.78	1.81	.53	.25		250
Burgin silty clay loam, Kentucky	None	6	1.76	2.19		.33		14x
Mauzy silt loam, Kentucky	do	5	2.16	1.91		.38		
Tilist silt loam, Kentucky	do	5	1.04	1.45	.507	.163	.224	495
Penn silt loam (Pennsylvania)		1	1.32	2.07	.694	.227	.19	
Bridgehampton very fine sandy loam, Rhode Island	NPK	15		1.03	.274			19x
Naragansett stony loam, Rhode Island	NPK	2	2.42	1.55	.35	.25	.15	
Summary:								
Maximum								
Minimum								
Mean								
Clover, red (<i>T. pratense</i>); leaves only in bloom; grown in Wales		1	2.82	3.07	.75	.52	.29	
Clover, red (<i>T. pratense</i>); stems only, in bloom; grown in Wales		1	.66	.61	.13	.11	.05	
Clover, red (<i>T. pratense</i>); flowering head, in bloom; grown in Wales		1	1.62(90)	1.53(127)	.37(104)	.22(112)	.16(75)	
Clover, hop (<i>T. procumbens</i> L.); above-ground portion, prebloom; grown in Tennessee.		1		2.030		.231		170
Clover, hop (<i>T. procumbens</i>); above-ground portion, in bloom; grown in Tennessee.		1		1.185	.193	.376		170
Clover, hop (<i>T. procumbens</i>); above-ground portion, in bloom; grown in Tennessee.		1	2.457	1.165		.424		580
Clover, hop (<i>T. procumbens</i>); above-ground portion, in bloom; grown in Tennessee.		1	1.718	1.351	.217	.354		580
Clover, hop (<i>T. procumbens</i>); above-ground portion, bloom dying; grown in Tennessee.		1	1.660	1.029	.127	.236		580
Clover, white (<i>T. repens</i> L.); above-ground portion, beginning bloom; Hagerslow loam, Pennsylvania.		1		1.59	.434	.310		495
Clover, white (<i>T. repens</i>); above-ground portion, end of bloom; summary.		2	3.38	1.05	.338	.34	.25	
Clover, white (<i>T. repens</i>); above-ground portion, maturity unknown;								
Mauzy silt loam, Kentucky		1	4.02	1.56		.428		14x
Merrimac fine sandy loam, Massachusetts		1	2.24	1.61	.25	.45		11
Puget sandy loam, Washington	(None)	5		.98		.61		231
Westmoreland silt loam, West Virginia	(NPK)	5		1.07		.52		
Dekalb silt loam, West Virginia	(NPK)	1		1.61		.291		17x
Huntington silt loam, West Virginia	(NPK)	4		1.96		.339		
		2		.91		.432		

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analy- ses	K	Ca	Mg	P	S	Refer- ence No.
Corn, stover, mature—Continued.	(None)	Number	Percent	Percent	Percent	Percent	Percent	
Dunkirk silty clay loam, New York	(L)	2	1.39	0.25	0.21	0.27	0.23	
	(P)	2	1.67	.29	.28	.21	.21	
	(L)	1	.77	.21	.16	.37	.14	
	(P,L)	1	.69	.26	.11	.25	.11	2x
Potoskey gritty sandy loam, New York	(P)	1	1.42	.15	.23	.36	.16	
	(P,L)	1	1.34	.21	.27	.42	.18	
	(None)	2	1.47	.12	.10	.32	.25	
Vohsias silt loam, New York	(L)	2	1.10	.12	.15	.24	.24	
	(P)	1	1.82	.24	.22	.32	.11	
	(P,L)	1	1.40	.29	.22	.37	.15	
	(None)	1			.22			
	(P)	1			.20			
	(K)	1			.10			
	(N,P)	2			.21			19x
	(N,K)	2			.12			
	(P,K)	2			.24			
	(N,P,K)	1			.21			
Summary:								
Maximum			1.89	.84	.51	.42	.31	
Minimum			.26	.10	.08	.04	.08	
Mean			.74(219)	.32(192)	.24(43)	.15(231)	.17(32)	
Standard deviation			±.37	±.11		±.08		
Mode			.50	.32	.06			
Corn, stover, leaves only, mature:								
Carrington silt loam, Illinois	(P)	14	.44	.37	.124			
Crisie silt loam, Illinois	(P)	26	.32	.31	.133			
Drummer clay loam, Illinois	(P)	6	.72	.25	.072			
Elliot silt loam, Illinois	(P)	5	.40	.15	.106			
Grundy clay loam, Illinois	(P)	10	.47	.41	.093			
Grundy silt loam, Illinois	(P)	29	.36	.31	.103			
Muscatine silt loam, Illinois	(P)	19	.46	.37	.098			
Onango sandy loam, Illinois	(P)	5	.78	.72	.203			596
Putnam silt loam, Illinois	(P)	9	.45	.43	.158			
Saybrook silt loam, Illinois	(P)	9	.42	.25	.120			
Seidell silt loam, Illinois	(P)	13	.48	.27	.118			
Summary:								
Maximum			1.02	.91	.234			
Minimum			.23	.11	.052			
Mean			.42(146)	.34(145)	.117(145)			

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
Flax, seed; summary:		Number	Percent	Percent	Percent	Percent	Percent	
Maximum						.85		
Minimum						.43		
Mean						.59(14)	0.06(1)	
Flax, straw; summary:						.135		
Maximum						.026		
Minimum						.066(8)		
Mean						.13		
Kafir, above-ground portion, cut before heads appeared; grown in Oklahoma		4		.44	.40			
Kafir, heads removed, mature plant; grown in Oklahoma		3		.60				133
Kafir, mature heads and grain only; grown in Oklahoma		3		.12				
Kafir, grain; grown in Ohio		1	.29	.013	.14		.12	192
Kafir, stover; grown in Africa		1	1.47	.18				291
Kafir, above-ground, portion, cut for forage; grown in Hawaii		2	2.23	.13				523
Kale, edible portion:								
Yolo silt loam, California		1	2.38	1.51	.74	.65	1.05	140
Sassaras sandy loam, Maryland		2	3.97	1.97	.25	.73	1.12	
Norfolk loam, Virginia		2	3.80	2.21	.31	.60	1.11	
Summary:								
Maximum			4.37	3.60	.74	.78	1.20	
Minimum			2.38	1.37	.24	.25	.64	
Mean			3.47(6)	2.16(9)	.36(6)	.62(9)	1.04(7)	
Kale, marrow-stem, above-ground portion; grown in England		4	3.35	2.13	.30	.35	1.16	600
Kale, thousand-head; above-ground portion; grown in England		2	2.72	1.75	.28	.35	.98	600
Kentucky bluegrass, young blades; grown in Kentucky		1	2.585	.470	.197	.952	.656	375
Kentucky bluegrass, above-ground portion, 4 to 5 inches high:								
Dunkirk silt loam, New York		1		.23		.59		304
Miami silt loam, Wisconsin	None	8		.57		.44		418
Kentucky bluegrass, above-ground portion, 8 to 10 inches high:	do	5		.55		.41		
Miami silt loam, Wisconsin								
Kentucky bluegrass, above-ground portion, 12 to 15 inches high; grown in Iowa	do	5		.28		.27		6x
Kentucky bluegrass, above-ground portion, clipped at 10-day intervals; Grundy silt loam, Iowa	do	51		.38		.30		6x
Kentucky bluegrass, above-ground portion, clipped at 40-day intervals; Grundy silt loam, Iowa	NPK	29		.38		.35		
Kentucky bluegrass, above-ground portion, first coming into bloom: Volusia silt loam, Ohio	NPK	30		.86		.27		
None		30		.35		.32		
Miami silt loam, Ohio	do	1	2.61	.222		.283		195
Miami clay, Ohio	do	1	1.75	.229		.284		
Yazoo clay, Ohio	do	1	1.86	.254		.247		

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analy- ses	K	Ca	Mg	P	S	Refer- ence No.
		Number	Percent	Percent	Percent	Percent	Percent	
Lespedeza , above-ground portion, maturity unknown:								
Burgin, silt clay loam, Kentucky		3	1.02	1.28		.30		14x
Decatur silt loam, Kentucky		4				.18		
Maury silt loam, Kentucky		3	.94	1.44		.31		22x
Tipton silt loam, Kentucky		6				.13		
Clarksville silt loam, Tennessee		4	1.00	.97	0.17	.16		580, 22x
Cumberland loam, Tennessee		28	1.01	1.04	.22	.24		
Hartsells silt loam, Tennessee		17	1.79	.94	.21	.14		580, 22x
Maury silt loam, Tennessee		6	1.26	1.27	.20	.20		
Lespedeza , above-ground portion, all stages of maturity; summary:								
Maximum			1.05	2.08	.42	.38		
Minimum			.62	.69	.13	.08		
Mean			1.02(144)	1.10(152)	.22(138)	.20(160)		
Standard deviation			±.24	±.22	±.05	±.07		
Mode			.70-.80	.90-1.00	.15-.20	.19-.23		
Lettuce , head:								
Cecil series, Alabama		1		.93	.05	1.05		13x
Norfolk series, Alabama		2		.80	.11	.48		
	(NPK	13		1.25		.75		5x
	(NPKMn	3		1.35		.76		
	(NPKCu	2		1.40		.76		19x
	(NPK	1		1.38		.74		
	(NPK	3	7.46			.65		
	(NPK	1	6.86			.69		
	(NPK	5	6.38					
Lettuce , head; summary:								
Maximum			7.91	1.38	.44	1.05	0.31	
Minimum			2.69	.77	.04	.19	.25	
Mean			5.98(16)	.77(17)	.21(11)	.56(30)	.28(3)	
Mangel-wurzel (<i>Beta vulgaris</i> var. <i>macrorhiza</i>), roots: Bridgehampton very fine sandy loam, Rhode Island								
Summary:								
Maximum		4		.24	.17	.08		19x
Minimum								
Mean			3.87	.31	.36	.29		
			.89	.13	.11	.07		
			2.20(5)	.21(6)	.21(5)	.18(10)	.23(1)	
Mangel-wurzel , crowns only: Bridgehampton very fine sandy loam, Rhode Island		4		.81	.24	.12		19x
Mangel-wurzel , leaves only: Bridgehampton very fine sandy loam, Rhode Island		4		1.76	.77	.16		19x
Millet , variety unknown, above-ground portion, fruiting; grown in Philippine Islands		1		.80		10 1.03		394
Millet , German Golden; above-ground portion, cut for hay: Bridgehampton very fine sandy loam, Rhode Island		5				.14		19x

	(NKL) {PKLMg}							
Millet, Japanese, above-ground portion, cut for hay: Bridgehampton very fine sandy loam, Rhode Island		5	7.42	1.12	.15	19x		
Millet, above-ground portion, variety and maturity unknown, cut for hay; summary:		2			.10			
Maximum			7.59	1.14	.62			
Minimum			1.54	.71	.00			
Mean			3.20(11)	.96(7)	.22(22)			
		1		.94	1.10	394		
Milo, above-ground portion, flowering; grown in Philippine Islands, Oklahoma		10		.58	.18	133		
Milo, above-ground portion, cut before heads appeared; grown in Philippine Islands		1		.13	.04	394		
Milo, above-ground portion, heads removed, mature plant; grown in Oklahoma		10		.64	.12	133		
Milo, mature heads and grain; grown in Oklahoma		10		.18	.30	133		
Milo, above-ground portion, maturity unknown; grown in Kentucky		3	3.06		.38	452, 453		
Mustard, edible leaves; Norfolk series, Alabama		2		1.95	.71	13x		
Summary:					.05			
Maximum			2.59		.97			
Minimum			1.76		.88			
Mean			2.13(5)		.71(5)			
Oats, grain: Bridgehampton very fine sandy loam, Rhode Island	NKL	7			.24	19x		
Summary:								
Maximum			.73	.19	.53			
Minimum			.28	.05	.15			
Mean			.48(99)	.10(170)	.34(233)			
Standard deviation			±.09	±.01	±.08			
Mode			.45-.50	.08-.10	.35-.40			
Oat straw; summary:								
Maximum			3.52	.67	.36			
Minimum			.59	.15	.09			
Mean			1.51(72)	.21(147)	.11(212)			
Above-ground portion, flowering stage; grown in France		1	2.01	.73	.48	58		
Penn silt loam, Pennsylvania					.57			
Chehalis clay loam, Washington		1	2.20	.30	.33	495		
	{None	1	.57	.24	.25			
	{(u)	7	.99	.27	.22			
	{None	1	.20	.20	.25			
	{(u)	4	.38	.23	.20			
	{None	1	.63	.17	.17			
	{(u)	4	.69	.20	.17			
	{None	1	.50	.28	.25			
	{(u)	5	.34	.31	.23			
	{None	1	1.44	.13	.24			
	{(u)	4	1.62	.14	.30			
	{None	2		.29	.29			
	{do	1	1.19	.19	.40			
	{(u)	5	1.51	.18	.27			

¹⁰ Omitted from summary following.
¹¹ Average of all fertilizer treatments. No consistent influence of fertilizers was noticeable, and no experiments were conducted using any single fertilizer elements except nitrogen.

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
		Number	Percent	Percent	Percent	Percent	Percent	
Oats, above-ground portion, grain matured; summary:								
Maximum			1.71	0.51	0.41	0.40	0.09	
Minimum			1.78	.21	.13	.16	.07	
Mean			1.15(6)	.35(9)	.25(9)	.23(6)	.08(5)	19x
Onions, bulb: Bridgehampton very fine sandy loam, Rhode Island.	NPK	1				.42		
Summary:								
Maximum			1.92	.59	.16	.74		
Minimum			1.03	.26	.11	.17		
Mean			1.52(29)	.39(28)	.13(3)	.26(33)	.60(1)	
Parsnip, roots:								
Bridgehampton very fine sandy loam, Rhode Island.	{NP	2	1.35			.31		19x
Summary:	{NPK	6						
Maximum			2.62	.33		.84		
Minimum			.86	.25		.20		
Mean			1.85(12)	.28(6)	.12(1)	.37(14)		
Peas, edible portion, all sizes:								
Dunkirk fine sand, New York.	{NP	11	1.42	.15				
Carrington silt loam, Wisconsin	{NPL	4	1.52	.14				505
Knox silt loam, Wisconsin	{NPK	8	1.49	.14				
Miami silt loam, Wisconsin	{NPKL	8	1.55	.17				
Summary:								
Maximum		17	1.7	.25		.59		4x
Minimum		4	1.18	.18		.58		
Mean		45	1.21	.21		.64		
Peas, foliage of plant:								
Norfolk sandy loam, Virginia.	{None	1	1.89	.43	.22	.78	.15	
Summary:	{L	6						
Maximum			1.79	.10	.15	.23	.09	
Minimum			1.41(54)	.19(127)	.18(6)	.57(100)	.13(4)	
Mean			1.84	4.84	.52	.18		450
Norfolk sandy loam, Virginia.			2.84	6.53	.26	.18		
Summary:								
Maximum			3.15	2.32	.46	.51	.29	
Minimum			1.74	.77	.27	.18	.18	
Mean			1.40(10)	1.37(27)	.37(9)	.28(36)	.23(4)	
Potatoes, tubers:								
Fort Collins loam, Colorado.	{None	5	2.13	.051	.103	.27	.09	257
Summary:	{N	5						
Maximum		6	2.29	.061	.107	.26	.09	
Minimum	{NPK	6	2.37	.18	.18	.32	.24	
Mean	{NPKMn	1	2.19	.19	.19	.36	.25	5x
Bladen series, Florida.	{NPKCu	1	2.56		.19	.35	.25	

Coxville series, Florida.....	(NPK.....	4	2.18	.22	.30	.24	5x
	{NPKMn.....	3	2.12	.22	.28	.25	
	{NPKCu.....	2	2.27	.19	.32	.22	
	{NPKCuMn.....	1	2.38	.20	.28	.25	
Leon series, Florida.....	NPK.....	4	2.32	.21	.45	.24	
Norfolk series, Florida.....	NPK.....	1	2.61	.23	.27	.21	
Gloucester stony loam, New Hampshire.....	NPK.....	1	2.73	.11	.24	.18	495
Hagerstown series, Pennsylvania.....	{NP.....	16	.028		.24		7x
	{NPK.....	1	.051		.24		
	{NPNa.....	5	1.72	.13	.24		19x
Bridgehampton very fine sandy loam, Rhode Island.....		1			.31		
Summary:							
Maximum.....			3.96	.23	.49	.42	
Minimum.....			1.05	.05	.11	.06	
Mean.....			2.28(773)	.13(315)	.25(588)	.12(420)	
Standard deviation.....			±.43	±.03	±.06	±.04	
Mode.....			2.05-2.20	1.11-.12	23-.25	-.09-.10	
Potatoes, vines only; grown in Germany.....		16	1.62	1.15	.26	.68	137,368
Pumpkin, edible portion; summary.....		3	2.47	4.12	.24	.25	
Pumpkin, entire fruit; grown in New Jersey.....		1	2.86	.16	.25	.25	593
Rape, above-ground portion, plant 6-8 inches high, green and succulent; grown in Iowa.....	None.....	1	1.67		.60		6x
Rape, above-ground portion, plant 15 inches high, half grown; grown in Iowa.....		1	2.46		.47		6x
Rape, above-ground portion, plant 20-27 inches high; grown in Iowa.....		3	1.92		.38		6x
Rape, above-ground portion, plant 30-34 inches high; grown in Iowa.....		3	1.96		.33		6x
Rape, above-ground portion, cut before bloom; grown in Massachusetts.....		1	4.36	.36	.28		215
Rape, above-ground portion, flowering stage; grown in France.....		2	3.41	.37	.51	1.16	
Rape, above-ground portion, maturity unknown.....		1		.03	.78		13x
Norfolk series, Alabama.....	{None.....	2			.39		75
	{P.....	8	1.50		.40		
Colts Neck loam, New Jersey.....	NPK.....	6			.22		19x
Summary:							
Maximum.....			5.27		.86	.99	
Minimum.....			1.37		.11	.16	
Mean.....			3.40(6)	.03(1)	.41(33)	.66(5)	
Rape, leaves only, mature plant; grown in Iowa.....		4	1.50(1)		.48		6x
Rape, stems only, mature plant; grown in Iowa.....		4	2.77		.30		6x
Rape, stalks only, mature plant; grown in Iowa.....		4	2.28		.27		6x
Redtop, above-ground portion, full bloom; Maury silt loam, Kentucky.....		2	1.59	.52	.28		14x
Redtop, above-ground portion, stage of maturity unknown:							
	{None.....	1	1.45	.170	.124	.314	14x
	{ML.....	1	1.25	.135	.118	.280	
	{M.....	1	1.28	.164	.143	.259	
	{LSP.....	1	1.28	.168	.161	.259	
	{None.....	4	.74		.44		14x
Maury silt loam, Kentucky.....							

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
Redtop, above-ground portion, stage of maturity unknown—Con.								
Tilist silt loam, Kentucky	{ None NPL NPS PK NPK	Number 1 1 1 1 5	Percent 1.52 1.34 1.28 1.65 1.51	Percent 0.316 0.407 0.290 0.31 0.33	Percent 0.210 0.245 0.11 0.12	Percent 0.112 0.111 0.148 0.20 0.19	Percent 0.250 0.296 0.331	14x 19x
Summary:								
Maximum			3.94	.63	.33	.41	.33	
Minimum			.71	.07	.07	.05	.09	
Mean			2.12(48)	.42(47)	.18(34)	.25(53)	.26(15)	
Redtop, above-ground portion, in seed; grown in Hawaii		1		.26		.11		490
Rice, paddy; summary:								
Maximum			.48	.138	.17	.43		
Minimum			.32	.057	.14	.29		
Mean			.38(15)	.090(10)	.16(7)	.36(18)		
Rice, straw; grown in Hawaii; summary		15	1.64	.30	.11	.13		313, 561
Rutabaga, roots:								
Bridgethampton very fine sandy loam, Rhode Island	{ NPK NKL NPKL	16 1 1				.25 .26 .32		19x
Summary:								
Maximum			4.77	.80	.19	.78		
Minimum			.60	.13	.11	.05		
Mean			1.91(24)	.40(23)	.14(7)	.27(66)	.30(1)	
Rutabaga, tops:								
Bridgethampton very fine sandy loam, Rhode Island	NPK	3				.34		19x
Summary:								
Maximum				2.92	.28	.93		
Minimum				2.12	.17	.29		
Mean			3.84(1)	2.50(5)	.22(4)	.49(5)		
Rye, grain; summary:								
Maximum			.63	.207	.19	.52	.41	
Minimum			.48	.043	.11	.21	.08	
Mean			.53(11)	.115(10)	.14(4)	.37(71)	.18(3)	
Rye-straw; summary:								
Maximum			1.51	.40	.133	.30	.20	
Minimum			.53	.16	.036	.04	.07	
Mean			.97(33)	.28(31)	.073(25)	.10(41)	.11(29)	
Rye, above-ground portion, cut before heads appeared; grown in Oklahoma		13		.290		.231		132
Sorghum, above-ground portion, cut before heads appeared; grown in Oklahoma		18		.36		.103		
Sorghum, above-ground portion, in Oklahoma; summary								
Maximum			1.70	.19	.27	.27	.05	523
Minimum			1.23	.22	.77	.19		420
Sorghum, above-ground portion, cut for fodder; grown in Hawaii		5						
Sorghum, green stalk; grown in Texas		1						

Sorghum, green heads and seed; grown in Texas	1	.55	.22	.11	.51	.06	240
Sorghum, mature stalk; grown in Oklahoma	3		.63		.59		133
Sorghum, mature heads and seed; grown in Oklahoma	3		.081		.22		133
Sorghum, straw; grown in Africa	1	1.20	.16		.13		291
Soybean, leaves; grown in France	1		3.18	.79	.16	.25	335
Soybean, pod; grown in France	1	1.95	.66	.41	.51	.22	335
Soybean, stem; grown in France	1	.67	.42	.42	.20	.27	335
Soybean, hulls and vines; grown in South Carolina	1	.96	.46	.23	.42		369
Soybean, vine, in full bloom; summary	9	2.31	1.47	.73	.39		
Soybean, vine, pods forming; grown in Massachusetts	1	.93	2.10	.76	.31	.18	215
Soybean, hay, maturity unknown:							
	1	1.46	.94	.441	.194	.274	
	1	1.01	1.15	.439	.182	.241	
	2	.96	1.07	.448	.356	.344	
	2	5.12	1.23	5.12	.274	.250	
	2	1.48	.99	3.53	.237	.201	
	2	(MLPK)	6.56	6.56	.214	.333	
	1	(None)	1.05	1.25	.214	.333	
	1	M	1.19	1.19	.148	.306	
	1	1.24	1.19	1.19	.200	.254	
	1	.96	1.20	6.08	.304	.322	
	2	.79	1.09	8.22	.257	.290	
	2	MPK	1.15	6.19	.292	.292	
	3	1.28	1.11	7.20	.265	.259	
	2	MLP	1.11	7.20	.154		
	1	1.16	1.32	.543			
			2.11	.86	.58	.52	
		2.31	.52	.16	.09	.14	
		.84	1.12(119)	.50(98)	.26(124)	.29(31)	
	10	1.24(64)			.40		452
		2.62					
Summary:							
Maximum			.333		.773		
Minimum			.161		.506		
Mean			.232(40)		.668(4)		
Soybean, entire plant including roots; grown in Kentucky.							
Soybean, green, shelled, matured as for food; summary:							
Maximum			.30	.34	1.08	.45	
Minimum			.19	.24	.50	.10	
Mean			.24(9)	.31(7)	.78(37)	.24(6)	
Soybean, mature seed; summary:							
Maximum		2.39	.58	.23	.68	.13x	
Minimum		.81	1.26	.17	.82	.140	
Mean		1.83(29)	1.97	.35	.86	.49	
Spinach, above-ground portion:							
Cecil series, Alabama	1						
Norfolk series, Alabama	2						
Sassafras sandy loam, Maryland	4	4.24	1.97	.35	.61		
	1	8.09	.76		.89		
	3	6.58	1.00	.52	.88		
	12	7.18	1.37		.87		
Bridgehampton, very fine sandy loam, Rhode Island.	2						
	2	9.09	2.78	1.24	1.17	.66	
	2	3.03	.51	.15	.27	.14	
	3	6.16(27)	1.21(45)	.55(27)	.72(62)	.39(11)	
	3	1.61			.83		
Norfolk loam, Virginia							
Summary:							
Maximum							
Minimum							
Mean							
Squash, entire fruit; Bridgehampton very fine sandy loam, Rhode Island.							
	3						

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
Strawberries, fruit: Derby fine sandy loam, Oklahoma	None	Number 16				0.13		252
Maximum			2.35	0.41		.37		
Minimum			1.58	.22		.10		
Mean			2.05(7)	.31(2)		.20(30)		
Sudan grass, above-ground portion, 6 to 12 inches high, entirely edible; grown in Iowa.	None	4		.56		.52		6x
Sudan grass, above-ground portion, 20 to 30 inches high, mostly edible; grown in Iowa.	None	4		.44		.43		6x
Sudan grass, above-ground portion, 30 to 60 inches high, slightly headed, partly edible; grown in Iowa.	None	4		.47		.27		6x
Sudan grass, above-ground portion, mature, partly edible; grown in Iowa.	None	2		.30		.21		6x
Sudan grass, above-ground portion, before flowering; grown in Philippine Islands.	None	1		.44		.88		394
Sudan grass, above-ground portion, flowering; grown in Philippine Islands.	None	1		.29		.59		394
Sudan grass, above-ground portion, early seed stage: Clarksville silt loam, Arkansas.	NPK	5	1.48					3x
Sudan grass, above-ground portion, maturity unknown; summary:								
Maximum				.86	0.43	.40		
Minimum				.43	.26	.19		
Mean				.73(6)	.35(5)	.29(2)	0.05(1)	
Sugar beets, roots; summary:								
Maximum			1.46(1)					
Minimum			1.39	.35	.23	.33	.13	
Mean			.37	.11	.11	.03	.07(9)	
Mean			.64(57)	.22(49)	.16(31)	.11(52)		
Sugar beets, leaves; summary:								
Maximum			7.64	2.83	1.07	.38	.61	
Minimum			1.01	.26	.31	.08	.31	
Mean			3.28(60)	1.14(60)	.51(59)	.22(60)	.48(20)	
Sugarcane leaves; summary:								
Maximum			1.58	.62		.22	.23	
Minimum			.36	.43		.07	.06	
Mean			1.10(10)	.54(6)	.40(1)	.13(10)	.13(5)	
Sugarcane, tops; summary:								
Maximum			2.65	.44	.17	.31		
Minimum			1.46	.13	.13	.10		
Mean			1.83(10)	.21(10)	.14(5)	.17(18)		
Sugarcane, stalks; summary:								
Maximum			1.44	.24	.13	.17	.15	
Minimum			.05	.02	.02	.02	.02	
Mean			.56(87)	.07(43)	.06(34)	.07(96)	.06(34)	

Sunflower, heads minus seeds and husks, mature plant; grown in District of Columbia.	1	9.43	2.49	1.26	.41	.46	592
Sunflower, seed husk, mature plant; grown in District of Columbia.	1	.92	.35	.23	.07	.05	592
Sunflower, seed kernel without husk, mature plant; grown in District of Columbia.	1	.96	.21	.40	1.01	.02	592
Sunflower, leaves, head formed, seed undeveloped; grown in Puerto Rico.	7	3.85	3.29	.088	1.08		209
Sunflower, leaves, mature plant; grown in District of Columbia.	1	1.62	7.64	3.15	.35	.43	592
Sunflower, stalks, head formed, seed undeveloped grown in Puerto Rico.	8	4.58	1.03	.63	.31		209
Sunflower, stalks, mature plant; grown in District of Columbia.	1	3.23	1.72	1.20	.07	.14	209
Sunflower, above-ground portion, cut before bloom; grown in Germany.	1	3.33	2.09	.49	.20	.13	368
Sunflower, above-ground portion, bud appearing on top of plant; Palouse silt loam, Idaho.	6	4.58	2.20	.42	.28	.28	431
Sunflower, above-ground portion, flower 3 inches in diameter, no seed; Palouse silt loam, Idaho.	6	4.20	1.89	.37	.28	.32	431
Sunflower, above-ground portion, cut in full bloom; grown in Germany.	1	2.11	2.06	.50	.18	.14	368
Sunflower, above-ground portion, cut before seed of first bloom was in dough stage; Palouse silt loam, Idaho.	6	4.08	1.80	.37	.27	.26	431
Sunflower, above-ground portion, seeds of first flower in dough stage; Palouse silt loam, Idaho.	6	3.69	1.61	.28	.27	.23	431
Sunflower, above-ground portion, seeds quite hard and rays fallen; Palouse silt loam, Idaho.	6	3.11	1.57	.37	.26	.25	431
Sweetpotatoes, tubers; summary:							
Maximum		1.74	.15	.21	.22		
Minimum		.68	.04	.11	.06		
Mean		1.21 (25)	.08 (9)	.16 (2)	.12 (33)		
Sweetclover (<i>Melilotus alba</i> Desv.), above-ground portion, very young; Palouse silt loam, Washington.	5		1.66		.46	.12 (1)	533
Sweetclover (<i>M. alba</i>), above-ground portion, flower buds just opening:							
Grown in Canada	2	1.88	1.55		.243		200
Palouse silt loam, Washington	1		1.19		.46		533
Sweetclover (<i>M. alba</i>), above-ground portion, in full bloom; summary:							
Maximum		1.81	2.64	.47	.41		
Minimum		1.48	.61		.05		
Mean		1.64 (3)	1.26 (175)	.34 (9)	.13 (176)		
Sweetclover (<i>M. alba</i>), above-ground portion, seeds well filled; grown in Canada.	2	1.27	1.36		1.94		200
Sweetclover (<i>M. alba</i>), above-ground portion, maturity unknown: Bluford silt loam, Illinois	3	1.49			.38	.369	40
Shelby loam, Missouri	1					.237	237
Hagerstown loam, Pennsylvania.	1	2.03	1.19	.293	.227	.70	495
Summary:							
Maximum		2.03	1.53	.35	.39	.70	
Minimum		60	1.19	.29	.12	.25	
Mean		1.30 (10)	1.33 (3)	.32 (2)	.31 (13)	.45 (3)	
Tendergreen, above-ground portion:							
Decatur series, Alabama	1		2.59	.09	.97		
Norfolk series, Alabama	1		2.38	.06	.74		

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
		Number	Percent	Percent	Percent	Percent	Percent	
Timothy, above-ground portion, 2-4 inches high; grown in Iowa.		3	2.25	0.59		0.37		6x
Timothy, above-ground portion, 4-5 inches high; Maury silt loam, Kentucky.		3		.49		.37		14x
Timothy, above-ground portion, 10-12 inches high; summary		2	1.98	.17		.27		
Timothy, above-ground portion, just beginning to head; summary		2	1.49	.23		.21		
Timothy, above-ground portion, full bloom:		2	1.16	.42		.31		14x
Maury silt loam, Kentucky		1	1.64	.20		.17		
Gloucester stony loam, New Hampshire		1	1.75	.16	0.09	.21	0.09	495
Hagerstown loam, Pennsylvania		1	2.08	.26	.15	.24	.19	
Penn silt loam, Pennsylvania		1						
Summary:								
Maximum			2.32	.44	.17	.41	.27	
Minimum			.92	.15	.09	.17	.02	
Mean			1.69(8)	.28(7)	.14(5)	.24(8)	.13(5)	
Timothy, above-ground portion, out of bloom, seed formed; grown in Missouri.		1	1.12			.17		565
Timothy, above-ground portion, seed ripe; summary		3	1.58	.16	.07	.17	.16	
Timothy, above-ground portion, seed all in dough; grown in Missouri.		1	.89			.14		565
Timothy, above-ground portion, cut for hay, maturity unknown:								
Tilist silt loam, Kentucky	(M)	1	1.46	.28	.17	.14	.17	
	(ML)	1	1.47	.31	.20	.20	.20	14x
	(MPL)	2	1.20	.34	.17	.16	.16	
Dunkirk silt loam, New York		2		.48	.57	.48		304
	(PK)	1	1.63	.21	.06	.19		
	(NPK)	1	1.14	.15	.04	.14		19x
	(NPKMn)	9		.24	.09			
Bridgethampton very fine sandy loam, Rhode Island.								
Summary:								
Maximum			2.89	1.02	.38	.60	.32	
Minimum			.79	.04	.03	.08	.07	
Mean			1.79(63)	.39(98)	.14(68)	.21(128)	.14(21)	
Timothy, leaves, cut for hay, maturity unknown:								
Grown in Wales		1	1.31		.19			170
Do		1	.27		.19			
Tobacco, leaves; dark, air-cured type; grown in Kentucky, summary:								
Maximum			3.41	3.52	1.01	.41	1.19	
Minimum			.51	2.91	.93	.21	.40	
Mean			2.31(33)	3.23(5)	.97(4)	.32(9)	.70(21)	
Tobacco, leaves, Burley: summary:								
Maximum			6.16	5.30	.80	.53		
Minimum			1.45	2.59	.25	.19		
Mean			3.56(72)	4.02(18)	.62(14)	.37(18)	.22(1)	

TABLE 20.—The principal chemical elements in some important crops—Continued

Plant or plant part, soil type, and location	Fertilizer used	Analyses	K	Ca	Mg	P	S	Reference No.
		Number	Percent	Percent	Percent	Percent	Percent	
Turnip, roots:								
Cecil series, Alabama		1		0.16		.06		13x
Decatur series, Alabama		3		.28		.08		
Norfolk series, Alabama		2		.23		.07		
Alton stony loam, Rhode Island	(NK)	2				.32		
	(NPK)	2				.34		
	(M)	2				.41		
	(NK)	3				.42		
	(NPKL)	3				.31		
	(None)	1				.22		
	(NK)	10				.28		
	(NP)	3				.35		
	(PK)	3				.36		
	(NPK)	8				.36		
	(NKL)	3				.33		
	(NPKL)	6				.39		
	(M)	3				.49		
	(NKL)	8				.26		
Bridgchampton very fine sandy loam, Rhode Island.								19x
Summary:								
Maximum			4.70	1.04	0.42	.79	0.78	
Minimum			1.07	.11	.11	.06	.12	
Mean			2.77 (97)	.51 (87)	.24 (48)	.36 (238)	.43 (5)	
Standard deviation			±.75	±.22		±.15		
Mode			2.00-3.00	.60-.64				
Turnip-tops, edible:								
Bridgchampton very fine sandy loam, Rhode Island.	NKL	7				.22		
Summary:								
Maximum			4.84	4.75	.95	.82	.86	
Minimum			1.27	1.49	.63	.14	.19	
Mean			3.00 (11)	3.28 (14)	.80 (4)	.39 (22)	.27 (2)	
Turnip, whole plant:								
Grown in France.		1		3.00		.848	1.494	66
Grown in Pennsylvania.		8		.69		.7x	.369	7x
Grown in Pennsylvania.		1	4.34			.32		291
Vetch, above-ground portions, cut before bloom; variety unknown; grown in Africa.								
Vetch, above-ground portion, full bloom; variety unknown; summary.		3	2.51	1.48	.43	.33	.28	
Vetch, above-ground portion, cut when pods were forming; grown in Germany.		1	1.70	1.51	.78	.62	.36	368
Vetch, above-ground portion, seeds beginning to ripen; variety unknown; grown in Germany.		1	1.48	1.40	.44	.31	.30	368
Vetch, purple (<i>Vicia arvensis</i>), above-ground portion, cut for hay; grown in South Carolina.		1		.53	.18	.38		414

	2	2.49	1.33	.36	
Vetch, common (<i>Vicia sativa</i> L.) above-ground portion, cut for hay; grown in Massachusetts; summary.					
Vetch, hairy (<i>Vicia villosa</i> Roth) above-ground portion, cut for hay; summary:					
Maximum		3.23	2.05	.68	
Minimum		1.38	.92	.22	
Mean		2.23(8)	1.28(28)	.36(36)	
Vetch, above-ground portion, cut for hay; variety unknown; summary:					
Maximum		2.99	3.39	.86	.20
Minimum		.81	.58	.07	.05
Mean		1.67(28)	1.14(36)	.29(32)	.14
Vetch, straw, cut when seeds were beginning to ripen; grown in Germany.					
Vetch, seed; grown in Germany; summary	1	1.44	1.54	.17	.14
Wheat, grain:	3	1.11	.12	.53	.14
Tilsit silt loam, Kentucky	1	.54	.068	.34	.19
Summary:	1	.52	.049	.35	.21
Maximum	1	.45	.057	.32	.21
Minimum	1	.48	.047	.44	.20
Mean	2	.62	.122	.54	.29
Standard deviation		.29	.005	.15	.10
Mode		.48(264)	.050(290)	.40(310)	.18(138)
Wheat, straw:					
Tilsit silt loam, Kentucky	1	.75	.19	.09	.26
Summary:	1	.98	.14	.27	.27
Maximum	1	1.29	.18	.06	.26
Minimum	2	.94	.18	.07	.28
Mean		1.54	.43	.17	.30
Standard deviation		.85(109)	.23(115)	.08(115)	.17(48)
Mode		±.24	±.07	±.03	±.03
Wheat, above-ground portion, in flower; grown in France.	17	.85-.92	.24-.26	.08-.09	.20
Wheat, above-ground portion, grown in France.	17				.22
Wheat, above-ground portion, milk stage; grown in Washington.	10	.21	.11	.15	.22
					533, 21x

TABLE 21.—The minor-element content of some important crops¹

[Moisture-free basis]

ARSENIC²

Plant or part of plant	Location	Reference No. ²	Arsenic	Remarks
Alfalfa, above-ground portion, cut for hay	France	300	Mg./kg. 0.5	
Alfalfa, above-ground portion	Virginia	594	14.	Hagerstown silt loam. Sample taken under trees sprayed with arsenic. Soil contained 60 p. p. m. of arsenic in first 3 inches.
Almond, edible portion	France	298	3	
Apple, edible portion	Switzerland	178	.19	
Do	do	178	.07	
Apple, fruit	France	298	.86	
Asparagus, edible portion	do	298	.75	
Banana, edible portion	do	298	.33	
Barley, grain	do	298	.55	
Do	do	298	1	
Beans, garden; edible pods and seeds	Kansas	594	1	
Do	Virginia	594	4	Soil contained 8 p. p. m. of arsenic.
Beets, leaves	France	298	.20	Chester clay loam. Contained 6 p. p. m. of arsenic. Two samples.
Beets, roots	do	300	.61	
Beets, tops	Virginia	594	1.3	Chester clay loam.
Do	do	594	.2	Do.
Do	New York	594	10.	Soil contained 140 p. p. m. of arsenic. Contaminated by spray.
Cabbage, edible portion	France	298	1.3	
Carrots, roots	do	298	.40	
Do	Virginia	594	.8	
Cauliflower, head	France	298	.86	Chester clay loam.
Celery, entire plant except flower	do	298	2.32	
Celery, stalks	Virginia	594	0.6	
Chestnuts, edible portion	France	300	.11	
Clover, red, above-ground portion	do	300	.37	
Clover, leaves	Virginia	594	12.	
Corn, grain	France	300	.86	
Do	Switzerland	178	.03	
Corn, sweet, edible portion	Virginia	594	.4	
Cress, water, edible portion	France	298	2.10	Chester clay loam.
Endive, above-ground portion	Switzerland	178	.21	
Hazelnuts, edible portion	France	298	.11	
Lettuce, edible portion	do	298	3.87	
Do	Switzerland	178	.43	
				<i>Trifolium pratense.</i> Hagerstown silt loam. Sample collected under trees sprayed with arsenic preparation.

TABLE 21.—The minor-element content of some important crops—Continued

BARIUM 3

Plant or part of plant	Location	Reference No.	Barium	Remarks
			<i>Mg./kg.</i>	
Alfalfa, above-ground portion, cut for hay	Kentucky	369	78	
Alfalfa, above-ground portion, mature	Missouri	495	130	Tatum silt loam. Contained 1,460 p. p. m. of barium.
Alfalfa, above-ground portion	Virginia	495	1,300	
Alfalfa, above-ground portion, before bloom	Pennsylvania	495	18	Hagerstown loam. Contained 540 p. p. m. of barium.
Alfalfa, above-ground portion, full bloom	do	495	70	Penn silt loam, Contained 470 p. p. m. of barium.
Apple, fruit	New Hampshire	495	3	Gloucester stony loam. Contained 4,750 p. p. m. of barium.
Beans, garden; above-ground portion, pods beginning to form	North Carolina	495	170	Cecil clay. Contained 350 p. p. m. of barium.
Beans, garden; edible green seeds	New Hampshire	495	3	Gloucester stony loam.
Cabbage, head	North Carolina	495	27	Cecil clay.
Carrots, roots	New Hampshire	495	27	Gloucester stony loam.
Clover, crimson; above-ground portion, full bloom	North Carolina	495	107	Cecil sandy loam. Contained 514 p. p. m. of barium.
Clover, red; above-ground portion, in bloom	Pennsylvania	495	74	Penn silt loam.
Do	do	495	54	Hagerstown loam.
Clover, red, above-ground portion, mature	New Hampshire	495	54	Gloucester stony loam.
Clover, sweet	do	495	13	Hagerstown loam.
Clover, white; above-ground portion, beginning to bloom	Pennsylvania	495	72	
Clover, white, above-ground portion, seeded	Maine	495	63	
Corn, above-ground portion; young	North Carolina	495	63	Norfolk sandy loam. Contained 36 p. p. m. of barium.
Do	South Carolina	495	18	York silt loam. Contained 2,570 p. p. m. of barium.
Corn, grain	Missouri	495	9	Decatur soil. Contained 1,790 p. p. m. of barium.
Corn, stover	Kentucky	369	82	
Cotton, plant, before bloom	North Carolina	495	36	Norfolk sandy loam.
Hemp, above-ground portion	Kentucky	369	21	
Kentucky bluegrass, above-ground portion, in seed	Pennsylvania	495	36	Penn silt loam.
Kentucky bluegrass, above-ground portion, stage of maturity unknown	Kentucky	369	46	
Lettuce, beginning to seed	Pennsylvania	495	72	Do.
Oats, above-ground portion, milk stage	do	495	18	Do.
Onions, bulb	New Hampshire	495	18	Gloucester stony loam.
Orchard grass	Pennsylvania	495	72	Penn silt loam.
Peas, above-ground portion, in blossom	North Carolina	495	63	Cecil clay.

Potatoes, tubers.....	495	New Hampshire.....	5	Gloucester stony loam.
Soybean, seed, mature.....	369	Kentucky.....	8	
Soybean, above-ground portion, cut for hay.....	369	do.....	29	
Soybean, stems and leaves.....	496	Missouri.....	1,090	Decatur soil.
Sweetclover, above-ground portion, before bloom.....	495	Pennsylvania.....	27	Hagerstown loam.
Timothy, above-ground portion, in bloom.....	495	do.....	36	Penn silt loam.
Do.....	495	do.....	27	Hagerstown loam.
Timothy, above-ground portion, in seed.....	495	New Hampshire.....	45	Gloucester stony loam.
Tobacco, above-ground portion, before bloom.....	495	Maine.....	18	
	495	North Carolina.....	160	Durham sandy loam. Contained 1,002 p. p. m. of barium.
Tobacco, dark, air-cured type; leaves.....	369	Kentucky.....	293	5 samples; range 56 to 435.
Tobacco, burley, leaves.....	369	do.....	88	2 samples.
Tobacco, burley, stalks.....	369	do.....	193	4 samples; range 53 to 400.
Tobacco, dark, air-cured type; stalks.....	369	do.....	273	4 samples; range 224 to 400.
Wheat, above-ground portion, mature.....	495	North Carolina.....	18	Durham sandy loam.
Do.....	495	Pennsylvania.....	18	Penn silt loam.
Wheat, grain.....	496	Missouri.....	8	Decatur soil.
Wheat, leaves.....	496	do.....	130	Do.
Wheat, stems.....	496	do.....	180	Do.

³ The barium content of locoweed and other weeds and grasses has been reported (41, 396). Barium in Brazil nuts (517, 578) and in tobacco and other plants (16) has been determined, but the moisture content of the materials was not reported.

TABLE 21.—The minor-element content of some important crops—Continued

BORON 4

Plant or part of plant	Location	Reference No.	Analyses	Boron			Remarks
				Maximum	Minimum	Mean	
Alfalfa, above-ground portion, cut for hay	France	66	Number 2	Mg./kg. 29	Mg./kg. 27	Mg./kg. 27	
Alfalfa, leaves	Washington and Canada	387	11	32	4	15	
Almond, edible portion	California	157	1			26	
Apples, edible portion	Nova Scotia	303	3		12	13	
Apple, fruit, immature	New Zealand	21	26	44	4	4	
Apple, fruit, mature			128	76	3	19	
Apple, parings	Nova Scotia	303	3	15	15	15	
Apricot, fruit	California	157	2	52	51	51	
Barley, grain	France	66	1			2	
Beans, garden; edible pods and seeds	France	66	7	37	2	17	
Cabbage, edible portion	do	66	1			37	
Carrots, whole plant	do	66	1			25	
Celery, entire plant except flower	do	66	3	15	.05	9	
Cherries, edible portion	California	157	1			42	
Clover, crimson; above-ground portion	France	68	1			70	<i>Trifolium pratense.</i>
Clover, red; above-ground portion	do	66	1			36	Do.
Clover, red; flower	Russia	78	1			40	Do.
Clover, red; leaves	do	78	1			57	Do.
Clover, red; stems	do	78	1			28	<i>T. incarnatum.</i>
Coffee, bean	Trinidad	156	1			17	
Cowpeas, above-ground portion	Florida	120	3			190	
Cranberries, edible portion	California	157	2	283	42	82	
Currants, edible portion	do	157	5	96	68	37	
Dandelion, whole plant	France	68	1	58	19	80	
Dates, edible portion	California	157	1			14	
Figs	do	157	1			80	
Gooseberries, edible portion	do	157	1			49	
Lupine, leaves	Russia	78	1			9	
Lupine, stems	do	78	1			6	
Lupine, entire flower	do	78	1			39	
Mustard, greens	France	68	2	53	22	38	
Orange, edible portion	California	157	4	38	10	20	
Peach, fruit	do	157	2	52	49	51	

Pear, fruit	do	157	1	7	2	19
Peas, green; edible portion	Netherlands	357	8	22	18	3
Peas, above-ground portion	Virginia	120	5	16	2	20
Potatoes, tubers	France	66	2	15	14	7
Potatoes, tops	California	157	3	7	7	14
Prunes, fruit	Florida	120	3	152	17	79
Radish, roots	New Zealand	488	4	11	7	7
Rutabaga, roots	do		2	3	3	9
Rye, above-ground portion, maturity unknown	Virginia	120	5	29	6	19
Soybean, seeds, mature	do	120	5	13	1	8
Soybean, above-ground portion	France	68	1			10
Spinach, edible portion	Russia	78	1			51
Tobacco, leaves	France	66	1			25
Tobacco, above-ground portion	Netherlands	157	1			191
Tomato, fruit	New Zealand	488	7	8	6	7
Turnip, roots	France	68	1			49
Turnip, whole plant	do	66	1			3
Wheat, above-ground portion						

⁴ The analyses of Askew and Thomson (28) and of de Long (147) showing the relation of the boron content of apples to internal cork have been omitted because they are given on the fresh basis without a statement as to the moisture content.

Summary (68, 553).

Summary (66, 68).

TABLE 21.—The minor-element content of some important crops—Continued

BROMINE ⁵

Plant or part of plant	Location	Reference No.	Bromine Mg./kg.	Remarks
Alfalfa, above-ground portion, cut for hay	France	130	6.4	
Do	do	130	1.9	
Do	do	131	9.8	
Artichoke, flower head	France	131	6.1	
Do	do	131	2.8	
Artichoke, Jerusalem	do	131	20.2	
Artichoke, Japanese; tuber	do	131	5.4	
Artichoke, Japanese; tuber	do	131	5.5	
Artichoke, edible portion	do	130	6.4	
Asparagus, edible portion	do	131	5.5	
Banana, edible portion	do	131	3.7	
Barley, grain	do	131	4.5	
Beans, garden; edible pods and seeds	do	131	94.5	
Beets, roots	do	131	3.9	
Do	do	131	7.3	
Cabbage, edible portion	do	131	6.7	
Cantaloup, edible portion	do	131	4.7	
Carrots, roots	do	131	3.8	
Cauliflower, head	do	131	Tr.	
Do	do	130	1.7	
Celery, entire plant except flower	do	131	1.7	
Do	do	131	.9	
Do	do	131	1.8	
Do	do	131	4.4	
Cherries, edible portion	do	131	2.0	
Corn, grain	do	131	2.2	
Currants, edible portion	do	130	6.2	
Do	do	131	5.3	
Figs	do	131	3.8	
Garlic, bulb	do	131	3.1	
Grapes, edible portion	do	131		
Hemp, plant	do	131		
Jerusalem-artichoke, tuber	do	130		
Mandarin, edible portion	do	131		
Millet, grain	do	131		
Oats, grain	do	130		
	do		3.1	

Onion, bulb.....	do	131	2.2
Do.....	do	131	1.0
Orange, edible portion.....	do	131	3.2
Peach, edible portion.....	do	131	4.7
Do.....	do	131	T _r
Peas, green; edible portion.....	do	130	2.1
Peas, pods only.....	do	130	6.3
Potatoes, tubers.....	do	131	14.3
Do.....	do	131	2.7
Radish, roots.....	do	131	9.2
Do.....	do	131	8.3
Rhubarb, stem.....	do	131	7.5
Rye, grain.....	do	130	1.9
Strawberries, edible portion.....	do	131	7.1
Turmp, edible portion.....	do	131	8.9
Do.....	do	131	3.1
Vetch, above-ground portion.....	do	130	2.1
Watermelon, edible portion.....	do	131	262

^a Analyses of bromine in wheat (286) and in several fruits and vegetables (159) have been reported, but the moisture content of the samples used is not given.

TABLE 21.—The minor-element content of some important crops—Continued

CHLORINE ⁶

Plant or part of plant	Location	Reference No.	Analyses	Chlorine			Remarks
				Maximum	Minimum	Mean	
Alfalfa, above-ground portion, prebud stage	England	601, 602	Number	Percent	Percent	Percent	
Alfalfa, above-ground portion, in bud	do	600, 601, 602	17	0.66	0.20	0.43	
			19	.56	.09	.34	
Alfalfa, above-ground portion, one-tenth to one-quarter bloom	New Hampshire	495	18	.71	.12	.35	Summary.
Alfalfa, above-ground portion, one-half bloom			3	.90	.30	.51	Do.
Alfalfa, above-ground portion, full bloom			9	.93	.25	.49	Do.
Alfalfa, above-ground portion, cut for hay			33	1.74	.01	.88	Do.
Apples, fruit			1			.63	Gloucester stony loam.
Do			2	.96	.03	.04	Summary.
Artichoke, flower head	France	131	1			.38	
Apricot, edible portion	do	131	1			.09	
Barley, grain			29	.60	.05	.17	Summary.
Beans, field; seed			14	.10	.02	.05	Summary.
Beans, garden; edible pods and seeds			17	.50	.05	.09	Summary.
Beans, garden; edible hulls	Kentucky	379	1			.05	
Beans, garden; immature seed	do	379	1			.06	
Beans, lima, mature seed	do	379	1			.03	
Beets, roots			19	.71	.11	.22	Do.
Broccoli, edible portion	Maryland	140	3	.89	.78	.85	
Cabbage, edible portion	Florida	5X	1			.46	Coxville soil.
Do	do	5X	1			.44	Leon soil (NPK).
Do	do	5X	8	.60	.39	.44	Leon soil (NPKMn).
Do	do	5X	2			.42	Leon soil (NPKCd).
Do	do	5X	2			.42	Parkwood soil (NPK).
Do	do	5X	1			.74	Ceal clay.
Do	do	5X	1			.85	Summary.
Do	do	495	52	1.50	.17	.51	Do.
Carrots, roots			6	1.50	.23	.52	
Cauliflower, head	France	131	2	.57	.57	.57	
Cauliflower, edible portion	California	140	2	.50	.47	.48	
Celery, edible portion	Florida	5X	3			2.28	Bladen soil.
Do			34	5.28	.12	1.92	Summary.
Cherries, edible portion	France	131	1			lr.	
Clover, alsike; above-ground portion			14	1.40	.15	.78	Summary. <i>Trifolium hybridum</i> .
Clover, crimson; above-ground portion	North Carolina	495	1			.37	<i>T. incarnatum</i> . Cecil sandy loam.
Do			11	1.19	.07	.63	Summary. <i>T. incarnatum</i> .
Clover, red; above-ground portion, mature, with blossoms and seeds.	New Hampshire	495	1			.30	Gloucester stony loam.
Clover, red; above-ground portion, in bloom	Pennsylvania	495	1			.20	<i>T. pratense</i> . Hagerstown loam.

TABLE 21.—The minor-element content of some important crops—Continued

CHLORINE—Continued

Plant or part of plant	Location	Reference No.	Analyses	Chlorine			Remarks
				Maximum	Minimum	Mean	
				Percent	Percent	Percent	
Potatoes, tubers.....	Florida.....	5x	3	Coxville soil (NPKMn),
Do.....	do.....	5x	2	Coxville soil (NPKCu),
Do.....	do.....	5x	1	Coxville soil (NPKMnCu),
Do.....	do.....	5x	4	0.43	0.20	Leon soil (NPK),
Do.....	do.....	5x	1	Norfolk soil (NPK),
Do.....	New Hampshire	5x	1	Norfolk soil (NPK),
Do.....	Do.....	5x	1	Gloucester stony loam.
Potatoes, tops.....	Germany.....	137, 368	496	.81	.03	Summary.
Prunes, edible portion.....	Germany.....	137, 192	16	3.36	.26	2.27
Redtop, above-ground portion, cut for hay.....	Wyoming.....	128	2	.09	.08	Summary.
Rye, grain.....	Do.....	4	.03	.02	Do.
Rye, straw.....	Do.....	25	.52	.06
Soybean, leaves.....	France.....	335	1
Soybean, pods.....	do.....	335	1
Soybean, stems.....	do.....	335	1
Soybean, seed, mature.....	Ohio.....	192	2	.04	.03
Spinach, edible portion.....	Maryland.....	140	2	.93	.66	Sassafras sandy loam.
Do.....	Virginia.....	140	6	.58	.52	Norfolk loam.
Do.....	Do.....	6	1.94	.52	Summary.
Do.....	France.....	131	1
Straw berries, edible portion.....	France.....	8	.92	.06	Summary.
Sugar beets, roots.....	Colorado.....	606	5	4.68	1.47	2.86
Sugar beets, young leaves.....	Germany.....	584	14	.33	.08
Sugar beets, leaves, mature.....	Africa.....	81	1
Sugarcane, leaves.....	Do.....	14	.57	.01	Summary.
Sugarcane, canes.....	District of Columbia	592	1
Sunflower, heads minus seeds and husks.....	do.....	592	1
Sunflower, seed husk.....	do.....	592	1
Sunflower, seed kernel without husk.....	do.....	592	1
Sunflower, leaves.....	do.....	592	1
Sunflower, stalks.....	do.....	592	1
Sunflower, above-ground portion, cut before bloom.....	Germany.....	308	1
Sunflower, above-ground portion, full bloom.....	do.....	308	1
Sunflower, above-ground portion, cut for forage.....	do.....	308	3	.05	.04
Sunflower, leaves and stems.....	do.....	308	3	.33	.22
Sunflower, seeds with husks.....	District of Columbia	592	1
Sweetclover, above-ground portion.....	Pennsylvania	495	1
Sweetpotatoes, tuber.....	Ohio.....	192	1	<i>Metilotus alba.</i>

TABLE 21.—The minor-element content of some important crops—Continued

COBALT

Plant or part of plant	Location	Reference No.	Cobalt	Remarks
Apricot, edible portion.....	France.....	61	Mg./kg. 0.032	
Beans, field, seed.....	do.....	61	.011	
Buckwheat, grain.....	do.....	61	.36	
Cabbage, edible portion.....	do.....	60	.07	
Carrots, roots.....	do.....	61	.02	
Carrots, leaves.....	do.....	61	.31	
Cherries, edible portion.....	do.....	61	.065	
Clover, white, above ground-portion.....	New Zealand.....	23	4.6	
Do.....	do.....	23	.28	
Coffee, bean.....	France.....	61	.002	
Corn, grain.....	do.....	61	.011	
Cress, water.....	do.....	61	.15	
Figs.....	do.....	61	.20	
Oats, grain.....	do.....	61	Tr.	
Onions, bulb.....	do.....	61	.13	
Pear, pulp.....	do.....	61	.18	
Peas, green; edible portion.....	do.....	61	.03	
Potatoes, tubers.....	do.....	61	.06	
Rice, polished.....	do.....	61	.006	
Spinach, edible portion.....	do.....	61	.074	
Tomato, fruit.....	do.....	61	.096	
Walnuts, edible portion.....	do.....	61	.05	
Wheat, grain.....	do.....	61	.012	
				<i>Phaseolus vulgaris.</i>
				<i>Trifolium repens.</i>
				Do.

TABLE 21.—The minor-element content of some important crops—Continued
COPPER 7

Plant or part of plant	Location	Reference No.	Analyses	Copper			Remarks
				Maximum	Minimum	Mean	
Alfalfa, above-ground portion, cut for hay			Number	Mg./kg.	Mg./kg.	Mg./kg.	
Apples, edible portion			8	15	4	9	Summary (166, 373, 14x).
Apples, fruit, mature			2	13	11	12	Summary (232, 355).
Artichoke, flower head	Wisconsin		3	7	5	6	Summary (232, 355).
Asparagus, edible portion		355	1	17	7	11	Summary (185, 347, 355).
Banana, edible portion			3	9	8	9	Summary (168, 232, 355).
Barley, grain			12	41	6	16	Summary (166, 230, 232).
Beans, field; seed			12	16	7	11	<i>Phaseolus vulgaris</i> . Summary (355, 379, 503).
Beans, garden; edible pod and seeds	Florida	5x	4	20	10	16	St. Lucie soil.
Do.			24	20	6	13	Summary (232, 348, 482, 5x).
Beans, garden; edible hulls	Kentucky	379	1			12	
Beans, garden; immature seed	do		1			10	
Beans, horse; seed	France	232, 392	3	11	10	10	<i>Vicia faba</i> .
Beans, lima; mature seed			3	10	8	9	Summary (355, 379, 503).
Beets, root			15	27	6	10	Summary (129, 232, 355, 482, 8x)
Beets, leaves		355	4	18	9	12	Summary (355, 482).
Blackberries, fruit	Wisconsin	355	1			10	
Blueberries, fruit	do	355	1			6	
Brazil nut, edible portion		355	1			15	
Broccoli, edible portion	Wisconsin	355	3	37	2	24	Summary (140, 347).
Brussels sprouts, edible portion	do		1			8	
Butternut, edible portion	Florida	355	1			15	
Cabbage, edible portion		5x	8	28	19	22	Coxville soil (NPK).
Do.	do	5x	2			13	Leon soil (NPK).
Do.	do	5x	2			26	Leon soil (NPKMn).
Do.	do	5x	2			10	Leon soil (NPKCu).
Do.	do	5x	1			14	Parkwood soil (NPK).
Do.	do	5x	26	28	4	14	Summary (185, 355, 482, 5x)
Cabbage, outer green leaf	England	129	2	5	5	5	
Cabbage, inner green leaf	do	129	2	6	4	5	
Cabbage, center vein of outer leaves	do	129	1			3	
Cabbage, center vein of inner leaves	do	129	1			2	
Cabbage, center stalk of plant	do	129	1			3	
Cabbage, outermost leaves, mesophyll	Japan	502	1	5	5	3	<i>Brassica pekinensis</i> (B. pe- <i>tsai</i>).
Cabbage, leaves, second layer, veins	do	502	6	4	4	4	Do.
Cabbage, leaves, second layer, mesophyll	do	502	6	4	4	3	Do.
Cabbage, leaves, second layer, veins	do	502	6	3	3	2	Do.
Cabbage, leaves, second layer, mesophyll	do	502	6	4	4	4	Do.

⁷ The analyses of Mader and Mader (390) of the copper content of potatoes and those of Grendel (287) and of van Leeuwen (312) were omitted because the moisture content of the material was not given. The data of Vedrode (573) have been omitted because of the very high values reported. The work was done in 1896.

TABLE 21.—The minor-element content of some important crops—Continued
COPPER

Plant or part of plant	Location	Reference No.	Analyses	Copper			Remarks
				Maximum	Minimum	Mean	
Cabbage, leaves, third layer, veins	Japan	502	Number	Mg./kg.	Mg./kg.	<i>Brassica pekinensis</i> (B. pe-tsai.) Do. Do.	
Cabbage, leaves, fourth layer, mesophyll	do.	502	5	3	3		
Cabbage, leaves, fourth layer, veins	do.	502	5	4	4		
Cabbage, center	do.	502	6	3	3	Do.	
Cantaloup, edible portion	do.	502	6	4	4		
Carrots, roots	Wisconsin	355	2	6	6	Summary (232, 355, 391, 482, 8x).	
Carrots, leafy part excepting lower leaves	do.	355	15	18	7		
Carrots, stalks of leaves	England	129	2	11	10		
Carrots, stalks of leaves	do.	129	2	11	10		
Carrots, yellow part of root	do.	129	1	—	5		
Carrots, red part of root	do.	129	1	—	5		
Cauliflower, edible portion	do.	129	1	—	5		
Celery, edible portion	do.	129	1	—	5		
Do.	Florida	5x	3	47	16		
Celery cabbage	do.	5x	3	560	2		
Cherries, edible portion	Wisconsin	355	1	—	10		Summary (232, 355, 5x).
Chestnut, edible portion	do.	355	1	—	10		
Citron, edible portion	Puerto Rico	179	2	16	12	Summary (232, 355).	
Citron, edible portion	do.	179	2	9	5		
Clover, alsike; above ground portion	Kentucky	14x	1	7	5	<i>Citrus medica</i> <i>Trifolium hybridum</i>	
Clover, red; above ground portion	do.	14x	5	11	9		
Do.	do.	14x	41	20	6	<i>T. pratense</i> . Tilsit silt loam. <i>T. pratense</i> . (166, 14x).	
Clover, red; seeds	do.	14x	1	—	17		
Clover, red; foliage	do.	14x	1	—	8	Do.	
Coconut, edible portion	Wisconsin	14x	1	—	11		
Corn, grain	do.	14x	6	—	8	Summary (166, 185, 232, 503).	
Corn, stover	Kentucky	14x	6	17	4		
Do.	do.	14x	2	6	3	Tilsit silt loam. Summary (166, 14x).	
Corn, sweet, edible portion	do.	14x	2	9	2		
Cotton, seed	Wisconsin	355	2	7	6	Norfolk soil.	
Cowpeas, seed	Mississippi	374	1	—	54		
Cranberries, edible portion	Alabama	13x	1	—	12	Summary (232, 355).	
Cress, water, edible portion	Wisconsin	355	2	—	12		
Cucumber, fruit	France	232	1	18	5	Summary (343, 355).	
Cucumber, edible portion	do.	232	1	30	18		
Currants, edible portion	Wisconsin	355	2	—	24	Summary (232, 355).	
Dandelion, edible portion	do.	355	1	—	17		
Dates, edible portion	do.	355	4	5	2	Summary (115, 232, 355).	
Eggplant, edible portion	do.	355	4	16	11		
Figs	Wisconsin	355	1	—	6	Summary (355, 482).	
Gooseberries	do.	355	1	—	8		
Grapefruit, edible portion	do.	355	1	—	5	Summary (232, 355, 503).	
Grapes, edible portion	do.	355	3	10	5		

Hickory nuts, edible portion.....	Wisconsin.....	355	1	-----	15
Kafir, grain.....	do.....	166	1	-----	8
Kale, edible portion.....	California.....	140	2	-----	24
Do.....	Maryland.....	140	2	56	31
Do.....	Virginia.....	140	2	39	35
Do.....	-----	140	6	56	24
Do.....	Kentucky.....	375	3	-----	14
Do.....	-----	-----	8	14	7
Kentucky bluegrass, above-ground portion, young blades.....	-----	-----	1	-----	60
Kentucky bluegrass, above ground portion, stage of maturity unknown.....	Kentucky.....	375	1	-----	15
Kentucky bluegrass, seeds.....	Wisconsin.....	355	1	-----	5
Kohlrabi, edible portion.....	-----	355	1	-----	5
Kumquat, edible portion.....	-----	236, 355	3	10	3
Lemon, fruit.....	California.....	236	9	2	3
Lemon, fruit, seed-free.....	do.....	236	2	4	3
Lemon, peel.....	do.....	236	1	-----	4
Lemon, pulp with seed.....	do.....	236	1	-----	5
Lettuce, head, outer leaves removed.....	Florida.....	5x	13	-----	32
Do.....	do.....	5x	3	83	26
Do.....	do.....	5x	2	-----	31
Do.....	do.....	5x	1	-----	31
Do.....	do.....	5x	1	-----	18
Do.....	do.....	5x	45	33	3
Lettuce, edible portion.....	-----	-----	-----	-----	-----
Lettuce, outer leaves.....	England.....	129	2	19	16
Lettuce, inner leaves.....	do.....	129	2	26	25
Lettuce, center stalk.....	do.....	129	1	-----	13
Lettuce, leaf.....	Wisconsin.....	355	1	-----	11
Mulberry, fruit.....	France.....	185	1	-----	4
Mushrooms, edible portion.....	Wisconsin.....	355	1	-----	62
Oats, grain.....	-----	-----	29	51	4
Oats, straw.....	-----	-----	26	54	3
Olives, fruit.....	France.....	232	1	-----	8
Okra, fruit.....	South Carolina.....	482	8	13	5
Onions, bulb.....	-----	-----	11	24	6
Orange, edible portion.....	California.....	236, 355	12	31	1
Orange, fruit.....	-----	-----	3	22	3
Parsley, edible portion.....	Wisconsin.....	355	1	-----	17
Parsnips, root.....	do.....	355	1	-----	7
Peanut, edible portion.....	do.....	355	1	-----	10
Pear, edible portion.....	-----	-----	2	13	6
Pear, green, edible portion.....	-----	-----	9	15	6
Peas, with pods, edible.....	France.....	391	2	12	7
Peas, pods only.....	-----	-----	2	5	3
Peas, black-eyed; seeds.....	North Carolina.....	503	2	-----	4
Pecans, edible portion.....	-----	-----	1	-----	7
Peppers, green; edible portion.....	Wisconsin.....	-----	1	-----	14
Pineapple, fruit.....	-----	-----	1	-----	16
Pistachio nut, edible portion.....	-----	-----	1	-----	8
Plums, edible portion.....	Wisconsin.....	-----	1	-----	12
Potatoes, tubers.....	Florida.....	5x	6	26	19
Do.....	do.....	5x	1	-----	22
Do.....	do.....	5x	1	-----	18
Do.....	do.....	5x	4	14	9
Do.....	do.....	-----	-----	-----	-----
Yolo silt loam.....	-----	-----	-----	-----	-----
Sassafras sandy loam.....	-----	-----	-----	-----	-----
Norfolk loam.....	-----	-----	-----	-----	-----
Summary.....	-----	-----	-----	-----	-----
Summary (166, 373, 381),.....	-----	-----	-----	-----	-----
Leon soil (NPK),.....	-----	-----	-----	-----	-----
Leon soil (NPKMn),.....	-----	-----	-----	-----	-----
Leon soil (NPKCu),.....	-----	-----	-----	-----	-----
Leon soil (NPK),.....	-----	-----	-----	-----	-----
Summary (343, 347, 355, 411, 482, 5x, 8x),.....	-----	-----	-----	-----	-----
Summary (232, 355, 482),.....	-----	-----	-----	-----	-----
Summary (232, 236),.....	-----	-----	-----	-----	-----
Summary (166, 220, 232, 473),.....	-----	-----	-----	-----	-----
Summary (166, 383, 473),.....	-----	-----	-----	-----	-----
Summary (232, 355),.....	-----	-----	-----	-----	-----
Summary (129, 185, 232, 355, 391, 392),.....	-----	-----	-----	-----	-----
Summary (129, 185),.....	-----	-----	-----	-----	-----
Bladen soil (NPK),.....	-----	-----	-----	-----	-----
Bladen soil (NPKMn),.....	-----	-----	-----	-----	-----
Bladen soil (NPKCu),.....	-----	-----	-----	-----	-----
Coxville soil (NPK),.....	-----	-----	-----	-----	-----

TABLE 21.—*The minor-element content of some important crops—Continued*
COPPER—Continued

Plant or part of plant	Location	Reference No.	Analyses	Copper			Remarks
				Maximum	Minimum	Mean	
Potatoes, tubers.	Florida	5x	Number	Mg./kg.	Mg./kg.		
Do.	do.	5x	3		8	Coxville soil (NPKMn).	
Do.	do.	5x	2		19	Coxville soil (NPKCu).	
Do.	do.	5x	1		13	Coxville soil (NPKMnCu).	
Do.	do.	5x	4		11	Leon soil (NPK).	
Do.	do.	5x	1	20	7	Norfolk soil (NPK).	
Do.	Wisconsin	12x	3	6	4	Miami silt loam.	
Do.	do.		143	24	1	Summary (129, 232, 343, 355, 389, 391, 503, 5x, 12x).	
Potatoes, tops.	England	129	1	1	11	Summary (391, 506).	
Prunes, edible portion.	Wisconsin	355	16	9	3		
Pumpkin, edible portion.	France	292	1		11		
Pumpkin, fruit.	Wisconsin	355	1		8		
Quince, edible portion.	do.	355	1		29		
Radish, roots.	Alabama	13x	1		Tr.	Norfolk soil.	
Rape, above-ground portion.	Wisconsin	166	1		8		
Do.	do.	355	1		8		
Raspberries, fruit.	Kentucky	14x	1		5	Dekalb silt loam; no fertilization.	
Redtop, above-ground portion, cut for hay.	do.	14x	1		4	Dekalb silt loam (M).	
Do.	do.	14x	1		4	Dekalb silt loam (ML).	
Do.	do.	14x	1		4	Dekalb silt loam (ML).	
Do.	do.	14x	1		4	Dekalb silt loam (PL).	
Do.	do.	14x	1		3	Tilist silt loam (PL).	
Do.	do.	14x	1		4	Tilist silt loam; no fertilization.	
Do.	do.	14x	1		3	Tilist silt loam (NP).	
Do.	do.	14x	1	5	3	Tilist silt loam (NPL).	
Rhubarb, edible portion.	Wisconsin	355	13		4		
Rice, brown.	do.	355	2		9	Summary (355, 380).	
Rice, polished.	do.	355	1	4	3	Summary (232, 355, 380).	
Rutabaga, roots.	Wisconsin	355	1	6	2	Summary (166, 232).	
Rye, grain.	do.		2	7	6		
Rye, straw.	Wisconsin	166	2		7		
Salsify, edible portion.	do.	355	1		4		
Soybean, leaves.	Kentucky	372	1		11		
Soybean, seeds, mature.	do.	372	1		8		
Soybean, seeds, green.	Wisconsin	166	1		12		
Soybean, above-ground portion, cut for hay.	Kentucky	14x	1		23		
Do.	do.	14x	1		9	Dekalb silt loam; no fertilization.	
Do.	do.	14x	2	10	9	Dekalb silt loam (ML).	
Do.	do.	14x	2	10	9	Dekalb silt loam (MP).	
Do.	do.	14x	2	10	9	Dekalb silt loam (MPPL).	
Do.	do.	14x	2	10	9	Dekalb silt loam (MPKL).	
Do.	do.	14x	1		8	Tilist silt loam; no fertilization.	

Do.....	do.....	14x	1	12	11	8	Tiltsit silt loam (M).
Do.....	do.....	14x	2	11		11	Tiltsit silt loam (MP).
Do.....	do.....	14x	3				Tiltsit silt loam (MPK).
Do.....	do.....	14x	1				Tiltsit silt loam (ML).
Do.....	do.....	14x	2		10	10	Tiltsit silt loam (MPL).
Do.....	do.....	14x	6		6	6	Tiltsit silt loam (MPKL).
Do.....	do.....	14x	32	7	4	9	Summary (166, 14x).
Spinach, edible portion.....	Maryland.....	140	2	45	43	44	Sassafras sandy loam.
Do.....	Virginia.....	140	2	73	66	69	Norfolk loam.
Do.....	do.....		40	73	3	15	Summary (140, 232, 347, 355, 392, 411, 482, 8x).
Do.....	do.....		34	24	3	9	Summary eliminating results of Davidson and LeClerc (140) (232, 347, 355, 392, 411, 482, 8x).
Squash, edible portion.....	South Carolina.....	482	6	15	11	13	Summary (185, 355).
Straw berries, edible portion.....	do.....		2	8	2	8	
Sugar beets, roots.....	Wisconsin.....	166	1			6	
Sugar beets, leaves.....	do.....	166	1			6	
Sweetclover, above-ground portion.....	do.....		2	12	9	10	<i>Melilotus alba</i> . Summary (166, 14x).
Sweetpotatoes, tubers.....	Wisconsin.....	355	11	9	3	4	Summary (482, 503).
Swiss chard, edible portion.....	do.....	355	1			13	
Tangerine, edible portion.....	do.....	355	1			6	
Timothy, above-ground portion, cut for hay.....	Kentucky.....	14x	4	7	4	5	Tiltsit silt loam.
Do.....	do.....		14	7	2	5	Summary (166, 14x).
Tobacco, leaves.....	Wisconsin.....	166	1			17	
Tomato, fruit.....	Florida.....	5x	26	19	5	11	Parkwood sandy loam (NPK).
Do.....	do.....		51	34	8	14	Summary (166, 232, 355, 378, 482, 5x).
Tomato, juice and pulp.....	Italy.....	349	7	9	5	7	
Tomato, seeds.....	do.....	349	7	12	9	10	
Tomato, skins.....	do.....	349	7	12	8	10	
Turnip, roots.....	do.....		8	18	4	9	Summary (129, 232, 355, 482).
Turnip, leaves.....	do.....		7	12	6	8	Summary (129, 347, 482).
Vetch, above-ground portion, cut for hay.....	Wisconsin.....	166	7			10	
Watermelon, edible portion.....	do.....	355	1			9	Tiltsit silt loam; no fertilization.
Wheat, grain.....	Kentucky.....	14x	1			6	Tiltsit silt loam (M).
Do.....	do.....	14x	1			8	Tiltsit silt loam (ML).
Do.....	do.....	14x	1			6	Tiltsit silt loam (MPL).
Do.....	do.....	14x	1	5	5	9	Summary (220, 232, 381, 391, 543, 581, 1x, 14x).
Do.....	do.....	14x	108	24	4	3	Tiltsit silt loam; no fertilization.
Wheat, straw.....	Kentucky.....	14x	1			4	Tiltsit silt loam (M).
Do.....	do.....	14x	1			3	Tiltsit silt loam (ML).
Do.....	do.....	14x	2	3	2	3	Tiltsit silt loam (MPL).
Do.....	do.....	14x	2	3	2	3	Summary (166, 185, 381, 14x).
Do.....	do.....	14x	24	5	1	3	

TABLE 21.—The minor-element content of some important crops—Continued

FLUORINE ⁸

Plant or part of plant	Refer- ence No.	Location	Fluorine Mg./kg.	Plant of part of plant	Refer- ence No.	Location	Fluorine Mg./kg.
Alfalfa, above-ground portion.....	203	France	56.5	Lentil.....	203	France	18.0
Apple, pulp.....	203	do	2.1	Lettuce.....	404	Austria	1.2
Apple, skin.....	203	do	27.8	Mustard, black; seeds.....	203	France	15.8
Apricot, edible portion.....	203	do	25.0	Mustard, black; leaves.....	203	do	68.0
Asparagus, young shoot.....	203	do	79.4	Onions, bulb.....	404	Austria	3.0
Banana, edible portion.....	203	do	3.8	Peach, pulp.....	203	France	39.3
Beans, garden; edible pods and seeds.....	404	Austria	6	Pear, pulp.....	203	do	1.7
Beets, leaves.....	203	France	134	Potatoes, tuber.....	203	do	3.0
Buckwheat.....	203	do	25.3	Radish, root.....	203	do	20.0
Cabbage, head.....	203	do	10.8	Rice, polished.....	203	do	9.4
Carrots, root.....	203	do	3.4	Rice, unpolished.....	203	do	30.0
Cauliflower, edible portion.....	203	do	25.7	Spinach, leaves.....	203	do	1.7
Cherries, pulp and skin.....	203	do	37.0	Do.....	404	Austria	1.3
Cress.....	203	do	12.0	Strawberries.....	203	France	14.0
Figs.....	203	do	19.8	Tomato, fruit.....	203	do	40.6
Grapes, edible portion.....	203	do	8.1	Tomato, edible portion.....	404	Austria	None
Kidney beans, mature seed.....	203	do	21.0	Turnip.....	203	France	20.2
Kidney beans, green seed.....	203	do	2.1	Walnuts, edible portion.....	203	do	7.8

⁸ The fluorine analyses quoted by Steinkoenig (592) are those of Gautier and Clausman (203) compiled in this table.

Plant or part of plant	Location	Reference No.	Analyses	Iodine (parts per billion)			Remarks
				Maximum	Minimum	Mean	
Alfalfa, above-ground portion, cut for hay.			Number				
Apple, fruit.			4	385	69	157	Summary (5, 414).
Asparagus, edible portion.	Oregon.	363	4	340	89	3	Summary (5, 45).
Barley, grain.			7	3,780	12	1,168	Summary (153, 258, 365, 481).
Beans, field; seed.	Germany.	45	5	124	73	550	Summary (45, 363).
Beans, garden; edible pods and seeds.	South Africa.	77	4	740	286	57	<i>Phaseolus vulgaris</i> .
Beets, roots.			32	1,560	29	579	Summary (5, 45, 258, 363, 385, 404, 587, 7x).
Beets, leaves.			18	416	8	241	Summary (45, 258, 264, 481, 587).
Cabbage, edible portion.			8	1,435	248	984	Summary (45, 258).
Carrots, roots.			85	2,400	16	218	Summary (5, 45, 258, 364, 445, 481, 587, 604).
			35		2	309	Summary (5, 46, 258, 264, 363, 365, 445, 481, 587, 604).
Celery, edible portion.			2	623	14	318	Summary (365, 587).
Cherries, edible portion.	Oregon.	363	1			33	<i>Citrus medica</i> L.
Citron, edible portion.	Puerto Rico.	179	1			26	<i>Trifolium hybridum</i> .
Clover, alsike; above-ground portion.	South Carolina.	414	1			181	Summary (381, 414).
Clover, red; above-ground portion.			23			121	<i>T. pratense</i> .
Clover, white; above-ground portion.	New Zealand.	525	1			5,590	<i>T. repens</i> .
Collards, edible portion.	Georgia.	274	9	191	34	74	
Corn, grain ⁹ .			21	650	63	177	Summary (377, 548).
Corn, sweet, milk stage.	Pennsylvania.	7x	10	210	100	130	Hagerstown soil.
Corn, sweet, edible portion.			21	250	52	127	Summary (7x, 363, 385).
Cowpeas, above-ground portion.	South Carolina.	414	7	560	37	167	
Cranberries, edible portion.	Massachusetts.	416, 417	1			259	
Cress, water, edible portion.	England.	445	1			626	
Cucumber, edible portion.			8	940	44	226	Summary (404, 587, 604).
Figs, edible portion.			1			69	
Hazelnuts, edible portion.	Switzerland.	177	1			15	
Lettuce, edible portion.			41	6,740	71	1,137	Summary (5, 153, 258, 282; 404, 445).
Loraberries.	Oregon.	363	1			160	
Millet, grain.			9	36	2	10	Summary (3, 4, 548).
Mustard, greens.			2	932	169	560	
Oats, grain.	Oklahoma.	258	13	175	10	57	Summary (45, 363, 604).
Oats, straw.			4	764	335	524	
Okra, fruit.	Germany.	45	3	1,075	412	697	Summary (258, 365).
Onion, bulb.			23	479	46	188	Summary (45, 404, 445).
Farsnips, root.	New Zealand.	264	2	180	100	140	

⁹ Data from the following references have been omitted because the moisture content of the material is not given: 76, 118, 175, 177.

¹⁰ McHargue (577) reported 250,000 p. p. b. of iodine in a sample of corn raised in Kentucky, while Adolph (6) reported that he found none in a sample raised in Nebraska.

TABLE 21.—The minor-element content of some important crops—Continued

IODINE—Continued

Plant or part of plant	Location	Reference No.	Analyses	Iodine (parts per billion)			Remarks
				Maximum	Minimum	Mean	
Peach, edible portion			Number				
Pear, edible portion			2	192	11	101	Summary (363, 365).
Peas, green, edible portion			2	15	15	15	Summary (5, 363).
Peas, seed	California	365	1			9	
Peas, field, above-ground portion, cut for hay	Germany	45	3	120	72	97	
Pineapple, fruit	South Carolina	414	2	1, 317	595	956	
Potatoes, tubers	Hawaii	587	1			1, 070	
Do	Pennsylvania	7x	10	446	162	238	Hagerstown soil.
			360	446	7	97	Summary (5, 45, 77, 198, 258, 364, 365, 385, 445, 481, 587, 604, 7x, 14x).
Prunes, fruit	Oregon	367	1			5	
Radish, roots	Oklahoma	265	1			994	
Redtop, above-ground portion, cut for hay	Kentucky	386	14			89	
Rice, polished	Manchuria	559	1			40	
Rutabaga, roots	New Zealand	271	4	850	460	615	
Rye, grain			5	133	3	83	Summary (45, 363).
Rye, straw	Germany	46	4	770	292	497	
Soybeans, above-ground portion, cut for hay	South Carolina	423	35	456	50	237	Summary (5, 45, 258, 274, 404, 481, 509, 510, 525, 587, 604).
Spinach			91	48, 650	19	9, 382	Iodine fertilization.
Do	Germany	{ 45, 609, }	24	48, 650	15, 000	26, 417	
	do	{ 509, 510, }	36	423	2	87	
Sugar beets, roots	do	{ 542, 604 }	25	2, 010	110	425	
Sugar beets, leaves	do	{ 510, 542, }	25				
		{ 604 }	4	115	70	92	Summary (258, 365, 481, 587).
Sweetpotatoes, roots	Oklahoma	265	1			515	
Swiss chard, edible portion	Kentucky	386	13			71	
Timothy, above-ground portion, cut for hay			22	660	20	233	Summary (45, 285, 365, 385, 404, 587, 7x).
Tomato, fruit			22	870	223	484	Summary (258, 264).
Turnip, roots			4	676	111	255	Summary (258, 274).
Turnip, leaves			22	2, 080	740	1, 434	No fertilization.
Turnip, whole plant	Pennsylvania	390	5	94, 960	19, 540	42, 304	Fertilized with K.I.
Do	do	390	5			546	
Vetch, purple, above-ground portion, cut for hay	South Carolina	414	1			547	
Vetch, hairy, above-ground portion, cut for hay	do	414	1			549	
Vetch, woolly-pod, above-ground portion, cut for hay	do	414	1			549	
Vetch, monantha, above-ground portion, cut for hay	do	414	1			328	
Vetch, above-ground portion, cut for hay	do	414	2	330	268	299	Variety unknown.
Wheat, grain			26	168	0	67	Summary (45, 265, 363).
Wheat, straw			6	772	67	455	Summary (45, 381).
Yams, tuber	Hawaii	587	1			173	

IRON ¹¹

Plant or part of plant	Location	Reference No.	Analyses	Iron			Remarks
				Maximum	Minimum	Mean	
Alfalfa, above-ground portion, prebud stage	Tennessee	580	Number	Mg./kg.	Mg./kg.		
Alfalfa, above-ground portion, full bloom	-----	-----	3	350	140	247	
Alfalfa, above-ground portion, cut for hay	Pennsylvania	495	4	1,000	161	428	
Do	do	495	1	-----	-----	287	
Do	Tennessee	22x	4	343	238	270	
Do	do	22x	4	308	217	262	
Do	-----	-----	4	629	132	275	
Do	-----	-----	19	42	42	42	
Almond, edible portion	Wisconsin	319, 455	2	38	21	29	
Apple, fruit, mature	-----	-----	8	69	1	15	
Artichoke, edible portion	-----	-----	24	337	69	203	
Artichoke, flower head	-----	-----	2	-----	-----	43	
Apricot, fruit	Wisconsin	455	1	979	80	337	
Asparagus, edible portion	-----	-----	5	575	12	264	
Avocado, edible portion	-----	-----	9	72	8	28	
Banana, edible portion	-----	-----	6	350	14	99	
Barley, grain	-----	-----	23	210	79	139	
Beans, field; seed	-----	-----	16	-----	-----	146	
Beans, garden; edible pod and seed	Florida	5x	4	175	125	146	
Do	-----	-----	30	769	125	231	
Beans, garden; edible hulls	Kentucky	379	1	-----	-----	270	
Beans, garden; immature seed	-----	-----	3	270	80	131	
Beans, lima; mature seed	do	371, 379	6	150	74	112	
Beets, roots	New Hampshire	495	3	-----	-----	104	
Do	-----	-----	14	280	69	133	
Beets, leaves	-----	-----	7	1,932	170	795	
Blackberries, fruit	Wisconsin	455	1	-----	-----	63	
Blueberries, fruit	do	455	1	-----	-----	22	
Brazil nut, edible portion	do	455	1	-----	-----	42	
Broccoli, edible portion	-----	-----	2	210	200	205	
Brussels sprouts, edible portion	-----	-----	2	210	177	193	
Buckwheat, grain	Wisconsin	455	1	-----	-----	34	
Buckwheat, above-ground portion, cut in bloom	France	58	1	-----	-----	100	
Bur clover; above-ground portion, early green	California	245	1	-----	-----	120	
Butternut, edible portion	Wisconsin	455	1	-----	-----	70	

¹¹ The analyses of Shackleton and McCance (593) and of Stiebelling (592) have been omitted because the moisture content of the materials is not given.

Summary (58, 580).
Hagerstown loam.
Penn silt loam.
Cumberland loam.
Fullerton silt loam.
Summary (373, 495, 522, 580, 14x, 22x).
Summary (91, 455).
Summary (276, 393).
Summary (393, 455).
Summary (347, 393, 455, 457).
Summary (393, 408, 562).
Summary (163, 319, 408, 455).
Summary (222, 368, 455, 462, 528).
Summary (368, 379, 455). *Phaseolus vul-*
garis.
St. Lucie soil.
Summary (393, 457, 482, 5x).
Summary (379, 455).
Gloucester stony loam.
Summary (393, 455, 482, 495, 8x).
Summary (393, 413, 455, 457, 482).
Summary (347, 393).
Summary (393, 455).
Medicago hispida.

TABLE 21.—The minor-element content of some important crops—Continued

IRON—Continued

Plant or part of plant	Location	Reference No.	Analyses	Iron			Remarks
				Maximum	Minimum	Mean	
Cabbage, edible portion.	Alabama	13x	Number	Mg./kg.	Mg./kg.		
Do.	do.	13x	1		208	Decatur soil.	
Do.	do.	13x	1		180	Hartsell soil.	
Do.	do.	13x	4	290	81	Norfolk soil.	
Do.	Florida	5x	8	200	140	Leon soil (NPK).	
Do.	do.	5x	2			Leon soil (NPKMn).	
Do.	do.	5x	2			Leon soil (NPKCu).	
Do.	do.	495	1			Cecil clay.	
Do.	North Carolina	495	1	305	11	Summary (393, 455, 457, 482, 495, 5x, 13x)	
Cabbage, Chinese, edible portion.	Hawaii	113	67	875	202	<i>Brassica pekinensis</i> (B. pe-tsai).	
Cabbage, white mustard; edible portion.	do.	113	2	490	461	<i>B. chinensis</i> .	
Cabbage, Chinese mustard; edible portion.	do.	113	1			<i>B. juncea</i> .	
Cantaloup, edible portion.	do.		7	210	54	Summary (393, 455).	
Carrots, roots.	Rhode Island	19x	8	350	140	Bridgeton very fine sandy loam (NPK).	
Do.	New Hampshire	495	1			Gloucester stony loam.	
Do.	do.		27	490	49	Summary (166, 393, 455, 482, 495, 8x, 19x).	
Carrots, leaves.	South Carolina	482	5	765	355		
Cashew, mature seed, edible portion.	Philippine Islands	393	1				
Cauliflower, edible portion.	do.		4	280	52	Summary (393, 455, 457).	
Celery, edible portion.	Florida	5x	3			Bladen soil.	
Do.	do.		32	350	110	Summary (393, 455, 5x).	
Celery cabbage, edible portion.	Wisconsin	455	1			Summary (393, 408, 455).	
Cherries, edible portion.	do.		3	140	28		
Chestnut, Italian; edible portion.	Wisconsin	455	1			<i>Cyrus medica</i> .	
Citron, edible portion.	Puerto Rico	179	1			Mauzy silt loam. <i>Trifolium agrarium</i> .	
Clover, hop; above-ground portion, before bloom.	Tennessee	22x	1			Do.	
Clover, hop; above-ground portion, in bloom.	do.	22x	2	181	133	Do.	
Clover, hop; above-ground portion.	do.	22x	1			Do.	
Clover, hop; leaves.	do.	22x	1			Do.	
Clover, hop; stems.	do.	22x	1			Do.	
Clover, hop; above-ground portion.	Kentucky	14x	2	490	410	Mauzy silt loam. <i>Trifolium agrarium</i> .	
Clover, alsike; above-ground portion.	do.	371	1			<i>T. hybridum</i> .	
Clover, alsike; seeds.	do.	371	1			Do.	
Clover, crimson; seeds.	do.	371	1			Do.	
Clover, low hop; above-ground portion, pre-bloom.	Tennessee	580	1			<i>T. incarnatum</i> .	
Clover, low hop; above-ground portion, in bloom.	do.	580	1			<i>T. procumbens</i> .	
Clover, low hop; above-ground portion, bloom dying.	do.	580	1			Do.	
Clover, red; above-ground portion.	Kentucky	14x	5	190	130	Tilsit silt loam. <i>T. pratense</i> .	
Do.	do.		42	1,300	100	Summary (524, 14x). <i>T. pratense</i> .	
Clover, red; seeds.	Kentucky	371, 373	2	336	21	<i>T. pratense</i> .	

Clover, red, foliage.	373	1	580	120	364
Clover, white; above-ground portion	19	18	35	20	<i>T. repens</i> .
Clover, white; seeds	371	1	44	39	Do.
Coconut, edible portion	13x	3	328	219	Summary (393, 455).
Collards, leaves	13x	1	534	111	Norfolk soil.
Do.		7	50	25	Vaiden soil.
Corn, grain	14x	4	188	160	Summary (413, 521, 13x).
Do.		2	345	94	Summary (393, 455, 528).
Corn, stover		16			Tilsit silt loam.
Do.		1			Summary (528, 14x).
Cotton, lint	374	1	662	81	Norfolk soil.
Cotton, seed	374	1	1,909	428	Summary (113, 13x).
Covpos, seed	13x	3	420	109	Summary (113, 393, 455).
Do.	416, 417	3	70	35	Summary (368, 455).
Cranberries, edible portion		1	266	45	Summary (115, 455).
Cress, water, edible portion	455	4	64	10	Summary (113, 393, 455, 586).
Cucumber, edible portion		2	118	85	Summary (408, 455).
Dandelion, edible portion		1	58	36	Summary (368, 371).
Dates, edible portion		2	210	37	Summary (393, 455).
Eggplant, edible portion	393	2	112	13	Summary (408, 455, 11x).
Endive, edible portion		6			
Figs		1			
Flax, seed	455	1			
Gooseberries	199	1			
Grapefruit, fruit		2			
Grapes, edible portion		2			
Hazelnuts, edible portion	455	1			
Hemp, seeds	371	1			
Hickory nuts, edible portion	455	1			
Horsebeans, forage	368	1			
Horsebeans, seed	368	1			
Horsebeans, straw	368	1			
Kafr, grain	528	1			
Kale, edible portion		2	979	399	<i>Vicia faba</i> .
Kale, thousand-head; above-ground portion	600	1	59	43	Do.
Kale, narrow stem; above-ground portion	600	2			Do.
Kentucky bluegrass, above-ground portion, young blades.	375	1			Summary (113, 393).
Kentucky bluegrass, above-ground portion, in seed.	495	1			
Kentucky bluegrass, above-ground portion, state of maturity unknown.		4	425	60	Penn silt loam.
Kentucky bluegrass, seeds	371, 375	2	460	350	Summary (373, 381, 524, 528).
Kohlrabi, edible portion		3	420	61	Summary (393, 455, 457).
Kumquat, edible portion	455	1			
Lemon, peel	455	1			
Lemon, pulp	393	1	231	119	Clarksville silt loam.
Lespedeza, above-ground portion, before bloom	22x	4	1,028	259	Cumberland loam.
Do.	22x	13			Decatur silt loam.
Do.	22x	1			Meury silt loam.
Do.	22x	2	552	245	Montevallo silt loam.
Do.	22x	1			

TABLE 21.—The minor-element content of some important crops—Continued

IRON—Continued

Plant or part of plant	Location	Reference No.	Analyses	Iron			Remarks
				Maximum	Minimum	Mean	
Lespedeza, above-ground portion, early bloom.	Tennessee.	22x	Number 3	Mg./kg. 196	Mg./kg. 133	Mg./kg. 159	Clarksville silt loam.
Do.	do.	22x	7	455	119	295	Cumberland loam.
Do.	do.	22x	7	266	217	240	Maury silt loam.
Lespedeza, above-ground portion, full bloom.	do.	22x	3	371	133	235	Clarksville silt loam.
Do.	do.	22x	13	685	252	355	Cumberland loam.
Do.	do.	22x	2	301	203	252	Dandridge silt loam.
Do.	do.	22x	1	—	—	224	Fullerton silt loam.
Do.	do.	22x	1	—	—	322	Montevallo silt loam.
Lespedeza, above-ground portion, late bloom.	do.	22x	6	643	112	241	Clarksville silt loam.
Do.	do.	22x	1	—	—	224	Maury silt loam.
Lespedeza, above-ground portion, in seed	do.	22x	1	—	—	217	Clarksville silt loam.
Do.	do.	22x	10	979	106	347	Cumberland loam.
Lespedeza, above-ground portion, maturity unknown.	do.	22x	1	—	—	259	Clarksville silt loam.
Do.	do.	580	3	231	126	166	Do.
Do.	do.	22x	28	601	161	232	Cumberland loam.
Do.	do.	580	17	1,329	182	445	Hartsells silt loam.
Do.	do.	22x	2	231	203	217	Maury silt loam.
Do.	do.	580	4	413	217	273	Do.
Lettuce, edible portion	Alabama.	13x	1	—	—	1,500	Cecil soil.
Do.	do.	13x	2	780	600	691	Norfolk soil.
Do.	do.	13x	2	4,830	65	1,034	Summary (113, 319, 347, 351, 455, 482, 8x, 13x).
Lettuce, leaf	Wisconsin.	455	1	—	—	334	Bridgehampton very fine sandy loam (NPK).
Mangel-wurzel, roots	Rhode Island	19x	4	210	140	175	Do.
Mangel-wurzel, crown	do.	19x	4	699	280	455	Do.
Mangel-wurzel, leaves	do.	19x	4	1,259	1,189	1,206	Do.
Millet, grain	Kentucky	371	1	—	—	52	Do.
Millet, above-ground portion, fruiting	Philippine Islands	364	1	—	—	560	Do.
Millet, Japanese; above-ground portion, cut for hay	Wisconsin	328	1	—	—	59	Do.
Milo, above-ground portion, flowering	Philippine Islands	394	1	—	—	420	Do.
Milo, above-ground portion, fruiting	do.	394	1	—	—	49	Do.
Mulberry, fruit	do.	393	1	—	—	280	Do.
Mushrooms, edible portion	Wisconsin	455	1	—	—	109	Do.
Mustard, greens	Alabama	13x	2	600	300	450	Norfolk soil.
Do.	do.	393	5	858	300	858	Do.
Mustard, Chinese; leaves	Philippine Islands	393	2	1,540	1,259	1,399	Summary (181, 521, 13x).
Mustard, above-ground portion, cut for fodder	Germany	368	2	350	350	350	Do.

Mustard, straw	368	1	1	350	70	
Oats, grain	370	34	7	370	79	Summary (222, 368, 371, 455, 528).
Oats, straw	58	203	61	370	203	Summary (382, 528).
Oats, above-ground portion, flowering stage	495	1			50	
Oats, above-ground portion, milk stage		1		140	154	Penn silt loam.
Okra, fruit		1		265	108	Summary (393, 482).
Onions, bulb		8		70	130	Summary (455, 457, 482).
Orange, edible portion	408	2			44	Summary (393, 408).
PaPaw, edible portion	393	1		210	17	<i>Carica papaya</i> L.
Do		3		1,609	117	Do.
Parsley, leaves	393	4		210	1,154	Summary (319, 393, 455).
Parsnip, root		3		28	140	Summary (393, 455).
Peach, edible portion		3		140	24	Summary (393, 455).
Peanut, edible portion	393	6		282	84	Summary (393, 455).
Pear, edible portion		2			126	Summary (113, 319, 371, 455).
Pears, green; edible portion		1			26	
Pecans, edible portion	455	1			68	
Peppers, green; edible portion	455	1			72	
Peppers, red; edible portion	455	1			72	
Peppers, sweet; fruit	393	1			630	<i>Capsicum annuum</i> L.
Pineapple, edible portion		4		280	144	Summary (393, 408).
Pineapple, fruit	455	1			40	
Pistachio nut, edible portion	455	1		140	825	Summary (393, 455).
Plums, edible portion		3			110	
Pomegranate, seeds	455	1		87	44	Bladen soil (NPK).
Potatoes, tubers	5x	6			75	Bladen soil (NPKMn).
Do	5x	1			83	Bladen soil (NPKMn).
Do	5x	1			74	Bladen soil (NPKCu).
Do	5x	4		107	92	Coxville soil (NPK).
Do	5x	3			101	Coxville soil (NPKMn).
Do	5x	2			85	Coxville soil (NPKCu).
Do	5x	1			94	Coxville soil (NPKMnCu).
Do	5x	4		71	54	Leon soil (NPK).
Do	5x	1			69	Norfolk soil (NPK).
Do	495	1			28	Norfolk stony loam.
Do		131		363	105	Summary (137, 213, 368, 393, 413, 455, 495, 5x, 20x).
Prunes, edible portion	506	15		75	55	Summary (113, 209, 393, 455).
Quince, edible portion	455	1			58	Norfolk soil.
Radish, roots		14		825	436	Summary (528, 13x).
Rape, above-ground portion	13x	1			210	
Do		2		244	176	
Rape, above-ground portion, flowering stage	58	1		87	84	
Rape seed	371	1			85	
Raspberries, fruit	455	1			62	
Redtop, above-ground portion, cut for hay	14x	1			130	Dekalb silt loam; no fertilization.
Do	14x	1			120	Dekalb silt loam (M).
Do	14x	1			160	Dekalb silt loam (ML).
Do	14x	1			110	Tilist silt loam; no fertilization.
Do	14x	1			110	Tilist silt loam (NP).
Do	14x	1			110	Tilist silt loam (NPL).
Do	14x	12		160	117	Summary (394, 14x).

TABLE 21.—The minor-element content of some important crops—Continued
IRON—Continued

Plant or part of plant	Location	Reference No.	Analyses	Iron			Remarks
				Maximum	Minimum	Mean	
			Number	Mg./kg.	Mg./kg.	Mg./kg.	
Redtop, seeds	Kentucky	371	1	559	154	160	Summary (393, 455).
Rhubarb, stem			2			356	
Rice, brown		380	1			76	Summary (380, 455).
Rice, polished			3	32	12	19	
Rutabagas, roots	Philippine Islands	393	3	559	56	275	Summary (393, 455).
Rutabagas, leaves	Wisconsin	528	1			1,389	
Rye, grain	do	455	1			57	Summary (113, 409, 528, 603).
Salsify, edible portion	Kentucky	372	1			336	
Soybean, leaves			41	161	59	101	Summary (371, 372, 393).
Soybean, seed, green			13	133	57	80	
Soybean, seed, mature			1			160	Summary (113, 409, 528, 603).
Soybean, above-ground portion, cut for hay	Kentucky	14x	1			140	
Do	do	14x	1			140	Summary (113, 409, 528, 603).
Do	do	14x	2			140	
Do	do	14x	2	170	100	140	Summary (371, 372, 393).
Do	do	14x	2	130	100	140	
Do	do	14x	2	150	110	125	Summary (113, 409, 528, 603).
Do	do	14x	2	160	110	125	
Do	do	14x	1			130	Summary (113, 409, 528, 603).
Do	do	14x	1			110	
Do	do	14x	1			110	Summary (113, 409, 528, 603).
Do	do	14x	1			110	
Do	do	14x	2	350	150	150	Summary (371, 372, 393).
Do	do	14x	2	300	180	140	
Do	do	14x	3			140	Summary (113, 409, 528, 603).
Do	do	14x	3			140	
Do	do	14x	2	130	110	120	Summary (113, 409, 528, 603).
Do	do	14x	2	350	320	130	
Do	do	14x	2	570	100	212	Summary (394, 528, 14x).
Do	do	14x	33			1,319	
Spinach, edible portion	Alabama	18x	1	517	299	408	Norfolk soil.
Do	do	18x	2	1,750	54	683	
Do			54			105	Summary (113, 181, 302, 319, 347, 393, 411, 455, 457, 482, 489, 8x, 13x).
Do	South Carolina	482	3	130	70	135	
Squash, edible portion			3	267	68	70	Summary (393, 408, 455).
Strawberries, edible portion	Philippine Islands	394	1			140	
Sudan grass, above-ground portion, before flowering	do		1			53	Summary (393, 408, 455).
Sudan grass, above-ground portion, flowering	Wisconsin	394	1			179	
Sugar beets, roots	do	528	1			140	Summary (393, 408, 455).
Sugar beets, leaves	do	528	1			179	

Sunflower, seeds.....	Kentucky.....	371	1		34				
Sweetclover, above-ground portion.....	do.....	381	18	140	130				<i>Melilotis alba.</i>
Sweetpotatoes, tubers.....			3	1,049	70				Summary (393, 482).
Swiss chard, edible portion.....	Wisconsin.....	455	1		690				Summary (113, 393, 455).
Tangerine, edible portion.....	Alabama.....	13x	1		44				
Tendergreen, edible portion.....	do.....	13x	1		1,415				Decatur soil.
Timothy, above-ground portion, in bloom.....	New Hampshire.....	495	1		262				Norfolk soil.
Do.....	Pennsylvania.....	495	1		154				Gloucester stony loam.
Do.....	do.....	495	1		238				Hagerstown loam.
Do.....	Maine.....	495	1		287				Penn silt loam.
Do.....	Kentucky.....	14x	4	160	135				Tiltsit silt loam.
Do.....	do.....	14x	14	160	116				Summary (528, 14x).
Do.....	Kentucky.....	371	1		61				
Do.....	do.....	371	1		240				
Tobacco, seed.....	Florida.....	5x	26	290	159				Parkwood fine sandy loam (NPK).
Tomato, fruit.....	do.....	13x	56	800	202				Summary (107, 378, 368, 455, 482, 5x).
Do.....	Alabama.....	13x	3	77	57				Norfolk soil.
Do.....	do.....	13x	1		92				Cecil soil.
Do.....	do.....	13x	1		66				Decatur soil.
Do.....	do.....	13x	9	210	92				Summary (368, 393, 455, 13x).
Turnip, tops.....	Germany.....	368	7	2,483	618				Summary (113, 181, 347, 368, 413, 482, 521).
Vetch, above-ground portion, in bloom.....	do.....	368	1		280				
Vetch, above-ground portion, seeds beginning to ripen.....	do.....	368	1		629				
Vetch, above-ground portion, cut for hay.....	Germany.....	368	2	354	317				Summary (166, 368).
Vetch, straw, seeds ripening.....	do.....	368	1		63				Summary (368, 371).
Vetch, seed.....	Germany.....	368	4	85	53				Summary (393, 455).
Walnuts, edible portion.....	do.....	14x	3	61	37				Summary (393, 408, 455).
Watermelon, edible portion.....	do.....	14x	3	280	109				Tiltsit silt loam; no fertilization.
Wheat, grain.....	Kentucky.....	14x	1		140				Tiltsit silt loam (M).
Do.....	do.....	14x	1		80				Tiltsit silt loam (M).
Do.....	do.....	14x	1		90				Tiltsit silt loam (ML).
Do.....	do.....	14x	2	70	65				Tiltsit silt loam (MPL).
Do.....	do.....	14x	131	420	68				Summary (222, 224, 238, 368, 371, 381, 528, 543, 1x, 14x).
Do.....	do.....	14x	1		150				Tiltsit silt loam; no fertilization.
Wheat, straw.....	Kentucky.....	14x	1		130				Tiltsit silt loam (M).
Do.....	do.....	14x	1		130				Tiltsit silt loam (ML).
Do.....	do.....	14x	2	130	125				Tiltsit silt loam (MLP).
Do.....	do.....	14x	39	630	177				Summary (166, 368, 381, 528, 14x).
Do.....	do.....	14x	1		340				
Yams, tuber.....	Hawaii.....	113	1						

TABLE 21.—The minor-element content of some important crops—Continued

MANGANESE

Plant or part of plant	Location	Refer- ence No.	Analyses	Manganese			Remarks
				Maximum	Minimum	Mean	
Alfalfa, above-ground portion, prebud stage	Tennessee	580	Number	Mg./kg.	Mg./kg.		
Alfalfa, above-ground portion, 1/10 bloom	Michigan	507	3	101	67		
Alfalfa, above-ground portion, full bloom			4	34	28	Summary (58, 253, 507, 580).	
Alfalfa, above-ground portion, cut for hay	Pennsylvania	495	8	936	14	Hagerstown loam.	
Do	do	495	1			Penn silt loam.	
Do	Tennessee	22x	1	54	77	Cumberland loam.	
Do	do	22x	4	124	70	Fullerton silt loam.	
Do	do	22x	32	124	16	Summary (80, 306, 373, 495, 580, 585, 14x, 22x).	
Alfalfa, leaves	Oregon	306	2	76	45		
Alfalfa, stems	do	306	2	13	11	Summary (306, 371, 373).	
Alfalfa, seeds			4	15	6	Summary (299, 461).	
Almond, edible portion			2	13	13	Gloucester stony loam.	
Apple, fruit	New Hampshire	495	1			Summary (299, 356, 495).	
Do	do		4	22	1	Summary (299, 461).	
Artichoke, flower head			2	26	23	Summary (63, 73, 299, 356).	
Asparagus, edible portion			5	29	12	Summary (117, 158, 163, 299, 356, 485).	
Banana, edible portion			6	117	7	Summary (73, 138, 356, 368, 485, 528).	
Barley, grain			12	38	7		
Barley, straw			1				
Beans, field; mature seed	Germany	368	16	36	14	Summary (73, 356, 368, 379, 485). <i>Phaseolus vulgaris</i> .	
Beans, garden; edible pods and seeds	Florida	5x	4	56	48	St. Lucie soil.	
Do	Wisconsin	498	2	44	43	Boone fine sandy loam.	
Do	do	498	3	39	36	Carrington silt loam.	
Do	do	498	1			Colby silt loam.	
Do	do	498	1			Kennan silt loam.	
Do	do	498	1			Miami gravelly sandy loam.	
Do	do	498	1	51	30	Miami silt loam.	
Do	do	498	6	57	30	Plainfield sand.	
Do	do	498	43	57	11	Summary (299, 421, 458, 461, 482, 5x).	
Beans, garden; edible hulls	Kentucky	379	1				
Beans, garden; immature seed	do		6	19	16	Summary (356, 379).	
Beans, lima; mature seed			2	25	12	Gloucester stony loam.	
Beets, roots	New Hampshire	495	1			Summary (73, 356, 413, 421, 482, 485, 495, 585, 5x).	
Do	do		20	104	19		
Beets, leaves			5	205	16	Summary (300, 482).	
Blackberries, fruit	Wisconsin	366	1				
Blueberries, fruit	do	356, 461	2	315	122		

Brazil nut, edible portion.....	461	1	32	21	10
Broccoli, edible portion.....	140	26			
Brussels sprouts, edible portion.....	356	21			
Buckwheat, above-ground portion, cut in bloom.....	58	1			
Cabbage, edible portion.....	5x	1	35	24	34
Do.....	5x	2			26
Do.....	5x	2			38
Do.....	5x	2			41
Do.....	5x	1			29
Do.....	495	1			49
Do.....	458	2	11	13	12
Do.....		68	436	5	31
Cabbage, main rib of leaf.....	63	1			21
Do.....	63	40			40
Cabbage, stalk.....	63	1			63
Cantaloup, edible portion.....	461	1			8
Carrots, root.....		19	91	6	31
Carrots, leaves.....		6	109	23	105
Cauliflower, edible portion.....		5	27	7	17
Celery, edible portion.....	5x	3	64	8	50
Do.....		34			30
Celery cabbage, edible portion.....	461	1			23
Cherries, edible portion.....	356	1			2
Chestnuts, edible portion.....	22x	2	56	16	36
Clover, hop; above-ground portion, before bloom.....	22x	1			85
Clover, hop; above-ground portion, in bloom.....	22x	2	70	54	62
Clover, hop; above-ground portion.....	22x	1			85
Clover, hop; leaves.....	22x	1			116
Clover, hop; stems.....	22x	1			54
Clover, alsike; above-ground portion.....		6	288	20	117
Clover, alsike; seeds.....	371	1			28
Clover, crimson; above-ground portion, full bloom.....	495	1			387
Clover, crimson; above-ground portion.....		4	387	24	246
Clover, crimson; seed.....		1			29
Clover, low hop; above-ground portion, prebloom.....	580	1			72
Clover, low hop; above-ground portion, in bloom.....	580	1			58
Clover, low hop; above-ground portion, bloom dying.....	580	1			72
Clover, red; above-ground portion.....	14x	5	120	60	94
Clover, red; above-ground portion, mature with blossoms and seeds.....	495	1			465
Clover, red; above-ground portion, in bloom.....	495	1			287
Clover, red; above-ground portion.....	495	1			542
Do.....		51	542	25	121
Clover, red; seeds.....	371, 373	2	38	25	31
Clover, red; foliage.....		1			200
Clover, red; bloom.....	306	2	66	30	48
Clover, red; leaves.....	306	2	84	40	62
Clover, red; seeds.....		2	12	6	9

Coxville soil (NPK).
 Leon soil (NPK).
 Leon soil (NPKMn).
 Leon soil (NPKCu).
 Parkwood soil (NPK).
 Cecil clay.
 Miami silt loam.
 Summary (65, 299, 356, 421, 482, 485, 495, 495, 585, 5x, 13x).

Summary (299, 356, 421, 482, 485, 495, 585, 8x).
 Summary (299, 482).
 Summary (140, 299, 421, 461).
 Bladen soil.
 Summary (356, 421, 585, 5x).

Summary (300, 461).
 Maury silt loam. *Trifolium agrarium*.
 Do.
 Do.
 Do.
 Do.
 Summary (306, 368, 585, 14x). *T. hybridum*.
T. hybridum.
 Cecil sand loam. *T. incarnatum*.
 Summary (368, 371, 495, 585). *T. incarnatum*.

T. incarnatum.
T. procumbens.
 Do.
 Do.

Thist silt loam. *T. pratense*.
 Gloucester stony loam. *T. pratense*.

Hagerstown loam. *T. pratense*.
 Penn silt loam. *T. pratense*.
 Summary (300, 368, 495, 585, 14x). *T. pratense*.

T. pratense.
 Do.
 Do.
 Do.
 Do.

TABLE 21.—The minor-element content of some important crops—Continued

MANGANESE—Continued

Plant or part of plant	Location	Refer- ence No.	Analyses	Manganese			Remarks
				Maximum	Minimum	Mean	
			Number	Mg./kg.	Mg./kg.	Mg./kg.	
Clover, red; stems.....	Oregon.....	306	2	20	15	17	<i>T. pratense</i> .
Clover, white; above-ground portion, beginning to bloom.	Pennsylvania.....	495	1			356	Hagerstown loam. <i>T. repens</i> .
Clover, white; above-ground portion, some blossoms seeded.	Maine.....	495	1			875	<i>T. repens</i> .
Clover, white; above-ground portion	Kentucky.....		3	144	18.5	102	Summary (368, 585). <i>T. repens</i> .
Clover, white; seeds	Wisconsin.....	371	1			22	<i>T. repens</i> .
Coconut, edible portion	Alabama.....	461	3	286	53	25	
Collards, leaves	South Carolina.....	413	1			147	Norfolk soil.
Corn, grain.....	Kentucky.....		13	19	5	13	Summary (138, 300, 356, 485, 528).
Do.		14x	2	114	103	108	Tilist silt loam.
Corn, sweet; edible portion			20	270	54	127	Summary (507, 528, 14x).
Cotton, lint	Mississippi.....	374	2	31	5	18	Summary (413, 461).
Cotton, seed	do.		1			190	
Cowpeas, seed	South Carolina.....	413	1			13	
Cranberries, edible portion			2	49	25	45	Summary (416, 417, 461).
Cress, water, edible portion			3	69	25	37	Summary (63, 290).
Cucumber, edible portion			4	48	24	56	Summary (461, 585).
Currants, edible portion	Wisconsin.....	356	1			4	
Dandelion, edible portion	do.	461	1			19	
Dates, edible portion			3	54	2	32	Summary (115, 356).
Eggplant			5	53	15	32	Summary (413, 461, 482).
Endive, edible portion	Wisconsin.....	461	1	102	38	61	Summary (368, 371).
Flax, seed			3			5	
Gooseberries	Wisconsin.....	356	1			5	
Grapefruit, fruit	Florida.....	199	2	5	4	5	
Grapefruit, edible portion	Wisconsin.....	356	1			Tr.	
Grapes, edible portion	do.	356	2			Tr.	
Hazelnuts, edible portion			2	43	33	38	Summary (299, 461).
Hemp, seeds			2	165	34	100	Summary (371, 585).
Horsebeans, seed	Germany.....	368, 585	2	15	14	14	<i>Vicia faba</i> .
do.	do.		2			23	Do.
Horsebeans, straw	do.	368	1			36	Do.
Horsebeans, forage			1			36	
Kafir, grain	Wisconsin.....	528	2	16	16	16	Yolo silt loam.
Kale, edible portion	California.....	140	1			60	Seasafra sandy loam.
Do.	Maryland.....	140	2	61	60	137	Norfolk silt loam.
Do.	Virginia.....	140	2	137	66	101	
Do.			6	137	11	68	Summary.

Kentucky bluegrass, above-ground portion, young blades.	Kentucky	375	1	-----	80	Penn silt loam.
Kentucky bluegrass, above-ground portion, in seed.	Pennsylvania	495	1	-----	29	Boise sandy loam.
Kentucky bluegrass, above-ground portion, stage of maturity unknown.	Idaho	80	10	107	69	Summary (80, 368, 373, 381, 528, 595).
Do.	Kentucky	371, 375	19	216	30	Clarksville silt loam.
Kentucky blue grass, seeds.	Wisconsin	356	1	110	85	Cumberland loam.
Kohlrabi, edible portion.	Wisconsin	356	1	-----	13	Decatur silt loam.
Lemon, edible portion.	France	63	1	-----	4	Maury silt loam.
Lespedeza, above-ground portion, before bloom.	Tennessee	22x	1	147	70	Montevillo silt loam.
Do.	do	22x	13	302	54	Clarksville silt loam.
Do.	do	22x	4	-----	114	Cumberland loam.
Do.	do	22x	1	-----	174	Decatur silt loam.
Do.	do	22x	2	-----	441	Maury silt loam.
Do.	do	22x	1	255	217	Montevillo silt loam.
Lespedeza, above-ground portion, early bloom.	do	22x	3	418	77	Clarksville silt loam.
Do.	do	22x	7	232	62	Cumberland loam.
Do.	do	22x	3	163	132	Maury silt loam.
Do.	do	22x	3	147	77	Clarksville silt loam.
Lespedeza, above-ground portion, full bloom.	do	22x	3	232	54	Cumberland loam.
Do.	do	22x	13	232	54	Dandridge silt loam.
Do.	do	22x	2	163	70	Fullerton silt loam.
Do.	do	22x	1	-----	178	Fullerton silt loam.
Do.	do	22x	1	-----	240	Montevillo silt loam.
Do.	do	22x	3	225	93	Cumberland loam.
Lespedeza, above-ground portion, late bloom.	do	22x	1	-----	173	Cumberland loam.
Do.	do	22x	1	-----	108	Maury silt loam.
Lespedeza, above-ground portion, in seed.	do	22x	1	-----	85	Clarksville silt loam.
Do.	do	22x	1	-----	199	Cumberland loam.
Do.	do	22x	10	364	101	Cumberland loam.
Do.	do	22x	1	-----	62	Clarksville silt loam.
Lespedeza, above-ground portion, maturity unknown.	do	22x	1	-----	178	Do.
Do.	do	580	8	331	86	Cumberland loam.
Do.	do	22x	28	217	54	Cumberland loam.
Do.	do	580	17	317	58	Hartsell silt loam.
Do.	do	22x	2	263	178	Maury silt loam.
Do.	do	580	4	223	130	Do.
Do.	do	580	4	-----	185	Do.
Lettuce, edible portion.	Alabama	13x	1	-----	89	Cecil soil.
Do.	do	13x	2	211	198	Norfolk soil.
Do.	do	13x	2	209	141	Leon soil (NPK).
Lettuce, head, outer leaves removed.	Florida	5x	13	209	141	Leon soil (NPKMn).
Do.	do	5x	3	389	302	Leon soil (NPKCu).
Do.	do	5x	2	-----	292	Portsmouth soil (NPK).
Do.	do	5x	2	-----	174	Portsmouth soil (NPK).
Do.	do	5x	1	-----	196	Summary (140, 299, 356, 413, 482, 485, 585, 5x, 8x, 13x).
Do.	do	5x	44	523	15	5x, 8x, 13x).
Lettuce, edible portion.	do	5x	44	523	15	5x, 8x, 13x).
Lettuce, leaf.	Wisconsin	461	1	-----	167	Do.
Lupine, yellow; seeds.	Netherlands	585	1	-----	678	Cumberland loam.
Mandarin, edible portion.	France	63, 299	2	3	2	Hartsell silt loam.
Millet, grain.	Kentucky	371	1	-----	17	Maury silt loam.
Millet, straw.	Africa	485	1	-----	25	Do.
Millet, German, above-ground portion, cut for hay.	Michigan	507	6	135	94	Do.
Millet, Japanese, above-ground portion, cut for hay.	Rhode Island	19x	1	-----	30	Bridgheampton very fine sandy loam (PKCaMg).
Do.	Wisconsin	528	1	-----	36	Do.

TABLE 21.—The minor-element content of some important crops—Continued
MANGANESE—Continued

Plant or part of plant	Location	Reference No.	Analyses	Manganese			Remarks
				Maximum Mg./kg.	Minimum Mg./kg.	Mean Mg./kg.	
Mushrooms, edible portion	Alabama	133	Number 2	42	7	23	Summary (299, 461). Norfolk soil.
Mustard, greens	France	63	2	160	150	153	
Mustard, above-ground portion, cut for fodder	Germany	368	2	39	27	33	
do	do	368	2	144	144	144	
Oats, grain	France	58	16	76	23	72	
Oats, straw	France	58	12	1,656	4	51	
Oats, above-ground portion, flowering stage	Pennsylvania	495	1			5	
Oats, above-ground portion, milk stage	France	299	1			116	
Oats, above-ground portion, maturity unknown	South Carolina	482	6	280	72	166	
Oats, fruit	France	482	5	62	36	48	
Onions, bulb	France	63, 299	16	96	2	26	Summary (368, 507). Penn silt loam.
Orange, edible portion	Wisconsin	356	2	3	3	3	
Parsley, edible portion	Wisconsin	356	1	16	2	9	Summary (356, 485). Toome fine sandy loam.
Peanut, edible portion	Wisconsin	356	2	4	2	3	
Pear, edible portion	Wisconsin	458	1	20	10	13	Carrington silt loam. Miami fine sandy loam.
Pears, edible portion	do	458	3	17	17	17	
Pears, edible portion, all sizes	do	458	2	25	10	15	Miami silt loam. Summary (299, 356, 357, 371, 413, 468, 485).
Do	do	458	7	25	4	12	
Peas, green, edible portion	France	299	3	12	4	9	Summary (73, 306, 585).
Peas, seed	Africa	485	1			15	
Peas, above-ground portion	France	299	1			27	Summary (299, 366).
Peas, pods only	Oregon	306	1			22	
Do	do	306	1			22	Toome fine sandy loam. Miami silt loam.
Peas, field; stems	do	306	1			14	
Peas, field; blooms	do	306	1			21	Miami silt loam. Summary (299, 356, 357, 371, 413, 468, 485).
Peas, field; leaves	do	306	1			38	
Peas, field; pods	do	306	1			21	Summary (299, 356, 357, 371, 413, 468, 485).
Peas, edible portion	do	461	1			36	
Peppers, green, edible portion	Wisconsin	461	1			10	Fort Collins stony loam; no fertilization. Fort Collins stony loam (N). Bladen soil (NPK). Bladen soil (NPK.Mn).
Peppers, red, edible portion	do	356	1			18	
Pineapple, fruit	Wisconsin	356	1			134	Summary (73, 306, 585).
Pistachio nut, edible portion	do	356	1			7	
Pumpkin, edible portion	Wisconsin	356	1			6	Summary (73, 306, 585).
Pumpkin, seeds	do	356	1			9	
Potatoes, tubers	Colorado	257	5	31	15	23	Fort Collins stony loam; no fertilization. Fort Collins stony loam (N). Bladen soil (NPK). Bladen soil (NPK.Mn).
Do	Florida	257	5	31	8	22	
Do	do	53	6	21	5	14	Summary (73, 306, 585).
Do	do	53	1			42	

Do	do	5x	1	17	10	40
Do	do	5x	4			14
Do	do	5x	3			31
Do	do	5x	2			38
Do	do	5x	1			40
Do	do	5x	4	11	7	10
Do	do	5x	1			11
Do	do	495	1			8
Do	Wisconsin	12x	3	7	4	7
Do			205	94		19
Potatoes, tops	Germany	368	4	108	86	99
Prunes, edible portion	California	506	15	7	4	6
Pumpkin, edible portion	Wisconsin	461	1			4
Pumpkin, fruit	France	289	1			13
Quince, edible portion	Wisconsin	356	1			Tr.
Radish, roots			2	45	14	29
Radish, flesh without skin	France	63	3	21	11	15
Rape, above-ground portion, maturity unknown	Alabama	13x	1			43
Rape, above-ground portion, flowering	France	58, 299	3	77	46	65
Rape, above-ground portion, maturity unknown			2	51	43	47
Rape, seed	Kentucky	371	1			38
Raspberries, fruit	Wisconsin	461	1			46
Redtop, above-ground portion, cut for hay	Kentucky	14x	1			330
Do	do	14x	1			310
Do	do	14x	1			110
Do	do	14x	1			120
Do	do	14x	1			370
Do	do	14x	1			510
Do	do	14x	1			180
Do	do	14x	1			226
Do	do	14x	25	510	79	510
Redtop, seeds		371	1			31
Rhubarb, edible portion	Wisconsin	461	1			23
Rice, brown			2	28	18	Summary (380, 485).
Rice, polished			4	14	10	Summary (373, 380, 461, 485).
Rutabaga, roots	Wisconsin	421, 461	7	10	8	9
Rye, grain			2	157	32	Summary (138, 528).
Salsify, root			2	27	12	Summary (297, 461).
Sorghum, above-ground portion, cut for hay	Michigan	507	10	210	82	131
Soybean, leaves	Kentucky	372	1			160
Soybean, seeds, green	Wisconsin	528	1			29
Soybean, seeds, mature		371, 372	11	41	21	28
Soybean, above-ground portion, cut for hay	Kentucky		1			200
Do	do	14x	1			Dekalb silt loam; no fertilization.
Do	do	14x	1			Dekalb silt loam (MPL).
Do	do	14x	2	280	170	Dekalb silt loam (MPL).
Do	do	14x	2	88	64	Dekalb silt loam (MPL).
Do	do	14x	2	64	45	Dekalb silt loam (MPKL).
Do	do	14x	1			Tiltsit silt loam; no fertilization.
Do	do	14x	1			Tiltsit silt loam (MP).
Do	do	14x	2	155	120	Tiltsit silt loam (MP).
Do	do	14x	3	130	88	Tiltsit silt loam (MPK).
Do	do	14x	1			Tiltsit silt loam (ML).
Do	do	14x	2	97	87	Tiltsit silt loam (MPL).
Bladen soil (NPKCu).						
Coxville soil (NPK).						
Coxville soil (NPKMn).						
Coxville soil (NPKCu).						
Coxville soil (NPKCuMn).						
Leon soil (NPK).						
Norfolk soil (NPK).						
Norfolk stony loam.						
Miami silt loam.						
Gloucester stony loam.						
Summary (79, 257, 356, 368, 413, 421, 485, 496, 5x, 12x, 20x).						
Summary (63, 297, 461).						
Norfolk soil.						
Summary (528, 13x).						
Dekalb silt loam; no fertilization.						
Dekalb silt loam (M).						
Dekalb silt loam (ML).						
Dekalb silt loam (PL).						
Tiltsit silt loam; no fertilization.						
Tiltsit silt loam (N.F.).						
Tiltsit silt loam (N.F.L.).						
Summary (80, 128, 14x).						
Summary (380, 485).						
Summary (373, 380, 461, 485).						
Summary (138, 528).						
Summary (297, 461).						
Dekalb silt loam; no fertilization.						
Dekalb silt loam (M.L).						
Dekalb silt loam (M.F.L).						
Dekalb silt loam (M.P.L).						
Dekalb silt loam (M.P.K.L).						
Tiltsit silt loam; no fertilization.						
Tiltsit silt loam (M).						
Tiltsit silt loam (MP).						
Tiltsit silt loam (MPK).						
Tiltsit silt loam (ML).						
Tiltsit silt loam (MPL).						

TABLE 21.—The minor-element content of some important crops—Continued

MANGANESE—Continued

Plant or part of plant	Location	Refer- ence No.	Analyses	Manganese			Remarks
				Maximum	Minimum	Mean	
Soybean, above-ground portion, cut for hay	Kentucky	14x	Number 2	Mg./kg. 100	Mg./kg. 95	Mg./kg. 95	Tiltsit silt loam (MPKL). Summary (328, 14x). Cecil soil.
Do.	do.	13x	32	280	45	126	
Spinach, edible portion	Alabama	13x	1			48	
Do.	do.	13x	2	280	244	267	Norfolk soil.
Do.	Maryland	140	2	694	84		Sassafras sandy loam.
Do.	Rhode Island	19x	1			52	Bridgehampton very fine sandy loam; no fertilization.
Do.	do.	19x	1			103	Bridgehampton very fine sandy loam (Mn).
Do.	do.	19x	8	56	10	36	Bridgehampton very fine sandy loam (N.P.K.).
Do.	do.	19x	8	120	10	66	Bridgehampton very fine sandy loam (N.P.K.Mn).
Do.	Virginia	140	2	70	65	67	Norfolk loam.
Do.	do.		76	694	10	101	Summary (73, 140, 207, 299, 356, 411, 421, 481, 485, 566, 585, 8x, 13x, 19x).
Squash, edible portion	South Carolina	482	6	28	19	24	Summary (63, 356).
Strawberries, edible portion	do.		2	40	6	23	
Sugar beets, roots	Wisconsin	528	1			206	
Sugar beets, leaves, mature	Michigan	507	21	79	.11	25	
Sugar beets, leaves	Wisconsin	528	1			712	
Sunflower, seeds	Kentucky	371	1			23	
Sunflower, above-ground portion, cut before bloom.	Germany	368	1			72	
Sunflower, above-ground portion, bud appearing on top of plant.	Idaho	431	6	1,066	778	852	Palouse silt loam.
Sunflower, above-ground portion, flower 3 inches in diameter.	do.	431	6	1,268	490	773	Do.
Sunflower, above-ground portion, full bloom	Germany	368	1			72	
Sunflower, above-ground portion, cut before seed of first bloom was in dough stage.	Idaho	431	6	1,263	864	1,015	Palouse silt loam.
Sunflower, above-ground portion, seeds of first flower in dough stage.	do.	431	6	1,224	576	818	Do.
Sunflower, above-ground portion, seeds quite hard and rays fallen.	do.	431	6	1,138	634	807	Do.
Sweetclover, above-ground portion	Pennsylvania	495	1			186	Hagerstown loam. <i>Melilotus alba</i> .
Do.	do.		2	186	50	118	Summary (495, 14x). <i>M. alba</i> .
Sweetpotatoes, tubers	Wisconsin	356	11	28	3	11	Summary (413, 482, 486).
Swiss chard, edible portion	do.	356	1			93	
Tangerine, edible portion	do.		1			3	

Tendergreen, edible portion	Alabama	13x	1			497	Decatur soil.
Do.	do.	13x	1			189	Norfolk soil.
Timothy, above-ground portion, in bloom	New Hampshire	495	1			58	Gloucester stony loam.
Do.	Pennsylvania	495	1			34	Hagerstown loam.
Do.	do.	495	1			36	Penn silt loam.
Timothy, above-ground portion, in seed	Maine	495	1			104	
Timothy, above-ground portion, cut for hay	Kentucky	14x	1		70	82	Tiltsit silt loam.
Do.	do.		28	110	11	86	Summary (80, 128, 254 528, 585, 14x).
Timothy, seed	Kentucky	371	1			72	
Tobacco, cigar wrapper type, leaves	Connecticut	10, 42	38	2, 262	70	657	
Tobacco, leaves, type unknown	do.	43	19	1, 729	144	701	
Tobacco, seed	Kentucky	371	1			70	Parkwood fine sandy loam (NPK).
Tomato, fruit	Florida	5x	26	58	43	51	Summary (378, 461, 482, 5x).
Do.	do.		47	98	18	46	Cecil soil.
Turnip, roots	Alabama	13x	1			5	Decatur soil.
Do.	do.	13x	1			21	Decatur soil.
Do.	do.	13x	3	65	12	51	Norfolk soil.
Do.	do.		17	56	5	19	Summary (48, 63, 297, 368, 413, 461, 485, 585, 13x).
Turnip, tops	Germany	368	11	900	36	185	Summary (63, 297, 368, 370, 482).
Vetch, above-ground portion, in bloom	do.	368	1			72	
Vetch, above-ground portion, seeds beginning to ripen	do.	368	1			144	
Vetch, above-ground portion, cut for hay	Germany	368	3	106	27	54	Summary (166, 300, 368).
Vetch, above-ground portion, pods forming	do.	368	1			360	
Vetch, straw, seeds ripening	do.	368	1			14	Summary (368, 371).
Vetch, seed	do.		4	72	14	34	Summary (299, 356).
Walnuts, edible portion	Wisconsin	356	2	33	19	26	
Watermelon, edible portion	Kentucky	14x	1			2	Tiltsit silt loam; no fertilization.
Do.	do.	14x	1			60	Tiltsit silt loam (M).
Do.	do.	14x	1			70	Tiltsit silt loam (ML).
Do.	do.	14x	1			50	Tiltsit silt loam (ML).
Do.	do.	14x	2	70	50	60	Tiltsit silt loam (MPL).
Do.	do.		109	260	5	49	Summary (73, 138, 255, 341, 368, 371, 381, 485, 528, 543, 585, 14x).
Wheat, straw	Kentucky	14x	1			60	Tiltsit silt loam; no fertilization.
Do.	do.	14x	1			90	Tiltsit silt loam (M).
Do.	do.	14x	1			60	Tiltsit silt loam (ML).
Do.	do.	14x	1			75	Tiltsit silt loam (ML).
Do.	do.	14x	2	80	70	60	Tiltsit silt loam (MPL).
Do.	do.		30	150	22	60	Summary (166, 368, 381, 528, 14x).
Yams, tuber	Africa	485	1			2	

TABLE 21.—The minor-element content of some important crops—Continued

NICKEL

Plant or part of plant	Location	Refer- ence No.	Nickel	Remarks	Plant or part of plant	Location	Refer- ence No.	Nickel	Remarks	
Apricots, edible portion	France	61	Mg./kg. 0.64	<i>Phaseolus vulgaris.</i>	Oats, grain	France	61	Mg./kg. 0.45		
Beans, field; seed	do.	61	.59		Onions, bulb	do.	61	.16		
Buckwheat, grain	do.	61	1.34		Pear, pulp	do.	61	1.30		
Cabbage, edible portion	do.	60	3.3		Peas, green, edible portion	do.	61	2.25		
Carrots, roots	do.	61	.29		Potatoes, tubers	do.	61	.25		
Do.	do.	61	.21		Rice, polished	do.	61	.02		
Carrots, leaves	do.	61	1.83		Soybeans, seed	Kentucky	372	3.92		
Cherries, edible portion	do.	61	.50		Spinach, edible portion	France	61	2.37		
Coffee, bean	do.	61	.38		Tomato, fruit	do.	61	.15		
Corn, grain	do.	61	.14		Walnuts, edible portion	do.	61	.60		
Cress, water	do.	61	.50		Wheat, grain	do.	61	.35		
Figs	do.	61	1.20							
					<i>Nasturtium officinale.</i>					

Plant or part of plant	Location	Reference No.	Selenium Mg./kg.	Remarks
Alfalfa.....	Colorado.....	104	3	Sandy loam. Contained 0.3 p. p. m. of selenium.
Do.....	Utah.....	104	100	Soil contained 1 p. p. m. of selenium.
Do.....	Colorado.....	104	0	Average of 17 samples, omitting the sample from Utah.
Do.....	Various Western States.....	104	2	Boyd clay. Contained 6 p. p. m. of selenium.
Aster, wild; young	South Dakota.....	100	670	Alluvial clay. Contained 2 p. p. m. of selenium.
Aster, wild	do.....	100	60	Boyd clay. Contained 3 p. p. m. of selenium.
Do.....	do.....	100	1,210	Boyd clay loam. Contained 2.5 p. p. m. of selenium.
Do.....	Nebraska.....	100	12	Boyd clay loam. Contained 7 p. p. m. of selenium.
Do.....	do.....	101	5,560	Niobrara clay. Contained 7 p. p. m. of selenium.
<i>Astragalus racemosus</i>	do.....	101	920	Niobrara clay. Contained 1 p. p. m. of selenium.
Do.....	do.....	101	70	Niobrara clay. Contained 1 p. p. m. of selenium.
Do.....	do.....	101	970	Pierre clay. Contained 0.7 p. p. m. of selenium.
Do.....	do.....	101	1	Do.
Do.....	do.....	101	1	Niobrara stony clay loam. Contained 1 p. p. m. of selenium.
Do.....	Various Western States.....	100, 101	9	Average of 89 samples. 75 percent of the plants examined contained from 1 to 15 p. p. m. of selenium.
Vetch, two-groove poison.....	Colorado.....	104	1,380	Pierre clay. Contained 3 p. p. m. of selenium.
Wheat, grain.....	Small plot in South Dakota.....	493	26	Soil contained from 3 to 16 p. p. m. of selenium.
Do.....	Various countries.....	493	0.5	29 samples. Range 0.1 to 1.9 p. p. m. of selenium.

¹² In this table there are listed a few representative samples of selenium contents of some plants grown in seleniferous and nonseleniferous areas. For more complete lists the reader is referred to the literature cited.

TABLE 21.—The minor-element content of some important crops—Continued
SODIUM 13

Plant or part of plant	Location	Reference No.	Analyses	Sodium in moisture-free material			Remarks
				Maximum	Minimum	Mean	
Alfalfa, above-ground portion, prebud stage.....	England.....	601	Number	Percent	Percent	Percent	
Alfalfa, above-ground portion, in bud.....	do.....	600, 601	7	0.18	0.02	0.12	
Alfalfa, above-ground portion, 1/6 to 1/4 bloom.....			13	.33	.07	.22	Summary.
Alfalfa, above-ground portion, 1/2 bloom.....			18	.36	.02	.16	Do.
Alfalfa, above-ground portion, full bloom.....			4	.16	.04	.06	Do.
Alfalfa, above-ground portion, cut for hay.....			38	1.76	.09	.43	Do.
Apples, fruit.....	New Hampshire.....	495	1	.01	.01	.01	Gloucester stony loam.
Barley, grain.....			3	.12	.01	.07	Summary.
Beans, field; seed.....			33	.15	.03	.07	Do.
Beans, garden; edible pod and seed.....			15	.38	.01	.10	Summary.
Beans, garden; edible hulls.....			2	.19	.01	.10	Summary.
Beans, garden; immature seed.....	Kentucky.....	379	1	<i>Phaseolus vulgaris</i> .
Beans, lima; mature seed.....	do.....	379	1	Summary.
Beets; roots.....	Kentucky.....	379	1	
Beets; leaves.....			28	3.78	.04	.74	Summary.
Broccoli, edible portion.....	Netherlands.....	294	20	3.76	.87	2.13	
Buckwheat, above-ground portion, cut in bloom.....	Maryland.....	140	3	.81	.40	.54	
Buckwheat, above-ground portion, past bloom.....	France.....	68	1	
Cabbage, edible portion.....	Massachusetts.....	214	1	
do.....	North Carolina.....	495	1	
Canada bluegrass, above-ground portion.....	Netherlands.....	292	4	.38	.02	.15	Cecil clay.
Carrrots, roots.....	do.....		7	.13	.05	.08	Summary.
Cauliflower, edible portion.....	California.....	140	11	2.88	.07	1.61	Do.
Celery, edible portion.....	Florida.....		2	.04	.60	.62	
do.....	do.....		30	3.49	1.18	2.32	Bladen soil.
Clover, alsike, above-ground portion.....	do.....		17	.58	.13	.46	Summary.
Clover, crimson; above-ground portion, full bloom.....	North Carolina.....	495	1	<i>Trifolium hybridum</i> .
Clover, crimson; above-ground portion.....	do.....		11	.80	.08	.39	Cecil sandy loam, <i>T. incarnatum</i> .
Clover, red; above-ground portion.....	Kentucky.....	14x	5	.24	.10	.17	Summary, <i>T. incarnatum</i> .
Clover, red; above-ground portion, mature with blossoms and seeds.....	New Hampshire.....	495	1	Tilist silt loam, <i>T. pratense</i> .
Clover, red; above-ground portion, in bloom.....	Pennsylvania.....	495	1	Gloucester stony loam, <i>T. pratense</i> .
Clover, red, above-ground portion.....	do.....		1	Hagerstown loam, <i>T. pratense</i> .
do.....	Rhode Island.....	19x	2	.10	.10	.10	Penn silt loam, <i>T. pratense</i> .
do.....	do.....		2	.10	.10	.10	Narragansett stony loam, <i>T. pratense</i> .
Clover, white; above-ground portion, some blossoms seeded.....	do.....		55	.85	.01	.21	Summary, <i>T. pratense</i> .
Clover, white; above-ground portion.....	Maire.....	495	1	<i>T. repens</i> .
do.....			34	.80	.07	.41	Summary, <i>T. repens</i> .

Coconut, kernel, without milk	Australia	39	1	.125	.001	.03	Summary.
Corn, grain	do	18	18	.06	.06	.048	Do.
Corn, sweet; edible portion	do	17	17	.05	.02	.03	Do.
Cotton, lint	do	3	3	.71	.01	.31	Do.
Cotton, seed	do	3	3				
Cowpeas, above-ground portion; full bloom	South Carolina	359	1			.09	
Cowpeas, above-ground portion; pods forming	do	359	1			.17	
Cowpeas, above-ground portion; pods formed	do	359	1			.03	
Cowpeas, above-ground portion; first pods beginning to turn yellow	Missouri	514	1			.11	
Cowpeas, above-ground portion	Ohio	193	1			.72	
Cowpeas, seeds	do	193	1			.30	
Horsebeans, forage	Germany	368	1			.19	<i>Vicia faba.</i>
Horsebeans, seed	do	368	1			.01	Do.
Horsebeans, straw	do	368	1			.30	Do.
Kaffr, grain	Ohio	193	1			.07	
Kale, edible portion	California	140	1			1.01	Yolo silt loam
Do	Maryland	140	2	.83	.83	.83	Sassaras sandy loam.
Do	Virginia	140	2	.77	.70	.73	Norfolk loam.
Do	England	600	6	1.01	.70	.85	Summary.
Kale, marrow stem; above-ground portion	do	600	4	.38	.16	.25	
Kale, thousand-head; above-ground portion	do	600	2	.16	.10	.13	
Kentucky bluegrass; above-ground portion, young blades	Kentucky	375	1			.24	
Kentucky bluegrass, ab-ve-ground portion, past bloom	Massachusetts	214	2	.14	.06	.10	
Kentucky bluegrass; above-ground portion, in seed-maturity unknown	Pennsylvania	495	12	.27	.01	.15	Penn silt loam.
Kentucky bluegrass; above-ground portion; stage of maturity unknown	Kentucky	375	1			.25	Summary.
Lettuce, edible portion	Germany	368	3	.65	.24	.42	
Lupine, straw only	do	368	1			.15	
Lupine, hulls only	do	368	1	.16	.10	.13	
Lupine, seeds	Ohio	193	1			.72	
Mangel-wurzel, roots	do	36	36	.23	.01	.09	Do.
Oats, grain	do	49	49	2.02	.06	.53	Do.
Oats, straw	do	49	49	.07	.03	.05	
Oats, above-ground portion, mature	Pennsylvania	466	5			.17	
Oats, above-ground portion, cut for hay	Germany	314, 368	9	.36	.01	.08	Do.
Onions, bulb	do	314, 368	2	.09	.07	.08	Do.
Onions, root	do	314, 368	2	.04	.01	.02	Do.
Parsnips, root	Germany	323	1			.03	Fort Collins loam; no fertilization.
Peas, seed	Colorado	257	5	.35	.01	.08	Fort Collins loam (N).
Potatoes, tubers	do	257	5	.38	.01	.10	Bladen soil (NPK).
Do	do	5x	6	.31	.17	.22	Bladen soil (NPKMn).
Do	Florida	5x	1			.20	Bladen soil (NPKCu).
Do	do	5x	1			.11	Coxville soil (NPK).
Do	do	5x	4	.23	.17	.16	Coxville soil (NPKMn).
Do	do	5x	3			.16	Coxville soil (NPKCu).
Do	do	5x	2			.22	

13 The summaries were calculated from data compiled from a number of sources.

TABLE 21.—The minor-element content of some important crops—Continued

SODIUM—Continued

Plant or part of plant	Location	Refer- ence No.	Analy- ses	Sodium in moisture- free material			Remarks
				Maxi- mum	Mini- mum	Mean	
				Percent	Percent	Percent	
Potatoes, tubers	Florida	5x	1				
Do	do	5x	4	0.23	0.18	Coxville soil (NPKCuMn).	
Do	do	5x	1		0.19	Leon soil (NPK).	
Do	New Hampshire	495	1		0.14	Norfolk soil (NPK).	
Do			188	38	01	Gloucester stony loam.	
Potatoes, tops	Germany	137, 368	16	1.18	0.13	Summary.	
Prunes, edible portion	France		13	0.28	0.04	Do.	
Rape, above-ground portion, flowering	do		3	0.41	0.36		
Rape, seed	do		1		0.01		
Redtop, above-ground portion, cut for hay	Kentucky	04	1		0.057	Dekalb silt loam; no fertilization.	
Do	do	14x	1		0.103	Dekalb silt loam (M).	
Do	do	14x	1		0.116	Dekalb silt loam (ML).	
Do	do	14x	1		0.044	Dekalb silt loam (LP).	
Do	do	14x	1		0.057	Tiltsit silt loam; no fertilization.	
Do	do	14x	1		0.116	Tiltsit silt loam (NP).	
Do	Rhode Island	19x	1		0.06	Bridgehampton very fine sandy loam (PK).	
Do	do	19x	1		0.03	Bridgehampton very fine sandy loam (NPK).	
Do	do		33	0.12	03	Summary.	
Rice, paddy	Hawaii	561	7	0.19	0.06	Do.	
Rice, brown	do		18	0.11	0.06	Do.	
Rice, polished	do	380	7	0.70	0.09	Do.	
Rice, straw	Hawaii	561	2	0.48	0.22	Do.	
Rutabaga, roots	do		4	0.08	0.01	Do.	
Rye, grain	do		31	0.27	0.04	Do.	
Rye, straw	do		1		0.27	Do.	
Sorghum, green stalk	Texas	240	1		0.57		
Sorghum, green head and stalk	do	240	1		0.25		
Soybean, leaves	France	335	1		0.01		
Soybean, pod	do	335	1		0.11		
Soybean, stems	do	335	1		0.05		
Soybean, hulls and vines	South Carolina	359	1		0.07		
Soybean, above-ground portion, full bloom	do	359	1		0.02		
Soybeans, seed, mature	Kentucky	14x	6	0.61	0.14	Do.	
Soybeans, above-ground portion, cut for hay	do	14x	1		0.24	Dekalb silt loam; no fertilization.	
Do	do	14x	1		0.34	Dekalb silt loam (ML).	
Do	do	14x	2	0.13	0.09	Dekalb silt loam (MP).	
Do	do	14x	2	0.10	0.04	Dekalb silt loam (MPL).	
Do	do	14x	2	0.10	0.04	Dekalb silt loam (MPKL).	

Do.	14x	1	05	Tiltsilt silt loam; no fertilization.
Do.	14x	1	13	Tiltsilt silt loam (M)
Do.	14x	2	06	Tiltsilt silt loam (MP)
Do.	14x	3	06	Tiltsilt silt loam (MPK)
Do.	14x	1	08	Tiltsilt silt loam (ML)
Do.	14x	2	07	Tiltsilt silt loam (MPL)
Do.	14x	2	12	Tiltsilt silt loam (MPKL)
Do.	14x	32	10	Summary.
Spinach, edible portion	140	2	2.37	Sassafras sandy loam.
Do.	140	2	1.62	Norfolk loam.
Do.	140	11	2.37	Summary.
Sugar beets, roots	606	8	4.76	Do.
Sugar beets, young leaves	606	5	4.81	Do.
Sugar beets, leaves, mature	584	14	6.65	Do.
Sugarcane, leaves	584	4	1.45	Do.
Sugarcane, canes	584	33	3.60	Do.
Sunflower, heads minus seeds and husks	592	1	06	Hagerstown loam, <i>Melilotus alba</i> .
Sunflower, seed husk	592	1	02	Summary. <i>M. alba</i> .
Sunflower, seed kernel without husk	592	1	07	Gloucester stony loam.
Sunflower, leaves	592	1	17	Hagerstown loam.
Sunflower, stalks	592	1	28	Summary.
Sunflower, above-ground portion, cut before bloom	368	1	1.00	Gloucester stony loam.
Sunflower, above-ground portion, full bloom	368	1	77	Hagerstown loam.
Sunflower, flower, cut for forage	368	3	12	Penn silt loam.
Sunflower, leaves and stems, cut for forage	368	3	08	Tiltsilt silt loam.
Sunflower, seeds with husks	368	1	07	Tiltsilt silt loam.
Sweetclover, above-ground portion	495	1	03	Bridgehampton very fine sandy loam.
Do.	495	4	07	Summary.
Sweetpotato, tuber	192	1	09	Summary.
Timothy, above-ground portion, in bloom	495	1	06	Parkwood fine sandy loam (NPK).
Do.	495	1	01	Summary.
Do.	495	1	02	Do.
Do.	495	1	01	Do.
Do.	278	2	81	Do.
Timothy, above-ground portion, in seed	495	1	35	Do.
Timothy, above-ground portion, cut for hay	14x	2	03	Do.
Do.	19x	19	41	Do.
Do.	141	32	3.47	Do.
Tobacco, burley; leaves	251	2	28	Do.
Tobacco, cigar wrapper type; leaves	141	2	14	Do.
Tobacco, dark fired type; leaves	62, 64	2	58	Do.
Tobacco, leaves, type unknown	5x	26	50	Do.
Tobacco, above-ground portion	5x	39	87	Do.
Tomato, fruit	5x	39	3.75	Do.
Do.	5x	3	1.81	Do.
Turnip, roots	368	1	53	Do.
Turnip, leaves	368	1	49	Do.
Vetch, above-ground portion, in bloom	368	1	53	Do.
Vetch, above-ground portion, seeds beginning to ripen	368	1	53	Do.
Vetch, above-ground portion, pods beginning to form	368	1	53	Do.
Vetch, straw, seeds ripening	368	1	11	Do.
Vetch, seeds	368	3	04	Do.
Walnuts, edible portion	235	5	15	Do.

TABLE 21.—The minor-element content of some important crops—Continued
SODIUM—Continued

Plant or part of plant	Location	Reference No.	Analyses	Sodium in moisture-free material			Remarks
				Maximum	Minimum	Mean	
Wheat, grain.....	Kentucky.....	14x	Number	Percent	Percent	Tilsit silt loam; no fertilization. Tilsit silt loam (M). Tilsit silt loam (ML). Tilsit silt loam (MPL). Summary. Tilsit silt loam; no fertilization. Tilsit silt loam (M). Tilsit silt loam (ML). Tilsit silt loam (MPL). Summary.	
Do.....	do.....	14x	1	.13	.13		
Do.....	do.....	14x	1	.12	.12		
Do.....	do.....	14x	1	.15	.15		
Do.....	do.....	14x	2	.12	.10		
Do.....	do.....	14x	100	.27	.01		
Do.....	do.....	14x	1	.19	.07		
Do.....	do.....	14x	1	.07	.07		
Do.....	do.....	14x	1	.20	.20		
Do.....	do.....	14x	2	.22	.08		
Do.....	do.....	14x	48	.30	.01	.14	
TITANIUM ¹⁴							
Plant or part of plant	Location	Reference No.	Titanium	Remarks			
Alfalfa, above-ground portion, cut for hay.....	France.....	67	Mg./kg.				
Almond, edible portion.....	do.....	67	3.6				
Apple, fruit.....	do.....	67	1.3				
Do.....	do.....	67	.9				
Barley, grain.....	Germany.....	204	4.0				
Beans, field; seed.....	France.....	67	.9				
Beets, roots.....	do.....	67	1.5				
Do.....	do.....	67	8.0	Phaseolus vulgaris.			
Beets, leaves.....	Germany.....	204	11.0				
Do.....	France.....	67	15.0				
Buckwheat, grain.....	Germany.....	204	42.0				
Buckwheat, straw.....	France.....	67	3.0				
Cabbage, leaves.....	do.....	67	6.6				
Do.....	do.....	67	1.5				
Cabbage, heart.....	do.....	67	1.2				
Cabbage, red.....	do.....	67	.9				
Carrots, roots.....	Germany.....	204	5.0				
Do.....	France.....	67	4.6				
Carrots, leaves.....	do.....	67	30.0				

TABLE 21.—The minor-element content of some important crops—Continued

ZINC 15

Plant or part of plant	Location	Refer- ence No.	Analyses	Zinc			Remarks
				Maximum Mg./kg.	Minimum Mg./kg.	Mean Mg./kg.	
Alfalfa, above-ground portion, full bloom.	France	58	Number	112	14	14	Summary (57, 373, 14x).
Alfalfa, above-ground portion, cut for hay	France	57	4	9	57	21	
Almond, edible portion.	do	57	2		6	3	
Apple, fruit	do	57	1		3	52	
Apricots, edible portion.	do	57	1			9	
Asparagus, edible portion.	do	57	1			21	
Banana.	do	57	1			35	
Barley, grain	do	57	1			8	
Beans, field; seed	do	57	10	56	23	23	
Beans, garden; edible pods and seeds.	France	57	1			8	
Beans, garden; edible hulls	Kentucky	379	1			53	
Beans, garden; immature seed	do	379	1			46	
Beans, lima; mature seed	do	379	1			46	
Beets, roots	France	57	2	69	25	47	
Buckwheat, grain	do	57	1			12	
Buckwheat, above-ground portion, cut in bloom	do	58	1			15	
Cabbage, leaves	do	56	2			24	
Carrots, roots	do	57	1	33	15	10	
Carrots, leaves	do	57	1			20	
Cauliflower, edible portion	do	57	1			25	
Cherries, edible portion	do	57	1			6	
Chestnut, edible portion	do	57	1			4	
Clover, alsike, above-ground portion.	Kentucky	14x	2	70	60	60	Trifolium hybridum. T. pratense.
Clover, red; above-ground portion	do	14x	5	60	30	44	
Do	do	14x	40	70	24	43	Do. Do.
Clover, red; seeds	do	14x	1			76	
Clover, red; foliage	do	14x	1			80	Do. Do.
Cocunut, edible portion	France	57	1			17	
Corn, grain	do	57	1			20	Tiltsilt silt loam.
Corn, stover	Kentucky	14x	2	13	9	11	
Do	do	14x	15	80	5	27	
Cotton, seed	Mississippi	374	1			320	84 101
Cress, water; edible portion	France	57	1			84	
Cress, garden; edible portion	do	56	1			44	99 4
Cucumber, fruit	do	57	1			44	
Dandelion, edible portion	do	56, 57	2	101	98	99	7
Dates, edible portion	do	57	1			4	
Endive, above-ground portion	do	57	1			7	

TABLE 21.—The minor-element content of some important crops—Continued
ZINC—Continued

Plant or part of plant	Location	Refer- ence No.	Analyses	Zinc			Remarks
				Maximum	Minimum	Mean	
Soybean, leaves	Kentucky	372	Number	Mg./kg.	Mg./kg.	Mg./kg.	
Soybean, seeds, mature	do	372	1	110	18	44	Dekalb silt loam; no fertilization.
Soybean, above-ground portion, cut for hay	do	14x	1	1	1	21	Dekalb silt loam (ML).
Do	do	14x	1	1	27	28	Dekalb silt loam (MP).
Do	do	14x	2	30	29	31	Dekalb silt loam (MPL).
Do	do	14x	2	34	32	36	Dekalb silt loam (MPKL).
Do	do	14x	1	40	34	50	Thist silt loam; no fertilization.
Do	do	14x	1	61	40	50	Thist silt loam (MP).
Do	do	14x	2	50	30	38	Thist silt loam (MPK).
Do	do	14x	3	50	30	32	Thist silt loam (ML).
Do	do	14x	1	40	33	36	Thist silt loam (MPL).
Do	do	14x	2	60	50	55	Thist silt loam (MPKL).
Do	do	14x	2	80	27	42	Summary.
Spinach	France	56, 57	31	121	67	94	
Strawberries, edible portion	do	57	1	1	4	19	<i>Melilotus alba</i> .
Sunflower, seeds	do	57	1	1	5	5	
Sweetclover, above-ground portion	Kentucky	14x	1	1	1	1	
Sweetpotatoes, tuber	France	57	1	1	1	1	
Timothy, above-ground portion, cut for hay	Kentucky	14x	4	60	40	60	Thist silt loam.
Do	do	14x	13	60	30	42	
Do	do	14x	2	67	12	39	Summary (57, 378).
Tomato, fruit	France	57	1	1	1	1	
Turnip, roots	do	57	1	1	1	1	
Turnip, leaves	do	57	1	1	1	1	
Vetch, seed	do	57	1	1	1	1	
Walnuts, edible portion	do	57	1	1	1	1	
Wheat, grain	Kentucky	14x	1	1	1	1	
Do	do	14x	1	1	1	1	
Do	do	14x	1	1	1	1	
Do	do	14x	2	90	70	80	Thist silt loam; no fertilization.
Do	do	14x	2	100	19	63	Thist silt loam (ML).
Wheat, straw	do	14x	27	100	19	20	Thist silt loam (MPL).
Do	Kentucky	14x	1	1	1	1	Summary (57, 67, 381, 543, 551, 14x).
Do	do	14x	1	1	1	1	Thist silt loam; no fertilization.
Do	do	14x	1	1	1	1	Thist silt loam (ML).
Do	do	14x	1	1	1	1	Thist silt loam; no fertilization.
Do	do	14x	2	20	20	20	Thist silt loam (ML).
Do	do	14x	2	25	7	17	Thist silt loam (MPL).
Do	do	14x	24	25	7	17	Summary (57, 381, 14x).

